

**Title:** Participation in physical activity decreased more in people with Rheumatoid Arthritis during the COVID-19 lockdown: a cross-sectional study

**Journal:** Rheumatology International

**Authors:** Christopher. Balchin<sup>1</sup>, Ai Lyn. Tan<sup>2, 3</sup>, Oliver. Wilson<sup>1</sup>, Jim. McKenna<sup>1</sup>, Antonios. Stavropoulos-Kalinoglou<sup>1</sup>

<sup>1</sup>Leeds Beckett University, Carnegie School of Sport, Leeds, UK

<sup>2</sup> University of Leeds, Leeds Institute of Rheumatic and Musculoskeletal Medicine, Chapel Allerton Hospital, Leeds, UK

<sup>3</sup> Leeds Teaching Hospitals NHS Trust, NIHR Leeds Biomedical Research Centre, Leeds, UK

**Corresponding author:** Dr Antonios Stavropoulos-Kalinoglou, Leeds Beckett University, Carnegie School of Sport, 225 Fairfax Hall, Headingley Campus, Churchwood Avenue, Leeds, LS6 3QS, [A.Stavropoulos@leedsbeckett.ac.uk](mailto:A.Stavropoulos@leedsbeckett.ac.uk)

**Online Resource 2:** Raw score to metric score conversion table for SWEMWBS. Transformed scores were subsequently used for parametric analyses

Raw Score (7-35)	Metric Score (7-35)
7	7.00
8	9.51
9	11.25
10	12.40
11	13.33
12	14.08
13	14.75
14	15.32
15	15.84
16	16.36
17	16.88
18	17.43
19	17.98
20	18.59
21	19.25
22	19.98
23	20.73
24	21.54
25	22.35
26	23.21
27	24.11
28	25.03
29	26.02
30	27.03
31	28.13
32	29.31
33	30.70
34	32.55
35	35.00