

**Title:** Participation in physical activity decreased more in people with Rheumatoid Arthritis during the COVID-19 lockdown: a cross-sectional study

**Journal:** Rheumatology International

**Authors:** Christopher. Balchin<sup>1</sup>, Ai Lyn. Tan<sup>2, 3</sup>, Oliver. Wilson<sup>1</sup>, Jim. McKenna<sup>1</sup>, Antonios. Stavropoulos-Kalinoglou<sup>1</sup>

<sup>1</sup>Leeds Beckett University, Carnegie School of Sport, Leeds, UK

<sup>2</sup> University of Leeds, Leeds Institute of Rheumatic and Musculoskeletal Medicine, Chapel Allerton Hospital, Leeds, UK

<sup>3</sup> Leeds Teaching Hospitals NHS Trust, NIHR Leeds Biomedical Research Centre, Leeds, UK

**Corresponding author:** Dr Antonios Stavropoulos-Kalinoglou, Leeds Beckett University, Carnegie School of Sport, 225 Fairfax Hall, Headingley Campus, Churchwood Avenue, Leeds, LS6 3QS, [A.Stavropoulos@leedsbeckett.ac.uk](mailto:A.Stavropoulos@leedsbeckett.ac.uk)

**Online Resource 4:** Mental wellbeing categories and % of RA and non-RA participants. Three-category versions of mental wellbeing were derived from transformed SWEMWBS scores: low, medium and high

Mental wellbeing categories	RA (n = 27)	Non-RA (n = 101)
Low (%)	44	28
Moderate	48	64
High	8	8