

A Comparison of Shear Wave Elastography between Normal Myometrium, Uterine Fibroids, and Adenomyosis: A Cross-Sectional Study

Sutee Pongpunprut, M.D.¹, Panyu Panburana, M.D.², Pornphan Wibulpolprasert, M.D.³, Wanwisa Waiyaput, M.Sc.⁴, Morakot Sroyraya, Ph.D.⁵, Tharintorn Chansoon, M.D.⁶, Areepan Sophonsritsuk, M.D., Ph.D.⁷

- 1. Department of Obstetrics and Gynaecology, Panyananthaphikkhu Chonprathan Medical Centre, Srinakharinwirot University, Nontaburi, Thailand
- 2. Foetal and Maternal Medicine Division, Department of Obstetrics and Gynaecology, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand
- 3. Division of Diagnostic Radiology, Department of Diagnostic and Therapeutic, Radiology, Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand
- 4. Office of Research Academic and Innovation, Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand
 5. Department of Anatomy, Faculty of Science, Mahidol University, Bangkok, Thailand
 - 6. Department of Pathology, Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand
- 7. Reproductive Endocrinology and Infertility Division, Department of Obstetrics and Gynaecology, Faculty of Medicine, Ramathidodi Hospital, Mahidol University, Bangkok, Thailand

Table S1: Comparison mean shear wave velocity by menopausal status

Normal myometrium	Premenopause	Postmenopause	P value
Control n (%) Mean SWV ± SD	4 (16) 3.04 ± 0.92	21 (84) 3.49 ± 1.10	0.390
Adenomyosis n (%) Mean SWV ± SD	20 (80) 4.91 ± 1.28	5 (20) 3.55 ± 1.76	0.061
Fibroid n (%) Mean SWV ± SD	19 (76) 4.58 ± 0.96	6 (24) 4.47 ± 1.53	0.843

The Student's t test was used to compare mean SWV between two groups. n; Sample number and SWV; Shear wave velocity.

Email: areepan.sop@mahidol.ac.th