Supplementary Table 1. Demographic and clinical comparisons between patients stratified according to the median Brief Resilience Scale (BRS) score (i.e., BRS \leq 2.3 vs. BRS > 2.3)

	Patients with BRS ≤ 2.3 (n = 25)	Patients with BRS > 2.3 (<i>n</i> = 25)	р
Age (years)	73.96 ± 8.66	70.00 ± 8.436	0.109
Sex (M/F)	15/10	20/5	0.067
Age at onset (years)	67.41 ± 8.62	63.47 ± 7.02	0.081
Disease duration (years)	6.55 ± 3.81	6.52 ± 4.17	0.976
MDS-UPDRS 3	25.48 ± 12.83	27.73 ± 11.61	0.521
HY stage	2.20 ± 0.71	2.17 ± 0.46	0.864
Consultation delay due to service shutdown (days)	98.71 ± 59.58	88.60 ± 57.59	0.547
COVID-19 related worries	7.44 ± 1.52	5.91 ± 2.82	0.018
COVID-19 related worries about health status	8.04 ± 1.51	6.56 ± 2.98	0.028
TSQ	2.70 ± 1.83	1.13 ± 0.96	0.001
EQ-5D	3.18 ± 2.20	1.52 ± 1.73	0.005
EQ-VAS	58.70 ± 17.89	70.86 ± 12.49	0.008

Values are expressed as mean ± standard deviation, unless othewise specified. *p < 0.05. BRS: Brief Resilience Scale, EQ-5D: EuroQoL-5 dimensions, EQ-VAS: EuroQoL-visual analogue scale, M: male, F: female, UPDRS: Unified Parkinson's Disease Rating Scale, HY: Hoehn and Yahr, TSQ: Trauma Screening Questionnaire.