## Supplementary Table

**Supplementary Table S1.** MoCA's mean subscores according to participants' group

|                           | Total sample<br>n = 227 | Healthy participants n=126 | Cardiovascular<br>disease n=101 |
|---------------------------|-------------------------|----------------------------|---------------------------------|
| Subscores                 |                         |                            |                                 |
| Visuoexecutive $M \pm SD$ | $4.2 \pm 0.9$           | $4.2 \pm 0.8$              | $4.2 \pm 1.0$                   |
| Naming $M \pm SD$         | $2.9 \pm .5$            | $2.9 \pm 0.4$              | $2.9 \pm 0.5$                   |
| Attention $M \pm SD$      | $5.7 \pm 0.7$           | $5.7 \pm 0.7$              | $5.7 \pm 0.6$                   |
| Langage $M \pm SD$        | $2.5 \pm 0.6$           | $2.6 \pm 0.6$              | $2.4 \pm 0.7$                   |
| Abstraction $M \pm SD$    | $1.7 \pm 0.5$           | $1.8 \pm 0.4$              | $1.7 \pm 0.6$                   |
| Memory $M \pm SD$         | $4.1 \pm 1.1$           | $4.2 \pm 1.0$              | $4.0 \pm 1.2$                   |
| Orientation $M \pm SD$    | $5.9 \pm 0.3$           | $5.9 \pm 0.3$              | $6.0 \pm 0.2$                   |