

**Supplemental Table S1. PLASMIC and French Scores for TTP and other TMA cohorts by age, 2006-2018**

		Age group (years)					
		18-39		40-59		≥ 60	
Risk score		TTP	Non-TTP TMA	TTP	Non-TTP TMA	TTP	Non-TTP TMA
PLASMIC	0-4	1/21 (4.8%)	10/18 (55.6%)	1/12 (8.3%)	4/12 (33.3%)	3/11 (27.3%)	7/14 (50.0%)
	5-7	20/21 (95.2%)	8/18 (44.4%)	11/12 (91.7%)	8/12 (66.7%)	8/11 (72.7%)	7/14 (50.0%)
French	0-1	7/21 (33.3%)	22/23 (95.7%)	5/12 (41.7%)	15/16 (93.8%)	6/11 (54.5%)	11/15 (73.3%)
	2	14/21 (66.7%)	1/23 (4.3%)	7/12 (58.3%)	1/16 (6.3%)	5/11 (45.5%)	4/15 (26.7%)

Abbreviations: TMA, thrombotic microangiopathy; TTP, thrombotic thrombocytopenic purpura.

Note: Data presented as number/total (%).

TTP diagnosis was confirmed by ADAMTS13 activity in patients who were diagnosed in 2006-2018.

For PLASMIC score, 1 point was given for each of the following: (1) platelet < 30 x 10<sup>9</sup>/L; (2) hemolysis [reticulocyte count > 2.5%, undetectable haptoglobin, or indirect bilirubin > 2 mg/dL]; (3) absence of active cancer; (4) absence of solid-organ or stem cell transplant; (5) MCV < 90 fL; (6) INR < 1.5; and (7) serum creatinine < 2.0 mg/dL.

For French score, 1 point was given for each of the following: (1) platelet < 30 x 10<sup>9</sup>/L and (2) serum creatinine ≤ 2.26 mg/dL.