

Appendix Figure 1. Flowchart for participants included in final analysis.



PATH, Population Assessment of Tobacco and Health Study.

**Appendix Figure 2.** Weighted prevalence of past-30-day cannabis use at follow-up waves by baseline e-cigarette use and mental health conditions.



IMH, internalizing mental health problems; EMH, externalizing mental health problems.

GAIN-SS Subscale <sup>a</sup>	Items
Internalizing mental health	Feeling very trapped, lonely, sad, blue, depressed, or hopeless about the future.
problems	Sleep trouble - such as bad dreams, sleeping restlessly or falling asleep during the day.
	Feeling very anxious, nervous, tense, scared, panicked or like something bad was going to happen.
	Becoming very distressed and upset when something reminded you of the past.
Externalizing mental health	Lied or conned to get things you wanted or to avoid having to do something.
problems	Had a hard time paying attention at school, work, or home.
	Had a hard time listening to instructions at school, work, or home.
	Were a bully or threatened other people.
	Started physical fights with other people.
	Felt restless or the need to run around or climb on things.
	Gave answers before the other person finished asking the question.

Appendix Table 1. Items for Internalizing and Externalizing Mental Health Problems

<sup>a</sup>GAIN-SS: Global Appraisal of Individual Needs – Short Screener.

Appendix Table 2. Percentage of Ever Cannabis Use at Each Follow-Up Wave by Baseline Covariates				
Covariates at corresponding baseline wave	Wave 1	Wave 2	Wave 3	
	% (95% CI)	% (95% CI)	% (95% CI)	
Overall ever cannabis use at follow-up waves	5.9 (5.3, 6.6)	5.7 (5.0, 6.3)	7.7 (7.1, 8.3)	
P30D e-cigarette use				
Yes	24.4 (10.5, 47.0)	19.9 (12.5, 30.3)	41.1 (31.7, 51.2)	
No	5.8 (5.2, 6.5)	5.5 (4.9, 6.2)	7.2 (6.6, 7.8)	
Internalizing mental health problems				
Low	3.9 (3.2, 4.8)	3.9 (3.3, 4.6)	5.9 (5.2, 6.8)	
Moderate	6.0 (4.9, 7.4)	6.2 (5.1, 7.6)	7.9 (6.6, 9.3)	
High	11.8 (9.6, 14.3)	10.0 (8.2, 12.1)	12.2 (10.8, 13.8)	
Externalizing mental health problems				
Low	3.1 (2.4, 4.1)	3.3 (2.7, 4.2)	5.1 (4.4, 6.0)	
Moderate	5.4 (4.3, 6.7)	5.4 (4.2, 6.8)	8.1 (6.8, 9.5)	
High	10.9 (9.3, 12.7)	9.6 (8.2, 11.3)	11.7 (10.5, 13.0)	
Age group, years				
12–14	5.8 (5.2, 6.6)	4.7 (4.1, 5.4)	5.6 (5.0, 6.2)	
15–17	8.1 (4.5, 14.1)	9.1 (7.6, 10.8)	11.6 (10.4, 12.9)	
Sex				
Male	5.5 (4.7, 6.5)	5.9 (5.1, 6.9)	7.2 (6.4, 8.0)	
Female	6.3 (5.4, 7.4)	5.4 (4.6, 6.3)	8.2 (7.3, 9.2)	
Race/ethnicity				
Non-Hispanic White	5.6 (4.8, 6.6)	6.0 (5.1, 7.1)	8.0 (7.1, 9.0)	
Non-Hispanic Black	7.8 (5.5, 10.9)	4.8 (3.4, 6.8)	8.5 (6.8, 10.5)	
Non-Hispanic other	2.7 (1.5, 4.9)	4.3 (2.5, 7.2)	5.3 (4.0, 7.1)	
Hispanic	6.8 (5.7, 8.3)	5.8 (4.8, 7.0)	8.0 (6.8, 9.4)	
Sexual orientation (ages $\geq 14$ years)				
Straight/heterosexual	8.4 (7.1, 10.1)	8.0 (7.0, 9.1)	10.1 (9.1, 11.3)	
Gay, lesbian, bisexual, or other	18.8 (9.4, 33.9)	14.6 (9.7, 21.3)	16.7 (13.1, 21.0)	
P30D combustible tobacco use <sup>a</sup>				
Yes	32.8 (19.1, 50.2)	28.1 (15.8, 44.8)	39.0 (26.6, 53.1)	
No	5.9 (5.2, 6.6)	5.5 (4.8, 6.2)	7.4 (6.9, 8.0)	
P30D alcohol use				

Yes	24.7 (15.8, 36.3)	17.5 (12.9, 23.2)	28.1 (23.1, 33.8)
No	5.6 (5.0, 6.3)	5.2 (4.6, 5.9)	6.8 (6.2, 7.3)
Parental education			
Less than high school	8.6 (7.0, 10.5)	5.1 (3.9, 6.6)	8.7 (7.4, 10.2)
High school graduate	7.4 (5.6, 9.6)	6.6 (5.1, 8.6)	7.3 (6.0, 8.9)
Some college or associate degree	5.7 (4.5, 7.2)	7.2 (6.0, 8.7)	8.8 (7.8, 10.0)
Bachelor's degree or higher	4.4 (3.5, 5.5)	4.4 (3.4, 5.7)	6.6 (5.6, 7.7)
State recreational cannabis law			
Legalized	9.1 (5.6, 14.4)	5.4 (3.2, 8.9)	8.2 (7.0, 9.6)
Not legalized	5.7 (5.1, 6.4)	5.7 (5.0, 6.4)	7.6 (6.9, 8.2)
	1 111 1 011		

<sup>a</sup>Combustible tobacco included cigarette, cigar, cigarillo, and filtered cigar.

P30D, past-30-day.

Appendix Table 3. AORs of Ever Cannabis Use During Follow-Up Among Baseline Cannabis-Naïve Adole					
	Model 1Model 2Model 3				
	No interaction	IMH interaction	EMH interaction		
Baseline characteristics	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)		
P30D e-cigarette use					
Yes	3.24 (2.09, 5.02)	6.56 (3.77, 11.43)	. , .		
No	ref	ref	ref		
Internalizing mental health problems (IMH)					
Low	ref	ref	ref		
Moderate	1.11 (0.93, 1.32)	1.13 (0.95, 1.34)	1.11 (0.93, 1.32)		
High	1.46 (1.18, 1.79)	1.53 (1.24, 1.89)	1.46 (1.18, 1.80)		
Externalizing mental health problems (EMH)					
Low	ref	ref	ref		
Moderate	1.55 (1.24, 1.93)	1.55 (1.24, 1.89)	1.58 (1.27, 1.98)		
High	2.27 (1.85, 2.78)	2.27 (1.85, 2.78)	2.31 (1.87, 2.85)		
P30D e-cigarette use X IMH					
Yes X High		0.21 (0.09, 0.48)			
Yes X Moderate		0.66 (0.27, 1.57)			
No X Low		ref			
P30D e-cigarette use X EMH					
Yes X High			0.49 (0.19, 1.26)		
Yes X Moderate			0.41 (0.11, 1.46)		
No X Low			ref		
Age group, years					
12–14	ref	ref	ref		
15–17	1.81 (1.56, 2.10)	1.81 (1.56, 2.10)	1.81 (1.56, 2.11)		
Sex					
Male	1.02 (0.88, 1.21)	1.01 (0.85, 1.20)	1.02 (0.88, 1.17)		
Female	ref	ref	ref		
Race/ethnicity					
Non-Hispanic White	ref	ref	ref		
Non-Hispanic Black	1.12 (0.89, 1.41)	1.12 (0.89, 1.41)	1.12 (0.89, 1.41)		
Non-Hispanic other	0.67 (0.52, 0.87)	0.67 (0.52, 0.87)	0.67 (0.51, 0.87)		

Hispanic	1.02 (0.86, 1.21)	1.01 (0.85, 1.20)	1.02 (0.86, 1.21)
P30D combustible tobacco use <sup>a</sup>			
Yes	2.97 (1.59, 5.54)	2.97 (1.64, 5.37)	2.90 (1.55, 5.41)
No	ref	ref	ref
P30D alcohol use			
Yes	3.25 (2.48, 4.27)	3.27 (2.49, 4.29)	3.27 (2.50, 4.29)
No	ref	ref	ref
Parental education			
Less than high school	ref	ref	ref
High school graduate	0.89 (0.72, 1.11)	0.90 (0.72, 1.12)	0.89 (0.72, 1.11)
Some college or associate degree	0.90 (0.73, 1.09)	0.89 (0.73, 1.09)	0.89 (0.73, 1.09)
Bachelor's degree or higher	0.65 (0.54, 0.79)	0.65 (0.54, 0.79)	0.65 (0.54, 0.79)
State recreational cannabis law			
Legalized	1.22 (0.96, 1.53)	1.23 (0.97, 1.54)	1.22 (0.97, 1.54)
Not legalized	ref	ref	ref

*Note*: Boldface indicates statistical significance (*p*<0.05).

<sup>a</sup>Combustible tobacco included cigarette, cigar, cigarillo, and filtered cigar.

P30D, past-30-day..

Baseline e-cigarette using status	Ever cannabis use for low internalizing problems	Ever cannabis use for moderate internalizing problems	Ever cannabis use for high internalizing problems
	<b>AOR</b> <sup>a</sup> (95% CI)	<b>AOR</b> <sup>a</sup> (95% CI)	<b>AOR</b> <sup>a</sup> (95% CI)
P30D e-cigarette use			
Yes	5.84 (3.09, 11.03)	5.25 (2.31, 11.92)	1.49 (0.73, 3.08)
No	ref	ref	ref

*Note*: Boldface indicates statistical significance (p < 0.05).

<sup>a</sup>Controlling for wave, sex, age, race/ethnicity, parental education, past-year externalizing mental health problems, P30D combustible tobacco use, P30D alcohol use, state-level cannabis legalization status at baseline.

P30D, past-30-day.

Appendix Table 5. AORs of P30D Cannabis Use at Follow-Up Among Baseline Cannabis-Naïve Adolescents			
	Model 1 Model 2		
	Without interaction	IMH interaction	EMH interaction
Baseline characteristics	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)
P30D e-cigarette use			
Yes	4.77 (2.92, 7.81)	14.58 (7.28, 29.19)	7.48 (2.95, 18.96)
No	ref	ref	ref
Internalizing mental health problems (IMH, continuous)	1.07 (0.99, 1.16)	1.10 (1.02, 1.19)	1.07 (0.99, 1.16)
Externalizing mental health problems (EMH, continuous)	1.13 (1.06, 1.20)	1.13 (1.06, 1.20)	1.14 (1.07, 1.22)
P30D e-cigarette use X IMH			
Yes <b>X</b> >0		0.60 (0.44, 0.82)	
No <b>X</b> 0		ref	
P30D e-cigarette use X EMH			
Yes <b>X</b> >0			0.88 (0.69, 1.12)
No <b>X</b> 0			ref
Age group, years			
12–14	ref	ref	ref
15–17	1.70 (1.35, 2.14)	1.67 (1.33, 2.11)	1.69 (1.34, 2.14)
Sex			
Male	0.91 (0.73, 1.14)	0.91 (0.73, 1.14)	0.92 (0.73, 1.14)
Female	ref	ref	ref
Race/ethnicity			
Non-Hispanic White	ref	ref	ref
Non-Hispanic Black	1.08 (0.74, 1.57)	1.07 (0.74, 1.55)	1.08 (0.74, 1.56)
Non-Hispanic other	0.52 (0.34, 0.81)	0.52 (0.34, 0.81)	0.52 (0.34, 0.81)
Hispanic	0.87 (0.66, 1.13)	0.85 (0.65, 1.10)	0.86 (0.66, 1.12)
P30D combustible tobacco use <sup>a</sup>			
Yes	2.19 (1.05, 4.58)	2.23 (1.09, 4.54)	2.16 (1.04, 4.50)
No	ref	ref	ref
P30D alcohol use			
Yes	3.35 (2.32, 4.84)	3.41 (2.36, 4.94)	3.37 (2.33, 4.86)
No	ref	ref	ref
Parental education			

Less than high school	ref	ref	ref
High school graduate	0.72 (0.49, 1.06)	0.73 (0.50, 1.06)	0.72 (0.49, 1.06)
Some college or associate degree	0.81 (0.63, 1.04)	0.79 (0.61, 1.01)	0.81 (0.63, 1.04)
Bachelor's degree or higher	0.59 (0.43, 0.82)	0.58 (0.42, 0.81)	0.59 (0.43, 0.82)
State recreational cannabis law			
Legalized	1.32 (0.95, 1.82)	1.35 (0.98, 1.86)	1.32 (0.96, 1.82)
Not legalized	ref	ref	ref

*Note*: Boldface indicates statistical significance (p < 0.05).

<sup>a</sup>Combustible tobacco included cigarette, cigar, cigarillo, and filtered cigar.

P30D, past-30-day.

Appendix Table 6. Percentage of P30D E-Cigarette Use at Baseline by Internalizing and Externalizing Mental Health Conditions

Covariates at corresponding baseline wave	Wave 1	Wave 2	Wave 3
	% (95% CI)	% (95% CI)	% (95% CI)
Internalizing mental health problems			
Low	0.4 (0.2, 0.7)	0.6 (0.3, 0.9)	1.3 (0.9, 1.7)
Moderate	0.5 (0.2, 1.2)	0.6 (0.4, 1.1)	0.9 (0.6, 1.4)
High	1.4 (0.8, 2.5)	1.9 (1.2, 3.1)	2.4 (1.7, 3.3)
Externalizing mental health problems			
Low	0.2 (0.1, 0.7)	0.5 (0.3, 0.9)	1.1 (0.8, 1.6)
Moderate	0.5 (0.2, 1.1)	0.6 (0.3, 1.1)	1.2 (0.7, 1.8)
High	1.2 (0.7, 1.9)	1.6 (1.0, 2.4)	2.1 (1.6, 2.1)