

Mechanisms, regulation and functions of the unfolded protein response

Claudio Hetz, Kezhong Zhang and Randal J. Kaufman

<https://doi.org/10.1038/s41580-020-0250-z>

Supplementary Table 1. IRE1 α binding partners and regulators.

A summary of the proteins that form complexes with IRE1. The experimental protocols and cell types used to detect the protein-protein interactions are indicated, in addition to conditions (endogenous proteins, overexpressed, tagged proteins). The complexes that were validated in vitro using purified proteins are also indicated.

Protein	Method	Endogenous	Ovexpressed IRE1	Ovexpressed interactor	In vitro binding	Cell type	References
		complex	(Tag: YES/NO)	(Tag: YES/NO)	recombinant/purified proteins		
cABL1	IP, PLA	-	YES	NO	Yes	INS-1, MEFs, HEK	1,52
BAK	PD/IP	YES	-	-	YES	MEF	2
BAX	PD/IP	YES	-	-	YES	MEF	2
PUMA	IPMS/PD/Y2H	YES	-	-	YES	MEF/HEK	3
BIM	IPMS/PD/Y2H	YES	-	-	YES	MEF/HEK	3
BI-1	IP/PD	YES	-	-	YES	MEF/HEK	4
BiD	PD/Y2H	-	NO	YES	YES	HEK	5
Caspases	not determined	-	-	-	-		6
CHIP	IPMS	YES	-	-	-	HEK	7
AIP1	IP	YES	-	-	-	EC	8
UFBP1	IP	-	YES	YES	-	HEK	9
ER protein targeting machineries	PAR/CLIP/IPMS	-	YES	-	-	HEK Trex	10
Filamin A	Y2H/IP/PD/CL	YES	-	-	YES	MEF/HEK/HUh7	11
Fortilin	PLA/IP/BLI	YES	-	-	YES	PC3	12
HSP47	IPMS/PLA/IP/PD/MST	YES	-	-	YES	MEF/HEK	13
HSP90	IP	YES	-	-	-	COS-7/INS-1	14, 15
HSP70	IP/PD	-	YES	NO	YES	PC12	16
BiP	IP/PD/MST	YES	-	-	YES	AT42J/HEK	17, 18
IP3R1-3	IP/PLA/PD	-	YES	NO	YES	MEF/HEK	19
JAB1	Y2H/IP/PD/CL	-	YES	YES	-	HEK	20
Myosin heavy chain IIB	IPMS/IP	-	YES	YES	-	HEK	21
NMI	MAPPIT-/P	-	NO	YES	-	HEK	22
Optineurin	IP	YES	-	-	-	MODE/HT29/Ileal crypts	23
PARP16	IP	-	YES	YES	-	HeLa	24
PDIA1	IP/PD	Yes	-	-	YES	HepG2	51
PDIA6	IP/MST	YES	-	-	YES	HEK	25, 26
PPM1I	IPMS	-	YES	YES	-	INS-1	27
PKA	Not tested	-	-	-	-	-	28
PKC	IP/PD	YES	-	-	YES	LO2	29
PSEN1	IP	-	YES	NO	-	SK-N-SH	30
PTP1B	Not tested	-	-	-	-	-	31
RACK1	IP	YES	-	-	-	INS-1/HEPG2	32
RIPK1	IP	-	YES	YES	-	HEK	33
RNF13	IP	-	YES	YES	-	HEK	34
RNH1	IPMS/PLA	YES	-	-	-	HK-2	35
RPAP2	IP	-	NO	YES	-	OVCAR-8	36
RTCB	IP	YES	-	-	YES	HEK	37-40
Sec61	IPMS/IP	-	YES	YES	YES	HEK	41, 42
SEL1L	IP	-	YES	NO	-	HEK	43
SIGMA-R1	IP	-	YES	YES	YES	CHO	44
HRD1	IP	YES	-	-	-	Synovial cells/HEK/HCT116	45
JIK	Y2H/IP	-	YES	YES	-	HEK	46
TRAF2-6	IP	YES	-	-	-	Primary macrophages/HEK	32
Ubiquitin D	MAPPIT	YES	-	-	-	HEK/EndoC- β H1 cells	47
USP14	Y2H/IP	YES	-	-	-	MEF/HEK/PC6.3	48, 49
YIP1	IP	YES	-	-	-	HeLa	50

IP: immunoprecipitation, PD: pull-down, IPMS: immunoprecipitation followed by mass spec analysis, Y2H: yeast two hybrid,

PLA: proximity ligation assay, CL: crosslinking, MAPPIT: mammalian protein-protein interaction trap

Supplementary references Table S1

1. Morita, S. et al. Targeting ABL-IRE1alpha Signaling Spares ER-Stressed Pancreatic beta Cells to Reverse Autoimmune Diabetes. *Cell Metab* **25**, 883-897 e8 (2017).
2. Hetz, C. et al. Proapoptotic BAX and BAK modulate the unfolded protein response by a direct interaction with IRE1alpha. *Science* **312**, 572-6 (2006).
3. Rodriguez, D.A. et al. BH3-only proteins are part of a regulatory network that control the sustained signalling of the unfolded protein response sensor IRE1alpha. *EMBO J* **31**, 2322-35 (2012).
4. Lisbona, F. et al. BAX inhibitor-1 is a negative regulator of the ER stress sensor IRE1alpha. *Molecular cell* **33**, 679-91 (2009).
5. Bashir, S. & Majid Fazili, K. Bid as a novel interacting partner of IRE1 differentially regulating its RNase activity. *BioRxiv* (2019).
6. Shemorry, A. et al. Caspase-mediated cleavage of IRE1 controls apoptotic cell commitment during endoplasmic reticulum stress. *Elife* **8** (2019).
7. Zhu, X. et al. Ubiquitination of inositol-requiring enzyme 1 (IRE1) by the E3 ligase CHIP mediates the IRE1/TRAF2/JNK pathway. *J Biol Chem* **289**, 30567-77 (2014).
8. Luo, D. et al. AIP1 is critical in transducing IRE1-mediated endoplasmic reticulum stress response. *J Biol Chem* **283**, 11905-12 (2008).
9. Liu, J. et al. A critical role of DDRGK1 in endoplasmic reticulum homoeostasis via regulation of IRE1alpha stability. *Nat Commun* **8**, 14186 (2017).
10. Acosta-Alvear, D. et al. The unfolded protein response and endoplasmic reticulum protein targeting machineries converge on the stress sensor IRE1. *Elife* **7** (2018).
11. Urra, H. et al. IRE1alpha governs cytoskeleton remodelling and cell migration through a direct interaction with filamin A. *Nat Cell Biol* **20**, 942-953 (2018).
12. Pinkaew, D. et al. Fortilin binds IRE1alpha and prevents ER stress from signaling apoptotic cell death. *Nat Commun* **8**, 18 (2017).
13. Sepulveda, D. et al. Interactome Screening Identifies the ER Luminal Chaperone Hsp47 as a Regulator of the Unfolded Protein Response Transducer IRE1alpha. *Mol Cell* **69**, 238-252 e7 (2018).
14. Marcu, M.G. et al. Heat shock protein 90 modulates the unfolded protein response by stabilizing IRE1alpha. *Mol Cell Biol* **22**, 8506-13 (2002).
15. Ota, A. & Wang, Y. Cdc37/Hsp90 protein-mediated regulation of IRE1alpha protein activity in endoplasmic reticulum stress response and insulin synthesis in INS-1 cells. *J Biol Chem* **287**, 6266-74 (2012).
16. Gupta, S. et al. HSP72 protects cells from ER stress-induced apoptosis via enhancement of IRE1alpha-XBP1 signaling through a physical interaction. *PLoS biology* **8**, e1000410 (2010).
17. Bertolotti, A., Zhang, Y., Hendershot, L.M., Harding, H.P. & Ron, D. Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. *Nat Cell Biol* **2**, 326-32 (2000).
18. Carrara, M., Prischi, F., Nowak, P.R., Kopp, M.C. & Ali, M.M. Noncanonical binding of BiP ATPase domain to Ire1 and Perk is dissociated by unfolded protein CH1 to initiate ER stress signaling. *Elife* **4** (2015).
19. Carreras-Sureda, A. et al. Non-canonical function of IRE1alpha determines mitochondria-associated endoplasmic reticulum composition to control calcium transfer and bioenergetics. *Nat Cell Biol* **21**, 755-767 (2019).
20. Oono, K. et al. JAB1 participates in unfolded protein responses by association and dissociation with IRE1. *Neurochem Int* **45**, 765-72 (2004).
21. He, Y. et al. Nonmuscle Myosin IIB Links Cytoskeleton to IRE1alpha Signaling during ER Stress. *Dev Cell* **23**, 1141-52 (2012).
22. Brozzi, F. et al. A combined "omics" approach identifies N-Myc interactor as a novel cytokine-induced regulator of IRE1 protein and c-Jun N-terminal kinase in pancreatic beta cells. *J Biol Chem* **289**, 20677-93 (2014).
23. Tschurtschenthaler, M. et al., Defective ATG16L1-mediated removal of IRE1alpha drives Crohn's disease-like ileitis. *J Exp Med* **214**:401–22 (2017) .
24. Jwa, M. & Chang, P. PARP16 is a tail-anchored endoplasmic reticulum protein required for the PERK- and IRE1alpha-mediated unfolded protein response. *Nat Cell Biol* **14**, 1223-30 (2012).
25. Eletto, D., Dersh, D., Gidalevitz, T. & Argon, Y. Protein disulfide isomerase A6 controls the decay of IRE1alpha signaling via disulfide-dependent association. *Mol Cell* **53**, 562-576 (2014).
26. Groenendyk, J. et al. Interplay between the oxidoreductase PDIA6 and microRNA-322 controls the response to disrupted endoplasmic reticulum calcium homeostasis. *Sci Signal* **7**, ra54 (2014).
27. Lu, G. et al. PPM1l encodes an inositol requiring-protein 1 (IRE1) specific phosphatase that regulates the functional outcome of the ER stress response. *Mol Metab* **2**, 405-16 (2013).
28. Mao, T. et al. PKA phosphorylation couples hepatic inositol-requiring enzyme 1alpha to glucagon signaling in glucose metabolism. *Proc Natl Acad Sci U S A* **108**, 15852-7 (2011).
29. Shin, G.C. et al. PRKCSH contributes to tumorigenesis by selective boosting of IRE1 signaling pathway. *Nat Commun* **10**, 3185 (2019).
30. Katayama, T. et al. Presenilin-1 mutations downregulate the signalling pathway of the unfolded-protein response. *Nat Cell Biol* **1**, 479-85 (1999).
31. Gu, F. et al. Protein-tyrosine phosphatase 1B potentiates IRE1 signaling during endoplasmic reticulum stress. *J Biol Chem* **279**, 49689-93 (2004).
32. Qiu, Y. et al. A crucial role for RACK1 in the regulation of glucose-stimulated IRE1alpha activation in pancreatic beta cells. *Science signaling* **3**, ra7 (2010).
33. Fidler, I.J. & Kripke, M.L. The challenge of targeting metastasis. *Cancer Metastasis Rev* **34**, 635-41 (2015).
34. Arshad, M. et al. RNF13, a RING finger protein, mediates endoplasmic reticulum stress-induced apoptosis through the inositol-requiring enzyme (IRE1alpha)/c-Jun NH₂-terminal kinase pathway. *J Biol Chem* **288**, 8726-36 (2013).
35. Tavernier, Q. et al. Regulation of IRE1 RNase activity by the Ribonuclease inhibitor 1 (RNH1). *Cell Cycle* **17**, 1901-1916 (2018).

36. Chang, T.K. et al. Coordination between Two Branches of the Unfolded Protein Response Determines Apoptotic Cell Fate. *Mol Cell* **71**, 629-636 e5 (2018).
37. Jurkin, J. et al. The mammalian tRNA ligase complex mediates splicing of XBP1 mRNA and controls antibody secretion in plasma cells. *EMBO J* **33**, 2922-36 (2014).
38. Kosmaczewski, S.G. et al. The RtcB RNA ligase is an essential component of the metazoan unfolded protein response. *EMBO Rep* **15**, 1278-85 (2014).
39. Lu, Y., Liang, F.X. & Wang, X. A synthetic biology approach identifies the mammalian UPR RNA ligase RtcB. *Mol Cell* **55**, 758-70 (2014).
40. Ray, A., Zhang, S., Rentas, C., Caldwell, K.A. & Caldwell, G.A. RTCB-1 mediates neuroprotection via XBP-1 mRNA splicing in the unfolded protein response pathway. *J Neurosci* **34**, 16076-85 (2014).
41. Plumb, R., Zhang, Z.R., Appathurai, S. & Mariappan, M. A functional link between the co-translational protein translocation pathway and the UPR. *Elife* **4** (2015).
42. Sundaram, A., Plumb, R., Appathurai, S. & Mariappan, M. The Sec61 translocon limits IRE1alpha signaling during the unfolded protein response. *Elife* **6** (2017).
43. Sun, S. et al. IRE1alpha is an endogenous substrate of endoplasmic-reticulum-associated degradation. *Nat Cell Biol* **17**, 1546-55 (2015).
44. Mori, T., Hayashi, T., Hayashi, E. & Su, T.P. Sigma-1 receptor chaperone at the ER-mitochondrion interface mediates the mitochondrion-ER-nucleus signaling for cellular survival. *PLoS One* **8**, e76941 (2013).
45. Gao, B. et al. Synoviolin promotes IRE1 ubiquitination and degradation in synovial fibroblasts from mice with collagen-induced arthritis. *EMBO Rep* **9**, 480-5 (2008).
46. Yoneda, T. et al. Activation of caspase-12, an endoplasmic reticulum (ER) resident caspase, through tumor necrosis factor receptor-associated factor 2-dependent mechanism in response to the ER stress. *J Biol Chem* **276**, 13935-40 (2001).
47. Brozzi, F. et al. Ubiquitin D Regulates IRE1alpha/c-Jun N-terminal Kinase (JNK) Protein-dependent Apoptosis in Pancreatic Beta Cells. *J Biol Chem* **291**, 12040-56 (2016).
48. Hyrskyluoto, A. et al. Ubiquitin-specific protease-14 reduces cellular aggregates and protects against mutant huntingtin-induced cell degeneration: involvement of the proteasome and ER stress-activated kinase IRE1alpha. *Hum Mol Genet* **23**, 5928-39 (2014).
49. Nagai, A. et al. USP14 inhibits ER-associated degradation via interaction with IRE1alpha. *Biochem Biophys Res Commun* **379**, 995-1000 (2009).
50. Taguchi, Y. et al. Yip1A, a novel host factor for the activation of the IRE1 pathway of the unfolded protein response during Brucella infection. *PLoS Pathog* **11**, e1004747 (2015).
51. Yu, J. et al. Phosphorylation switches protein disulfide isomerase activity to maintain proteostasis and attenuate ER stress. *EMBO J*, e103841, doi:10.15252/embj.2019103841 (2020).
52. Dufey E. et al. Genotoxic stress triggers the selective activation of IRE1 α -dependent RNA decay to modulate the DNA damage response. *Nature Com.* In press. doi: 10.1038/s41467-020-15694-y (2020).