

**Supplementary Material 3: Age-standardized rates per 100,000 inhabitants and proportions for ICD Chapter XVIII outcomes observed in 2020, and expected (mean of 2015-2019), for epidemiological weeks 10-48, their absolute difference, and risk ratio. Belo Horizonte MG, Brazil.**

	Observed	Expected	Absolute Difference	Risk Ratio
Deaths	33 (30;37)	39 (36;43)	-6.4	0.84 (0.73;0.96)*
Deaths at Home	21 (19;24)	25 (22;28)	-3.7	0.85 (0.72;1.01)
% Deaths at Home	64 (59;69)	63 (59;68)	1.0	1.02 (0.86;1.2)
Hosp. Admissions	100 (95;106)	98 (92;104)	2.4	1.02 (0.94;1.11)
ICU	20 (18;23)	19 (16;21)	1.7	1.09 (0.9;1.31)
% ICU	20 (18;22)	19 (17;21)	1.2	1.06 (0.88;1.28)
In-Hospital Deaths	7 (5;9)	8 (6;9)	-0.7	0.91 (0.67;1.23)
% In-Hospital Deaths	7 (5;8)	8 (6;9)	-0.9	0.89 (0.66;1.2)

ICD: International Classification of Diseases, ICU: intensive care unit