

SUPPLEMENTAL TABLES

Supplemental Table 1Plasma cytokine levels (pg/mL) by average levels of perceived stressor severity ($N = 92$)

	Interleukin-6	Interleukin-8	Interleukin-1 β	Tumor necrosis factor- α
Overall average perceived stressor severity				
Low tertile	0.651 (0.527-1.073)	2.652 (2.052-3.931)	0.065 (0.053-0.088)	1.560 (1.432-1.781)
Mid-tertile	0.681 (0.444-0.923)	2.662 (2.131-3.283)	0.079 (0.047-0.117)	1.611 (1.425-1.723)
High tertile	0.776 (0.586-1.035)	3.634 (2.588-4.521)	0.089 (0.073-0.105)	1.545 (1.300-1.953)
Chronic average perceived stressor severity				
Low tertile	0.605 (0.469-1.073)	2.611 (2.180-4.173)	0.070 (0.050-0.097)	1.551 (1.432-1.667)
Mid-tertile	0.712 (0.561-0.945)	2.700 (2.240-3.766)	0.094 (0.061-0.142)	1.545 (1.385-1.761)
High tertile	0.794 (0.578-1.278)	3.281 (2.745-4.050)	0.074 (0.058-0.086)	1.676 (1.458-1.953)
Acute average perceived stressor severity				
Low tertile	0.606 (0.527-1.303)	2.604 (1.903-3.748)	0.060 (0.042-0.075)	1.560 (1.329-1.781)
Mid-tertile	0.816 (0.563-0.984)	2.956 (2.247-3.745)	0.084 (0.053-0.137)	1.533 (1.425-1.976)
High tertile	0.687 (0.403-1.023)	3.455 (2.542-4.917)	0.093 (0.073-0.117)	1.584 (1.454-1.955)

Notes: Results are presented in pg/mL as median (interquartile range)