

Additional file 1. PICo Framework (Modified PICO).

<b>Population</b>	<b>TB/HIV high burden countries as per WHO for the period 2016 to 2020</b>
	<ul style="list-style-type: none"> <li>• Angola</li> <li>• Botswana</li> <li>• Brazil</li> <li>• Cameroon</li> <li>• The central African Republic</li> <li>• Chad</li> <li>• China</li> <li>• Congo</li> <li>• Democratic Republic of the Congo, DR Congo, DRC</li> <li>• Ethiopia</li> <li>• Ghana</li> <li>• Guinea-Bissau</li> <li>• India</li> <li>• Indonesia</li> <li>• Kenya</li> <li>• Lesotho</li> <li>• Liberia</li> <li>• Malawi</li> <li>• Mozambique</li> <li>• Myanmar</li> <li>• Namibia</li> <li>• Nigeria</li> <li>• Papua New Guinea, PNG</li> <li>• South Africa</li> <li>• Swaziland</li> <li>• Thailand</li> <li>• Uganda</li> <li>• United Republic of Tanzania, UR Tanzania, Tanzania</li> <li>• Zambia</li> <li>• Zimbabwe</li> </ul>
<b>Interest</b>	<b>Barriers and Facilitators</b> barrier*, facilitator*
<b>Context</b>	<b>Cotrimoxazole preventive therapy (CPT), Isoniazid preventive therapy (IPT)</b> Cotrimoxazol*, Trimethoprim, Sulfamethoxazole Drug Combination, Isoniazid

Note: The asterisk (\*) is used as a truncation indicator.