

Table SII. Additional outcomes

Assessment	Retrospective data	Visit 1 Lockdown		Visit 2 Partially opening		Visit 3 3-months Follow-up	
	Median (Q1–Q3)	Median (Q1–Q3)	<i>p</i> -value	Median (Q1–Q3)	<i>p</i> -value	Median (Q1–Q3)	<i>p</i> -value
Activities-specific Balance Confidence Scale	NA	76.88 (53.75-93.13)	0.146	86.09 (51.88-95.63)	0.491	79.38 (58.28-92.81)	<0.001
International Physical Activity Questionnaire							
Domestic and Garden MET-min/week	15 (0–360)	75 (0–675)	0.471	300 (15–540)	0.602	180 (0–1,380)	0.010
Leisure-Time MET-min/week	240 (0–858)	371.25 (0–804.38)	0.565	693 (140–1,453.50)	0.426	891 (66–1,386)	0.079
Transport MET-min/week	167.75 (0–798.75)	30 (0–359.55)	0.216	177 (0–297)	0.646	288 (60–648)	0.317
Work MET-min/week	0 (0–0)	0 (0–0)	0.136	0 (0–0)	0.820	0 (0–0)	0.978
Patient-Reported Outcomes Measurement Information System – 29 Version							
Anxiety, depression, sleep, and pain subscale	NA	29.5 (26.75–39)	0.937	32 (28–38.25)	0.422	33 (27.50–38.50)	0.957
Fatigue Severity Scale	36.5 (24–44)	35.5 (24.75–44.25)	0.734	37 (28–50)	0.596	37 (28–44)	0.578
National Institutes of Health Stroke Scale	2 (1–6)	NA	NA	2 (0–4)	0.110	2 (0–4)	0.004

MET: metabolic equivalent of tasks; NA: not applicable; Q: quartile. Bold font indicates *p*-values with a statistical significance of < 0.05.