

S3 Table. Rates (per 10,000 population) of RSV-associated hospitalization by congestive heart failure (CHF) status and by surveillance year, RSV-NET, 2015–2017 (N = 2042)

	Crude				Adjusted for under-detection ^a and age				Adjusted for under-detection ^a and age			
	CHF (95% CI)		no CHF (95% CI)		CHF (95% CI)		no CHF (95% CI)		Rate difference	95% CI	Rate ratio	95% CI
2015–2016												
All ages	5.1	4.3, 6.0	0.5	0.5, 0.5	24.5	20.4, 29.2	3.2	3.2, 3.2	21.3	17.2, 26.0	7.7	6.4, 9.2
<65 years	3.7	3.1, 4.5	0.3	0.3, 0.3	22.1	18.1, 26.8	1.7	1.7, 1.7	20.3	16.4, 25.0	12.7	10.4, 15.3
≥65 years	5.9	5.1, 6.8	1.7	1.7, 1.7	36.8	31.9, 42.3	10.6	10.4, 10.8	26.2	21.6, 31.5	3.5	3.1, 3.9
2016–2017												
All ages	7.5	6.4, 8.8	0.7	0.7, 0.7	28.4	23.7, 33.9	3.4	3.4, 3.4	25.0	20.3, 30.6	8.4	7.0, 10.0
<65 years	5.3	4.4, 6.5	0.3	0.3, 0.3	25.9	21.2, 31.4	1.6	1.6, 1.6	24.3	19.6, 29.8	15.8	13.0, 19.1
≥65 years	8.8	7.7, 10.2	2.5	2.4, 2.5	39.5	34.2, 45.4	11.0	10.8, 11.2	28.5	23.5, 34.2	3.6	3.2, 4.0

^a Adjusted for the under-detection of RSV (based on clinician testing practices and sensitivities of the laboratory tests).