

Supplementary Table S2

Characteristics of all patients receiving a bedaquiline (BDQ) containing treatment regimen against multidrug resistant tuberculosis in the Republic of Moldova (2016-2018). IQR= interquartile range, PTC=previously treated case

Patient characteristics	All patients receiving BDQ 2016-2018 % (n)	Patients included % (n)	Patients excluded % (n)	P-value
Urban	48.8 (99/203)	41.9 (26/62)	51.8 (73/141)	0.2240
Male	75.9 (154/203)	80.6 (50/62)	73.8 (104/141)	0.3736
Age, median (IQR)	37 (31-45)	39 (34-45)	36 (30-45)	0.0938
Microscopy +	72.4 (147/203)	74.2 (46/62)	71.6 (101/141)	0.7367
Case definition				
New case	25.1 (51/203)	29.0 (18/62)	23.4 (33/141)	0.4824
PTC-relapse	14.8 (30/203)	12.9 (8/62)	15.6 (22/141)	0.6741
PTC-default	13.8 (28/203)	17.7 (11/62)	12.1 (17/141)	0.2784
PTC-failure	46.3 (94/203)	40.3 (25/62)	48.9 (69/141)	0.2868
Treatment outcome				
still on treatment	6.9 (14/203)	6.5 (4/62)	7.1 (10/141)	1.0000
lost to follow-up	7.4 (15/203)	9.7 (6/62)	6.4 (9/141)	0.3975
cured	66.7 (116/174)	71.2 (37/52)	64.8 (79/122)	0.4838
failure	25.9 (45/174)	23.1 (12/52)	27.0 (33/122)	0.7059
died	7.5 (13/174)	5.8 (3/52)	8.2 (10/122)	0.7571
HIV status				
HIV +	12.7 (25/197)	12.9 (8/62)	12.6 (17/135)	1.0000
HIV status unknown	3.0 (6/203)	0.00 (0/62)	4.3 (6/141)	0.1804