

SUPPLEMENTARY ONLINE MATERIAL 1**Table 1:** Biomarkers included in the carbohydrate cluster

(1) Carbohydrates cluster		
Biomarker	Bio-fluid	Reference
Glucose	Plasma	Cosentino et al., 2020 ¹
HOMA-IR	-	Bloomgarden, 2003 ² Govers et al., 2015 ³ Shashaj et al., 2016 ⁴
Insulin	Serum	Aleksandrova et al., 2018 ⁵ Bloomgarden, 2003 ² Govers et al., 2015 ³
Leptin	Serum	Chen et al., 2014 ⁶ Finucane et al., 2019 ⁷ López-Jaramillo et al., 2014 ⁸
Adiponectin	Serum	Dastani et al., 2012 ⁹ Li et al., 2009 ¹⁰ Liu et al., 2016 ¹¹ Wang et al., 2018 ¹²
Lactate	Serum	Berhane et al., 2015 ¹³ Choi et al., 2002 ¹⁴ Lovejoy et al., 1992 ¹⁵ Shantha et al., 2013 ¹⁶
Glutamate	Serum	Martin and Price, 2018 ¹⁷ Ottosson et al., 2018 ¹⁸
Uric acid	Serum	Darmawan et al., 2018 ¹⁹ Fabbrini et al., 2014 ²⁰ Johnson et al., 2013 ²¹ van der Schaft et al., 2017 ²²
Propionylcarnitine	Plasma	Mai et al., 2013 ²³ Mihalik et al., 2010 ²⁴ Zhang et al., 2014 ²⁵
BCAA (Valine, Leucine, Isoleucine)	Serum	Chen et al., 2016 ²⁶ Katagiri et al., 2018 ²⁷ Lotta et al., 2016 ²⁸ Newgard et al., 2009 ²⁹ Okekunle et al., 2019 ³⁰
Phenylalanine	Serum	Chen et al., 2019 ³¹ Suzuki et al., 2019 ³² Wang et al., 2011 ³³
Tyrosine	Serum	Chen et al., 2019 ³¹ Hellmuth et al., 2016 ³⁴ Wang et al., 2011 ³³
Glutamine	Serum	Chen et al., 2019 ³¹ Guasch-Ferré et al., 2016 ³⁵ Liu et al., 2019 ³⁶ Rhee et al., 2018 ³⁷

Abbreviations: HOMA-IR, homeostatic model assessment of insulin resistance; BCAA, branched chain amino acids.

Table 2: Biomarkers included in the lipid cluster

(2) Lipid cluster		
Biomarker	Bio-fluid	Reference
LDL-cholesterol	Plasma	Grundy et al., 2019 ³⁸
HDL-cholesterol	Plasma	Grundy et al., 2019 ³⁸
Triglycerides	Plasma	Grundy et al., 2019 ³⁸
Total cholesterol	Plasma	Grundy et al., 2019 ³⁸
PUFAs (total)	Serum	Koga et al., 2019 ³⁹
LPCs (total)	Plasma	Law et al., 2019 ⁴⁰
Linoleic acid C18:2	Serum	Koga et al., 2019 ³⁹
Oleic acid C18:1	Serum	Griel and Kris-Etherton, 2006 ⁴¹ Yamagishi et al., 2013 ⁴²
Leptin	Serum	Mantzoros and Flier, 2000 ⁴³
Adiponectin	Serum	Abdella and Mojmíniyi, 2018 ⁴⁴ Liu et al., 2018 ⁴⁵
SFAs	Serum	Liu et al., 2019 ⁴⁶
3-hydroxybutyrate	Serum	Margolis and O'Fallon, 2020 ⁴⁷
MUFAs (total)	Serum	Griel and Kris-Etherton, 2006 ⁴¹ Yamagishi et al., 2013 ⁴²
Propionylcarnitine	Plasma	Ottosson et al., 2018 ¹⁸
DHA C20:3	Serum	Koga et al., 2019 ³⁹
Choline	Serum	Ottosson et al., 2018 ¹⁸

Abbreviations: LDL, low-density lipoprotein; HDL, high-density lipoprotein; PUFAs, poly unsaturated fatty acids; LPC, lysophosphatidylcholine; SFAs, saturated fatty acids; MUFAs, monounsaturated fatty acids; DHA, docosahexaenoic acid.

Table 3: Biomarkers included in the inflammation cluster

(3) Inflammation cluster		
Biomarker	Bio-fluid	Reference
CRP	Serum	Amsterdam, 2003 ⁴⁸ Knight, 2015 ⁴⁹ Liu et al., 2016 ¹¹ Musunuru et al., 2008 ⁵⁰ Wang et al., 2013 ⁵¹
IL-6	Plasma	Liu et al., 2016 ¹¹ Wang et al., 2013 ⁵¹
N-acetylglycoproteins	Serum	Gruppen et al., 2019 ⁵² Ritchie et al., 2015 ⁵³
TNF α	Plasma	Liu et al., 2016 ¹¹
MCP-1	Plasma	Georgakis et al., 2019 ⁵⁴
IL-10	Plasma	Charles et al., 2011 ⁵⁵ Leon-Cabrera et al., 2015 ⁵⁶ Meng et al., 2019 ⁵⁷
sICAM1	Plasma	El Amine et al., 2010 ⁵⁸ Luc et al., 2003 ⁵⁹ Straczkowski et al., 2002 ⁶⁰
LBP	Plasma	Moreno-Navarrete et al., 2012 ⁶¹
sCD14	Plasma	de Courten et al., 2016 ⁶²
LPCs (total)	Plasma	Iwase et al., 2008 ⁶³
Linoleic acid C18:2	Serum	Steffen et al., 2012 ⁶⁴ Yli-Jama et al., 2002 ⁶⁵
DHA C20:3	Serum	Steffen et al., 2012 ⁶⁴ Yli-Jama et al., 2002 ⁶⁵

Abbreviations: CRP, C-reactive protein; IL-6, interleukin-6; IL-10, interleukin-10; TNF α , tumor necrosis factor alpha; MCP-1, monocyte chemoattractant protein-1; sICAM-1, soluble intercellular adhesion molecule-1; LBP, lipopolysaccharide binding protein; sCD14, soluble CD14; LPC, lysophosphatidylcholines; DHA: docosahexaenoic acid.

Table 4: Biomarkers included in the oxidative stress cluster

(4) Oxidative stress cluster		
Biomarker	Bio-fluid	Reference
8-iso-PGF2α	Urine	Davies and Roberts, 2011 ⁶⁶ Kim et al., 2012 ⁶⁷ Milne et al., 2015 ⁶⁸ van't Erve, 2018 ⁶⁹
8-OHdG	Urine	Di Minno et al., 2016 ⁷⁰ Kroese and Scheffe, 2014 ⁷¹
LDLox	Plasma	Barbosa et al., 2011 ⁷² Gao et al., 2017 ⁷³
Uric acid	Serum	Darmawan et al., 2018 ¹⁹ Fabbrini et al., 2014 ²⁰ van der Schaft et al., 2017 ²²
Allantoin	Urine	Il'yasova et al., 2012 ⁷⁴
Betaine	Urine	Svingen et al., 2016 ⁷⁵ Walford et al., 2016 ⁷⁶
Pseudouridine	Urine	Topp et al., 2008 ⁷⁷
Dimethylglycine	Urine	Svingen et al., 2016 ⁷⁵
Glycine	Serum	Sekhar et al., 2011 ⁷⁸
Methionine	Serum	Grizales et al., 2018 ⁷⁹

Abbreviations: 8-iso-PGF2α, 8-iso-prostaglandin F2 alpha; 8-OHdG, 8-hydroxydeoxyguanosine; LDLox, oxidized low density lipoprotein.

Table 5: Biomarkers included in the microbiota cluster

(5) Microbiota cluster		
Biomarker	Bio-fluid	Reference
TMAO	Serum	Bain et al., 2006 ⁸⁰ Chen et al., 2016 ⁸¹ Ge et al., 2020 ⁸² Heianza et al., 2017 ⁸³ Schiattarella et al., 2017 ⁸⁴ Yang et al., 2019 ⁸⁵ Yao et al., 2020 ⁸⁶
TMA	Urine	Aragonès et al., 2019 ⁸⁷ Bouatra et al., 2013 ⁸⁸ Chen et al., 2016 ⁸¹ Palau-Rodriguez et al., 2015 ⁸⁹ Yang et al., 2019 ⁸⁵
DMA	Urine	Bouatra et al., 2013 ⁸⁸ Palau-Rodriguez et al., 2015 ⁸⁹
Betaine	Urine	Chen et al., 2016 ⁸¹ Palau-Rodriguez et al., 2015 ⁸⁹ Yang et al., 2019 ⁸⁵
Choline	Serum	Aragonès et al., 2019 ⁸⁷ Chen et al., 2016 ⁸¹ Palau-Rodriguez et al., 2015 ⁸⁹ Yang et al., 2019 ⁸⁵
Dimethylglycine	Urine	Palau-Rodriguez et al., 2015 ⁸⁹
Succinate	Plasma	Psychogios et al., 2011 ⁹⁰ Serena et al., 2018 ⁹¹
Lactate	Serum	Psychogios et al., 2011 ⁹⁰ Vojinovic et al., 2019 ⁹²
LBP	Plasma	Awoyemi et al., 2018 ⁹³ Barengolts et al., 2019 ⁹⁴ Bouatra et al., 2013 ⁸⁸ Citronberg et al., 2018 ⁹⁵
Acetate	Urine	Liu et al., 2014 ⁹⁶

Abbreviations: TMAO, trimethylamine N-oxide; TMA, trimethylamine; DMA, dimethylamine; LBP, lipopolysaccharide binding protein.

References:

1. Cosentino F, Grant PJ, Aboyans V, et al. 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J* 2020;41(2):255-323.
2. Bloomgarden ZT. American Association of Clinical Endocrinologists (AACE) consensus conference on the insulin resistance syndrome: 25-26 August 2002, Washington, DC. *Diabetes Care* 2003;26(4):1297-303.
3. Govers E, Slof E, Verkoelen H, et al. Guideline for the Management of Insulin Resistance. *Int J Endocrinol Metab Disord* 2015;1(4).
4. Shashaj B, Luciano R, Contoli B, et al. Reference ranges of HOMA-IR in normal-weight and obese young Caucasians. *Acta Diabetol* 2016;53(2):251-60.
5. Aleksandrova K, Mozaffarian D, Pischedda T. Addressing the Perfect Storm: Biomarkers in Obesity and Pathophysiology of Cardiometabolic Risk. *Clin Chem* 2018;64(1):142-53.
6. Chen GC, Qin LQ, Ye JK. Leptin levels and risk of type 2 diabetes: gender-specific meta-analysis. *Obes Rev* 2014;15(2):134-42.
7. Finucane FM, Luan J, Wareham NJ, et al. Correlation of the leptin:adiponectin ratio with measures of insulin resistance in non-diabetic individuals. *Diabetologia* 2009;52(11):2345-49.
8. López-Jaramillo P, Gómez-Arbeláez D, López-López J, et al. The role of leptin/adiponectin ratio in metabolic syndrome and diabetes. *Horm Mol Biol Clin Investig* 2014;18(1):37-45.
9. Dastani Z, Hivert MF, Timpson N, et al. Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS Genet* 2012;8(3):e1002607.
10. Li S, Shin HJ, Ding EL, et al. Adiponectin levels and risk of type 2 diabetes: a systematic review and meta-analysis. *Jama* 2009;302(2):179-88.
11. Liu C, Feng X, Li Q, et al. Adiponectin, TNF- α and inflammatory cytokines and risk of type 2 diabetes: A systematic review and meta-analysis. *Cytokine* 2016;86:100-09.
12. Wang Y, Meng RW, Kunutsor SK, et al. Plasma adiponectin levels and type 2 diabetes risk: a nested case-control study in a Chinese population and an updated meta-analysis. *Sci Rep* 2018;8(1):406.
13. Berhane F, Fite A, Daboul N, et al. Plasma Lactate Levels Increase during Hyperinsulinemic Euglycemic Clamp and Oral Glucose Tolerance Test. *J Diabetes Res* 2015;2015:102054.
14. Choi CS, Kim YB, Lee FN, et al. Lactate induces insulin resistance in skeletal muscle by suppressing glycolysis and impairing insulin signaling. *Am J Physiol Endocrinol Metab* 2002;283(2):E233-40.
15. Lovejoy J, Newby FD, Gebhart SSP, et al. Insulin resistance in obesity is associated with elevated basal lactate levels and diminished lactate appearance following intravenous glucose and insulin. *Metabolism* 1992;41(1):22-27.
16. Shantha GP, Wasserman B, Astor BC, et al. Association of blood lactate with carotid atherosclerosis: the Atherosclerosis Risk in Communities (ARIC) Carotid MRI Study. *Atherosclerosis* 2013;228(1):249-55.
17. Martin AJ, Price CI. A Systematic Review and Meta-Analysis of Molecular Biomarkers Associated with Early Neurological Deterioration Following Acute Stroke. *Cerebrovasc Dis* 2018;46(5-6):230-41.
18. Ottosson F, Brunkwall L, Ericson U, et al. Connection Between BMI-Related Plasma Metabolite Profile and Gut Microbiota. *J Clin Endocrinol Metab* 2018;103(4):1491-501.
19. Darmawan G, Hamijoyo L, Hasan I. Association between Serum Uric Acid and Non-Alcoholic Fatty Liver Disease: A Meta-Analysis. *Acta Med Indones* 2017;49(2):136-47.
20. Fabbriani E, Serafini M, Colic Baric I, et al. Effect of plasma uric acid on antioxidant capacity, oxidative stress, and insulin sensitivity in obese subjects. *Diabetes* 2014;63(3):976-81.
21. Johnson RJ, Nakagawa T, Sanchez-Lozada LG, et al. Sugar, uric acid, and the etiology of diabetes and obesity. *Diabetes* 2013;62(10):3307-15.
22. van der Schaft N, Brahimaj A, Wen KX, et al. The association between serum uric acid and the incidence of prediabetes and type 2 diabetes mellitus: The Rotterdam Study. *PLoS One* 2017;12(6):e0179482.
23. Mai M, Tönjes A, Kovacs P, et al. Serum levels of acylcarnitines are altered in prediabetic conditions. *PLoS One* 2013;8(12):e82459.
24. Mihalik SJ, Goodpaster BH, Kelley DE, et al. Increased levels of plasma acylcarnitines in obesity and type 2 diabetes and identification of a marker of glucolipotoxicity. *Obesity (Silver Spring)* 2010;18(9):1695-700.
25. Zhang X, Zhang C, Chen L, et al. Human serum acylcarnitine profiles in different glucose tolerance states. *Diabetes Res Clin Pract* 2014;104(3):376-82.
26. Chen T, Ni Y, Ma X, et al. Branched-chain and aromatic amino acid profiles and diabetes risk in Chinese populations. *Sci Rep* 2016;6:20594.
27. Katagiri R, Goto A, Budhathoki S, et al. Association between plasma concentrations of branched-chain amino acids and adipokines in Japanese adults without diabetes. *Sci Rep* 2018;8(1):1043.

28. Lotta LA, Scott RA, Sharp SJ, et al. Genetic Predisposition to an Impaired Metabolism of the Branched-Chain Amino Acids and Risk of Type 2 Diabetes: A Mendelian Randomisation Analysis. *PLoS Med* 2016;13(11):e1002179.
29. Newgard CB, An J, Bain JR, et al. A branched-chain amino acid-related metabolic signature that differentiates obese and lean humans and contributes to insulin resistance. *Cell Metab* 2009;9(4):311-26.
30. Okekunle AP, Zhang M, Wang Z, et al. Dietary branched-chain amino acids intake exhibited a different relationship with type 2 diabetes and obesity risk: a meta-analysis. *Acta Diabetol* 2019;56(2):187-95.
31. Chen S, Akter S, Kuwahara K, et al. Serum amino acid profiles and risk of type 2 diabetes among Japanese adults in the Hitachi Health Study. *Sci Rep* 2019;9(1):7010.
32. Suzuki Y, Kido J, Matsumoto S, et al. Associations among amino acid, lipid, and glucose metabolic profiles in childhood obesity. *BMC Pediatr* 2019;19(1):273.
33. Wang TJ, Larson MG, Vasan RS, et al. Metabolite profiles and the risk of developing diabetes. *Nat Med* 2011;17(4):448-53.
34. Hellmuth C, Kirchberg FF, Lass N, et al. Tyrosine Is Associated with Insulin Resistance in Longitudinal Metabolomic Profiling of Obese Children. *J Diabetes Res* 2016;2016:2108909.
35. Guasch-Ferré M, Hruby A, Toledo E, et al. Metabolomics in Prediabetes and Diabetes: A Systematic Review and Meta-analysis. *Diabetes Care* 2016;39(5):833-46.
36. Liu X, Zheng Y, Guasch-Ferré M, et al. High plasma glutamate and low glutamine-to-glutamate ratio are associated with type 2 diabetes: Case-cohort study within the PREDIMED trial. *Nutr Metab Cardiovasc Dis* 2019;29(10):1040-49.
37. Rhee SY, Jung ES, Park HM, et al. Plasma glutamine and glutamic acid are potential biomarkers for predicting diabetic retinopathy. *Metabolomics* 2018;14(7):89.
38. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2019;73(24):e285-e350.
39. Koga N, Ogura J, Yoshida F, et al. Altered polyunsaturated fatty acid levels in relation to proinflammatory cytokines, fatty acid desaturase genotype, and diet in bipolar disorder. *Transl Psychiatry* 2019;9(1):208.
40. Law SH, Chan ML, Marathe GK, et al. An Updated Review of Lysophosphatidylcholine Metabolism in Human Diseases. *Int J Mol Sci* 2019;20(5).
41. Griel AE, Kris-Etherton PM. Tree nuts and the lipid profile: a review of clinical studies. *Br J Nutr* 2006;96 Suppl 2:S68-78.
42. Yamagishi K, Folsom AR, Steffen LM. Plasma fatty acid composition and incident ischemic stroke in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) Study. *Cerebrovasc Dis* 2013;36(1):38-46.
43. Mantzoros CS, Flier JS. Editorial: leptin as a therapeutic agent--trials and tribulations. *J Clin Endocrinol Metab* 2000;85(11):4000-2.
44. Abdella NA, Mojiminiyi OA. Clinical Applications of Adiponectin Measurements in Type 2 Diabetes Mellitus: Screening, Diagnosis, and Marker of Diabetes Control. *Dis Markers* 2018;2018:5187940.
45. Liu Z, Liang S, Que S, et al. Meta-Analysis of Adiponectin as a Biomarker for the Detection of Metabolic Syndrome. *Front Physiol* 2018;9:1238.
46. Liu Q, Matthan NR, Manson JE, et al. Plasma Phospholipid Fatty Acids and Coronary Heart Disease Risk: A Matched Case-Control Study within the Women's Health Initiative Observational Study. *Nutrients* 2019;11(7).
47. Margolis LM, O'Fallon KS. Utility of Ketone Supplementation to Enhance Physical Performance: A Systematic Review. *Adv Nutr* 2020;11(2):412-19.
48. Amsterdam EA. C-reactive protein: a guideline for its application. *Prev Cardiol* 2003;6(2):70.
49. Knight ML. The Application of High-Sensitivity C-Reactive Protein in Clinical Practice: A 2015 Update. *US Pharm* 2015;40(2):50-53.
50. Musunuru K, Krai BG, Blumenthal RS, et al. The use of high-sensitivity assays for C-reactive protein in clinical practice. *Nat Clin Pract Cardiovasc Med* 2008;5(10):621-35.
51. Wang X, Bao W, Liu J, et al. Inflammatory markers and risk of type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care* 2013;36(1):166-75.
52. Gruppen EG, Kunutsor SK, Kienecker LM, et al. GlycA, a novel pro-inflammatory glycoprotein biomarker is associated with mortality: results from the PREVEND study and meta-analysis. *J Intern Med* 2019;286(5):596-609.
53. Ritchie SC, Würtz P, Nath AP, et al. The Biomarker GlycA Is Associated with Chronic Inflammation and Predicts Long-Term Risk of Severe Infection. *Cell Syst* 2015;1(4):293-301.

54. Georgakis MK, Malik R, Björkbacka H, et al. Circulating Monocyte Chemoattractant Protein-1 and Risk of Stroke: Meta-Analysis of Population-Based Studies Involving 17 180 Individuals. *Circ Res* 2019;125(8):773-82.
55. Charles BA, Doumatey A, Huang H, et al. The roles of IL-6, IL-10, and IL-1RA in obesity and insulin resistance in African-Americans. *J Clin Endocrinol Metab* 2011;96(12):E2018-22.
56. Leon-Cabrera S, Arana-Lechuga Y, Esqueda-León E, et al. Reduced systemic levels of IL-10 are associated with the severity of obstructive sleep apnea and insulin resistance in morbidly obese humans. *Mediators Inflamm* 2015;2015:493409.
57. Meng D, Liang L, Guo X. Serum interleukin-10 level in patients with inflammatory bowel disease: A meta-analysis. *Eur J Inflamm* 2019;17:1-7.
58. El Amine M, Sohawon S, Lagneau L, et al. Plasma levels of ICAM-1 and circulating endothelial cells are elevated in unstable types 1 and 2 diabetes. *Endocr Regul* 2010;44(1):17-24.
59. Luc G, Arveiler D, Evans A, et al. Circulating soluble adhesion molecules ICAM-1 and VCAM-1 and incident coronary heart disease: the PRIME Study. *Atherosclerosis* 2003;170(1):169-76.
60. Straczkowski M, Lewczuk P, Dzienis-Straczkowska S, et al. Elevated soluble intercellular adhesion molecule-1 levels in obesity: relationship to insulin resistance and tumor necrosis factor-alpha system activity. *Metabolism* 2002;51(1):75-8.
61. Moreno-Navarrete JM, Ortega F, Serino M, et al. Circulating lipopolysaccharide-binding protein (LBP) as a marker of obesity-related insulin resistance. *Int J Obes (Lond)* 2012;36(11):1442-9.
62. de Courten B, Moreno-Navarrete JM, Lyons J, et al. Contrasting association of circulating sCD14 with insulin sensitivity in non-obese and morbidly obese subjects. *Mol Nutr Food Res* 2016;60(1):103-9.
63. Iwase M, Sonoki K, Sasaki N, et al. Lysophosphatidylcholine contents in plasma LDL in patients with type 2 diabetes mellitus: relation with lipoprotein-associated phospholipase A2 and effects of simvastatin treatment. *Atherosclerosis* 2008;196(2):931-6.
64. Steffen BT, Steffen LM, Tracy R, et al. Obesity modifies the association between plasma phospholipid polyunsaturated fatty acids and markers of inflammation: the Multi-Ethnic Study of Atherosclerosis. *Int J Obes (Lond)* 2012;36(6):797-804.
65. Yli-Jama P, Seljeflot I, Meyer HE, et al. Serum non-esterified very long-chain PUFA are associated with markers of endothelial dysfunction. *Atherosclerosis* 2002;164(2):275-81.
66. Davies SS, Roberts LJ, 2nd. F2-isoprostanes as an indicator and risk factor for coronary heart disease. *Free Radic Biol Med* 2011;50(5):559-66.
67. Kim JY, Lee JW, Youn YJ, et al. Urinary levels of 8-iso-prostaglandin f2 α and 8-hydroxydeoxyguanine as markers of oxidative stress in patients with coronary artery disease. *Korean Circ J* 2012;42(9):614-7.
68. Milne GL, Dai Q, Roberts LJ, 2nd. The isoprostanes--25 years later. *Biochim Biophys Acta* 2015;1851(4):433-45.
69. van 't Erve TJ. Strategies to decrease oxidative stress biomarker levels in human medical conditions: A meta-analysis on 8-iso-prostaglandin F(2 α). *Redox Biol* 2018;17:284-96.
70. Di Minno A, Turnu L, Porro B, et al. 8-Hydroxy-2'-Deoxyguanosine Levels and Cardiovascular Disease: A Systematic Review and Meta-Analysis of the Literature. *Antioxid Redox Signal* 2016;24(10):548-55.
71. Kroese LJ, Scheffer PG. 8-hydroxy-2'-deoxyguanosine and cardiovascular disease: a systematic review. *Curr Atheroscler Rep* 2014;16(11):452.
72. Barbosa KB, Volp AC, Hermsdorff HH, et al. Relationship of oxidized low density lipoprotein with lipid profile and oxidative stress markers in healthy young adults: a translational study. *Lipids Health Dis* 2011;10:61.
73. Gao S, Zhao D, Wang M, et al. Association Between Circulating Oxidized LDL and Atherosclerotic Cardiovascular Disease: A Meta-analysis of Observational Studies. *Can J Cardiol* 2017;33(12):1624-32.
74. Il'yasova D, Scarborough P, Spasojevic I. Urinary biomarkers of oxidative status. *Clin Chim Acta* 2012;413(19-20):1446-53.
75. Svingen GF, Schartum-Hansen H, Pedersen ER, et al. Prospective Associations of Systemic and Urinary Choline Metabolites with Incident Type 2 Diabetes. *Clin Chem* 2016;62(5):755-65.
76. Walford GA, Ma Y, Clish C, et al. Metabolite Profiles of Diabetes Incidence and Intervention Response in the Diabetes Prevention Program. *Diabetes* 2016;65(5):1424-33.
77. Topp H, Fusch G, Schöch G, et al. Noninvasive markers of oxidative DNA stress, RNA degradation and protein degradation are differentially correlated with resting metabolic rate and energy intake in children and adolescents. *Pediatr Res* 2008;64(3):246-50.
78. Sekhar RV, McKay SV, Patel SG, et al. Glutathione synthesis is diminished in patients with uncontrolled diabetes and restored by dietary supplementation with cysteine and glycine. *Diabetes Care* 2011;34(1):162-7.
79. Grizales AM, Patti ME, Lin AP, et al. Metabolic Effects of Betaine: A Randomized Clinical Trial of Betaine Supplementation in Prediabetes. *J Clin Endocrinol Metab* 2018;103(8):3038-49.

80. Bain MA, Faull R, Fornasini G, et al. Accumulation of trimethylamine and trimethylamine-N-oxide in end-stage renal disease patients undergoing haemodialysis. *Nephrol Dial Transplant* 2006;21(5):1300-4.
81. Chen YM, Liu Y, Zhou RF, et al. Associations of gut-flora-dependent metabolite trimethylamine-N-oxide, betaine and choline with non-alcoholic fatty liver disease in adults. *Sci Rep* 2016;6:19076.
82. Ge X, Zheng L, Zhuang R, et al. The Gut Microbial Metabolite Trimethylamine N-Oxide and Hypertension Risk: A Systematic Review and Dose-Response Meta-analysis. *Adv Nutr* 2020;11(1):66-76.
83. Heianza Y, Ma W, Manson JE, et al. Gut Microbiota Metabolites and Risk of Major Adverse Cardiovascular Disease Events and Death: A Systematic Review and Meta-Analysis of Prospective Studies. *J Am Heart Assoc* 2017;6(7)
84. Schiattarella GG, Sannino A, Toscano E, et al. Gut microbe-generated metabolite trimethylamine-N-oxide as cardiovascular risk biomarker: a systematic review and dose-response meta-analysis. *Eur Heart J* 2017;38(39):2948-56.
85. Yang S, Li X, Yang F, et al. Gut Microbiota-Dependent Marker TMAO in Promoting Cardiovascular Disease: Inflammation Mechanism, Clinical Prognostic, and Potential as a Therapeutic Target. *Front Pharmacol* 2019;10:1360.
86. Yao ME, Liao PD, Zhao XJ, et al. Trimethylamine-N-oxide has prognostic value in coronary heart disease: a meta-analysis and dose-response analysis. *BMC Cardiovasc Disord* 2020;20(1):7.
87. Aragonès G, González-García S, Aguilar C, et al. Gut Microbiota-Derived Mediators as Potential Markers in Nonalcoholic Fatty Liver Disease. *Biomed Res Int* 2019;2019:8507583.
88. Bouatra S, Aziat F, Mandal R, et al. The human urine metabolome. *PLoS One* 2013;8(9):e73076.
89. Palau-Rodriguez M, Tulipani S, Isabel Queipo-Ortuño M, et al. Metabolomic insights into the intricate gut microbial-host interaction in the development of obesity and type 2 diabetes. *Front Microbiol* 2015;6:1151.
90. Psychogios N, Hau DD, Peng J, et al. The human serum metabolome. *PLoS One* 2011;6(2):e16957.
91. Serena C, Ceperuelo-Mallafré V, Keiran N, et al. Elevated circulating levels of succinate in human obesity are linked to specific gut microbiota. *Isme j* 2018;12(7):1642-57.
92. Vojinovic D, Radjabzadeh D, Kurilshikov A, et al. Relationship between gut microbiota and circulating metabolites in population-based cohorts. *Nat Commun* 2019;10(1):5813.
93. Awoyemi A, Trøseid M, Arnesen H, et al. Markers of metabolic endotoxemia as related to metabolic syndrome in an elderly male population at high cardiovascular risk: a cross-sectional study. *Diabetol Metab Syndr* 2018;10:59.
94. Barengolts E, Green SJ, Chlipala GE, et al. Predictors of Obesity among Gut Microbiota Biomarkers in African American Men with and without Diabetes. *Microorganisms* 2019;7(9).
95. Citronberg JS, Curtis KR, White E, et al. Association of gut microbial communities with plasma lipopolysaccharide-binding protein (LBP) in premenopausal women. *Isme j* 2018;12(7):1631-41.
96. Liu X, Lu L, Yao P, et al. Lipopolysaccharide binding protein, obesity status and incidence of metabolic syndrome: a prospective study among middle-aged and older Chinese. *Diabetologia* 2014;57(9):1834-41.