

Appendix

Table 1. Severity Fractions for Mental Health and Substance Use Disorders Used in this Study

DSM-5 Disorder Groupings	Disorder-Specific Definitions (If Subtypes Exist)	Severity Level	Severity Fraction
Depressive Disorders	Major Depressive Disorder	Asymptomatic	0.13
		Mild	0.59
		Moderate	0.17
		Severe	0.10
	Dysthymia	Asymptomatic	0.29
		Symptomatic	0.71
Anxiety Disorders	Anxiety Disorders	Asymptomatic	0.20
		Mild	0.50
		Moderate	0.19
		Severe	0.11
Alcohol Use Disorders	Alcohol-Use Disorders	Asymptomatic	0.32
		Very mild	0.59
		Mild	0.038
		Moderate	0.033
		Severe	0.024
Opioid-Use Disorders	Opioid-Use Disorders	Asymptomatic	0.16
		Mild	0.37
		Moderate/Severe	0.47
Cannabis-Use Disorders	Cannabis-Use Disorders	Asymptomatic	0.58
		Mild	0.36
		Moderate/Severe	0.06
Schizophrenia Spectrum and Other Psychotic Disorders	Schizophrenia	Acute state	0.63
		Residual state	0.37
Bipolar and Related Disorders	Bipolar I and Bipolar II	Manic episode	0.23
		Depressive episode	0.27
		Residual state	0.50
Attention-Deficit/Hyperactivity Disorder	ADHD	Asymptomatic	0.33
		Mild	0.36
		Moderate	0.18
		Severe	0.14
Feeding and Eating	Bulimia Nervosa and Anorexia	Asymptomatic/Mild	0.71

Disorders	Nervosa	Moderate	0.15
		Severe	0.14
Intellectual Disabilities	Intellectual Disabilities and Developmental Disorders	Asymptomatic/Mild	0.85
		Moderate	0.10
		Severe	0.03
		Profound	0.02

Figure 1. Step 9: Severity-adjusted Number of Cases in British Columbia. Data corresponding to Anxiety Disorders is shown.

