

## Supplementary

**Table S1** Bodyweight and histological changes of the experimental groups

Group	Group A	Group B	Group C
	4NQO	4NQO+SAB	4NQO+SAB-PLC-NP
<b>Bodyweight (mean, g)</b>			
Maximum (SD)	27.41 (1.73)	27.09 (1.61)	25.76 (1.19)
Minimum (SD)	18.36 (0.99)	18.91 (0.63)	19.05 (1.55)
Weight loss (SD)	9.05 (0.74)	8.17 (0.98)	6.71 (0.37)
Histological examination at 8w (n)	8	8	7
Normal	4	6	7
Mild dysplasia	2	2	0
Moderate/severe dysplasia	2	0	0
Carcinoma	0	0	0
Histological examination at 14w (n)	8	7	7
Mild dysplasia	2	3	4
Moderate/severe dysplasia	5	4	3
Carcinoma	1	0	0
Histological examination at 18w (n)	9	7	8
Mild dysplasia	1	1	3
Moderate/severe dysplasia	5	5	4
Carcinoma	3	1	1
Cumulative incidence at 18w (n)	25	22	22
No/mild dysplasia	9	12	14
Moderate/severe dysplasia	12	9	7
Carcinoma	4	1	1
Histological examination at 22w (n)	10	8	8
Mild dysplasia	0	0	2
Moderate/severe dysplasia	2	4	4
Carcinoma	8	4	2
Cumulative incidence at 22w (n)	35	30	30
No/mild dysplasia	9	12	16
Moderate/severe dysplasia	14	13	11
Carcinoma	12	5	3

SD, standard deviation; 4NQO, 4-nitroquinoline-N-oxide; SAB, salvianolic acid B; PLC-NP, phospholipid complex loaded nanoparticles.