

## **Supplemental Online Content**

Rao P, Jiang SF, Kipnis P, et al. Evaluation of outcomes following hospital-wide implementation of a subcutaneous insulin protocol for diabetic ketoacidosis. *JAMA Netw Open*. 2022;5(4):e226417. doi:10.1001/jamanetworkopen.2022.6417

**eTable.** Missingness of Laboratory Values Data in Diabetic Ketoacidosis Cohort

This supplemental material has been provided by the authors to give readers additional information about their work.

**eTable. Missingness of Laboratory Values Data in Diabetic Ketoacidosis Cohort**

	Intervention Site		Standard Care Site	
	Pre	Post	Pre	Post
Hospital encounters	n = 298	n = 122	n = 4,441	n = 3,128
Unique patients	n = 173	n = 87	n = 2,703	n = 2,083
Glucose	0(0.0)	0(0.0)	0(0.0)	0(0.0)
Anion gap	0(0.0)	0(0.0)	1(0.02)	0(0.0)
Blood urea nitrogen	0(0.0)	0(0.0)	2(0.05)	0(0.0)
Bicarbonate	0(0.0)	0(0.0)	2(0.05)	0(0.0)
Sodium	0(0.0)	0(0.0)	0(0.0)	0(0.0)
White blood cell count	0(0.0)	0(0.0)	9(0.2)	1(0.03)
Creatinine	0(0.0)	0(0.0)	0(0.0)	0(0.0)
Potassium	1(0.3)	0(0.0)	175(3.9)	95(3.0)
Chloride	0(0.0)	0(0.0)	6(0.1)	6(0.2)
Arterial PH	130(43.6)	96(78.6)	1854(41.7)	2376(76.0)
Venous PH	194(65.1)	53(43.4)	2778(62.6)	1132(36.2)