

Increasing stratification as observed by Satellite Sea Surface Salinity measurements.

October 2021

E. Olmedo^{(1)(*)}, A. Turiel⁽¹⁾, V. González-Gambau⁽¹⁾, C. González-Haro⁽¹⁾, A. García-Espriu⁽¹⁾, C. Gabarró⁽¹⁾, M. Portabella⁽¹⁾, I. Corbella⁽²⁾, M. Martin-Neira⁽³⁾, M. Arias⁽¹⁾, R. Catany⁽⁴⁾, R. Sabia⁽⁵⁾, R. Oliva⁽⁶⁾, K. Scipal⁽³⁾

(*) Corresponding author: Estrella Olmedo, olmedo@icm.csic.es

(1) Barcelona Expert Center (BEC), Institute of Marine Sciences (ICM) and Consejo Superior de Investigaciones Científicas (CSIC), P. Marítim de la Barceloneta, 37-49, 08003 Barcelona, Spain

(2) Barcelona Expert Center (BEC), Universitat Politècnica de Catalunya (UPC), Barcelona, Spain

(3) European Space Agency, ESTEC, Noordwijk, The Netherlands

(4) Argans UK Ltd., Derriford, PL6 8BX Plymouth, U.K

(5) Telespazio-UK, for ESA-ESRIN, Largo Galileo Galilei 1, 00044 Frascati, Italy

(6) Zenithal Blue Technologies, 08023 Barcelona, Spain

1 SUPPLEMENTARY MATERIAL

Near Surface Salinity average in 2011-2018 as observed by Argo floats

Near Surface Salinity Trends in 2011-2018 as observed by Argo floats




Figure 1: Left: Near-surface salinity average (left) and linear trend (right) in 2011-2018 as observed by Argo floats. Maps are plotted with Panoply v 4.12.0 (<https://www.giss.nasa.gov/tools/panoply/>).