

**S2 Table. Comparison of different objective functions and pre-processing techniques.**

Dataset	$J_{MVIB-T}$		$J_{MVIB}$	
	D	J	D	J
IBD	0.922 (0.020)	<b>0.936</b> <b>(0.014)</b>	0.915 (0.018)	0.915 (0.018)
EW-T2D	<b>0.859</b> <b>(0.023)</b>	0.853 (0.025)	0.859 (0.024)	0.855 (0.027)
C-T2D	0.750 (0.009)	<b>0.758</b> <b>(0.012)</b>	0.754 (0.014)	0.756 (0.015)
Obesity	0.662 (0.024)	0.666 (0.027)	<b>0.673</b> <b>(0.028)</b>	0.672 (0.030)
Cirrhosis	0.925 (0.005)	0.924 (0.005)	<b>0.930</b> <b>(0.002)</b>	0.928 (0.002)
Colorectal	0.780 (0.071)	0.777 (0.069)	0.788 (0.059)	<b>0.796</b> <b>(0.055)</b>
Obesity-Joint	0.815 (0.019)	0.818 (0.018)	0.825 (0.019)	<b>0.825</b> <b>(0.018)</b>
Colorectal-EMBL	0.811 (0.010)	0.814 (0.013)	0.827 (0.012)	<b>0.830</b> <b>(0.011)</b>
Early-Colorectal-EMBL	0.535 (0.050)	<b>0.543</b> <b>(0.048)</b>	0.525 (0.051)	0.533 (0.053)
Hypertension	0.602 (0.045)	0.608 (0.043)	0.622 (0.048)	<b>0.624</b> <b>(0.049)</b>