

**SUPPLEMENTARY TABLE 6. sICH (FPR vs REcanalized non-FPR)**

<b>AUTHORS, YEAR</b>	<b>FPR DEFINITION</b>	<b>sICH DEFINITION</b>	<b>FPR</b>	<b>FPR-TOTAL</b>	<b>NON-FPR</b>	<b>NON-FPR TOTAL</b>	<b>THROMBECTOMY TECHNIQUE</b>
<b>MOHAMMADEN ET AL, 2020<sup>67</sup></b>	2c	Not provided	5	167	5	113	Aspiration and/or Stent-retriever
<b>TORNEL ET AL, 2019<sup>62</sup></b>	2b	Not provided	11	213	16	246	Aspiration
<b>Z Aidat ET AL, 2018<sup>4</sup></b>	2b	any intracranial hemorrhage associated with neurological deterioration of $\geq 4$ points on the NIHSS score at 24 hours	5	43	9	70	Stent-retriever
<b>HUN KANG ET AL, 2019<sup>65</sup></b>	2c	Intracranial hemorrhage was regarded as symptomatic (sICH) if the NIHSS score increased to $\geq 4$ .	4	66	10	278	Aspiration

FPR:First pass revascularization; sICH: symptomatic intracranial hemorrhage.