

**Supplementary Figures and Tables**

<p>Q4.1 <b>SECTION 2A</b></p> <p>Q4.2</p>	<p>Have you read the new <i>BAPM Framework for Practice on Perinatal Management of Extreme Preterm Birth before 27 weeks of gestation</i>?</p> <p><input type="radio"/> Yes</p> <p><input type="radio"/> No</p>
<p>Q4.3</p>	<p>Are you using the new <i>BAPM Framework for Practice on Perinatal Management of Extreme Preterm Birth before 27 weeks of gestation</i> to guide your clinical practice?</p> <p><input type="radio"/> Yes</p> <p><input type="radio"/> No</p>
<p>Q4.4</p>	<p>If no, why not?</p>
<p>Q4.5 <b>SECTION 2B</b></p>	<p>We would like you to read the following scenarios and use the new <i>BAPM Framework for Practice on Perinatal Management of Extreme Preterm Birth before 27 weeks of gestation</i> to consider the risk you would assign to each case (using the information in figure 1: <i>Visual tool for refinement of risk</i> and box 1: <i>Consensus for risk categorisation</i>).</p> <p>After assigning the risk we would like to think about what outcome you would estimate for the baby and how you would manage the case around delivery (using the information from figure 2: <i>Flow diagram for decision making around management of delivery</i>).</p> <p>If you would like to review these figures, they will appear at the bottom of the screen for all of the following questions.</p> <p>For each case, please assume that that an early dating scan has been performed to estimate gestational age, and that there are no other known medical conditions affecting the fetus or mother.</p>

Note that the cases will appear in random order.




Figure 1: Proposed visual tool for refinement of risk

**BOX 1**

**Extremely high risk:** The Working Group considered that babies with a > 90% chance of either dying or surviving with severe impairment if active care is instigated would fit into this category. For example, this would include:

- babies at 22<sup>0</sup> - 22<sup>6</sup> weeks of gestation with unfavourable risk factors
- some babies at 23<sup>0</sup> - 23<sup>6</sup> weeks of gestation with unfavourable risk factors, including severe fetal growth restriction
- (rarely) babies ≥ 24<sup>0</sup> weeks of gestation with significant unfavourable risk factors, including severe fetal growth restriction


**High risk:** The Working Group considered that babies with a 50-90% chance of either dying or surviving with severe impairment if active care is instituted would fit into this category. For example, this would include:

- babies at 22<sup>0</sup> - 23<sup>6</sup> weeks of gestation with favourable risk factors
- some babies ≥ 24<sup>0</sup> weeks of gestation with unfavourable risk factors and/or co-morbidities

**Moderate risk:** The Working Group considered that babies with a < 50% chance of either dying or surviving with severe impairment if active care is instituted would fit into this category. For example, this would include:

- most babies ≥ 24<sup>0</sup> weeks of gestation
- some babies at 23<sup>0</sup> - 23<sup>6</sup> weeks of gestation with favourable risk factors.

Box 1 represents the consensus of the Working Group in regard to risk categories for the purposes of this framework.



\*However, assess for modifiable risk factors and reassess risk if/when circumstances change

Figure 2. Decision-making around management of delivery, following risk assessment and after consultation with parents.

Q5.1	<b>Case A;</b> A mother has gone into extremely preterm labour at <b>23+3</b> weeks' gestation. She is in a hospital with a <b>NICU</b> and has received a full course of <b>steroids</b> . The fetus is a <b>singleton</b> , has normal growth (estimated fetal weight of <b>590g</b> ) and is known to be <b>female</b> .
Q5.2	Using the risk categories taken from the BAPM Framework for Practice what level of risk would you assign to this case? (Scroll down to the bottom of this screen if you would like to review the figures from the guidance).  <input type="radio"/> Extremely high risk  <input type="radio"/> High risk  <input type="radio"/> Moderate risk
Q5.3	If the baby receives active resuscitation, what is your estimate for the chance of survival to discharge from hospital to the nearest 5%?
Q5.4	If the baby survives to discharge, what do you think is the chance of severe disability to the nearest 5%?
Q5.5	Based on your understanding of the BAPM Framework for Practice, what management is recommended in this case? (Scroll down to the bottom of this screen if you would like to review the figures from the guidance).  <input type="radio"/> Recommend palliative (comfort focused) management  <input type="radio"/> Be guided by parent's views on whether to choose palliative (comfort focused) management or active (survival focused) management  <input type="radio"/> Recommend active (survival focused) management
Q5.9	You have indicated the BAPM Framework for Practice would recommend active (survival focused) management. If you were caring for this patient, would you support this management plan?  <input type="radio"/> Yes  <input type="radio"/> No
Q5.10	If no, please elaborate on why:
Q5.11	Would you be prepared to offer palliative (comfort focused) management at parents' request?  <input type="radio"/> Yes  <input type="radio"/> No

Q5.12	<p>You have indicated the BAPM Framework for Practice would recommend palliative (comfort focused) management. If you were caring for this patient, would you support this management plan?</p> <p><input type="radio"/> Yes</p> <p><input type="radio"/> No</p>
Q5.13	If no, please elaborate on why:
Q5.14	<p>If the parents are insistent that they wish for active resuscitation, should this be an option?</p> <p><input type="radio"/> Yes</p> <p><input type="radio"/> No</p>
Q5.15	<p>You have indicated the BAPM Framework for Practice would recommend being guided by parent's views on whether to choose palliative (comfort focused) management or active (survival focused) management. If you were caring for this patient, would you support this management plan?</p> <p><input type="radio"/> Yes</p> <p><input type="radio"/> No</p>
Q5.16	If no, please elaborate on why:
Questions 5.2 -5.16 repeated for each of the following case:	
	Q6.1 Case D; A mother has gone into extremely preterm labour at 22+3 weeks gestation. She is currently in a hospital with a NICU and has received a full course of steroids. The fetus is a singleton, normally grown (estimated fetal weight 500g) and is known to be female.
	Q7.1 Case E; A mother has gone into extremely preterm labour at 23+4 weeks gestation. She is currently in a local hospital and has had no steroids. The fetus is a singleton, with growth restriction (estimated fetal weight 450g) and is known to be male.
	Q8.1 Case B; A mother has gone into extremely preterm labour at 25+0 weeks gestation, with a twin pregnancy. She is currently in a local hospital and has had no steroids. Both twins are male and have growth restriction (estimated fetal weight of twin 1 is 520g and estimated fetal weight of twin 2 is 560g).

	Q9.1 Case C; A mother has gone into extremely preterm labour at 22+6 weeks gestation. She is currently in a hospital with a NICU and has had a full course of steroids. The male fetus has normal growth (estimated fetal weight 600g).
--	---

**Supplementary Table 1:** Full text of survey questions. The study was approved by the University of Oxford Central University Research Ethics Council (R69766/RE001) in June 2020 and was registered with the Monash University Human Research Ethics Committee (Ref 23763).

Characteristic (n= number of responses)	n (%)
<b>Professional Role (n= 336)</b>	
Consultant	167 (50)
Registrar	109 (32)
ANNP	42 (13)
Other	18 (5)
<b>Years working with extremely preterm infants (n= 200)</b>	
0-3	12 (6)
4-7	57 (29)
8-15	50 (25)
>16	81 (40)
<b>Age (years) (n=193)</b>	
18-30	3 (2)
31-40	89 (46)
41-50	41 (21)
51-60	51 (26)
61-70	7 (4)
>71	2 (1)
<b>Neonatal centre usually worked in (n= 203)</b>	
NICU	158 (78)
LNU	40 (20)
SCU	4 (2)
<b>Gender (n=198)</b>	
Female	136 (68)
Male	61 (31)
Other	1 (1)

**Supplementary Table 2:** Baseline characteristics of respondents.

Special Care Units (SCU): These provide special care for their own local population. They also provide, by agreement with their neonatal network, some high dependency services.

Local Neonatal Units (LNU): These provide special care and high dependency care and a restricted volume of intensive care (as agreed locally) and would expect to transfer babies who require complex or longer-term intensive care to a Neonatal Intensive Care Unit.

Neonatal Intensive Care Unit (NICU): These are larger intensive care units that provide the whole range of medical (and sometimes surgical) neonatal care for their local population and additional care for babies and their families referred from the neonatal network in which they are based, and also from other networks when necessary, to deal with peaks of demand or requests for specialist care not available elsewhere. Many will be sited within perinatal centres that are able to offer similarly complex obstetric care. These units will also require close working arrangements with all of the relevant paediatric sub-specialties.

(As referenced from British Association of Perinatal Medicine Service Standards for Hospitals Providing Neonatal Care (3<sup>rd</sup> Edition 2010)

Case	Risk categorisation (n) vs management recommendation (n) with P Value		
	Extremely high vs palliative care	High vs Seeking Parents' Wishes	Moderate vs active treatment
A (23+3F)	12 vs 3, P=0.007	168 vs 153, P=0.032	44 vs 68, P<0.001
B (25+0U)	43 vs 17, P< 0.001	159 vs 93, P<0.001	19 vs 111, P<0.001
C (22+6F)	72 vs 28, P< 0.001	151 vs 193, P<0.001	3 vs 5, P=0.68
D (22+3F)	89 vs 52, P< 0.001	135 vs 172, P<0.001	2 vs 2, P=0.62
E (23+4U)	202 vs 128, P< 0.001	19 vs 91, P<0.001	0 vs 2, P=1

**Supplementary Table 3;** Number of respondents categorising extremely high, high and moderate risk versus number of respondents recommending palliative care, seeking parents' wishes and active treatment respectively. P values represent chance of difference between risk categorisation and corresponding management recommendation from the BAPM guideline using McNemar's test. Statistical significance was taken as  $p < 0.05$ .

Each case has been given an abbreviation of the gestational age followed by U or F, depending whether there is a majority of unfavourable (U) or favourable (F) risk factors.

**Comparison of risk management between groups**

A significantly higher number of ANNPs (43%) compared to consultants (14%) and registrars/fellows (20%) classified the risk for case B (25+0U) as extremely-high ( $p < 0.001$  and  $0.02$  respectively). Fewer consultants (3%) recommended palliative care compared to registrars/fellows (13%) and ANNPs (13%) ( $p = 0.08$  and  $0.03$  respectively).

A higher proportion of respondents working in a local neonatal unit (LNU) or a special care unit (SCU) compared to those working in a NICU recommended palliative care compared to from NICUs for case C (22+6F) (25 vs 10%,  $p = 0.007$ ).

CASE A (23+3F)	Number (%)			X <sup>2</sup> and P value for difference between groups		
	Consultant	Reg/ Fellow	ANNP	Consultant and Reg/Fellow	Consultant and ANNP	Reg/Fellow and ANNP
Risk and Management Allocation						
Extremely High	11 (10)	1 (1)	0 (0)	X <sup>2</sup> 6.2 p 0.01	X <sup>2</sup> 0.82 p 0.36	X <sup>2</sup> 0.88 p 0.35
High	79 (69)	71 (83)	18 (75)	X <sup>2</sup> 4.6 p 0.03	X <sup>2</sup> 0.31 p 0.58	X <sup>2</sup> 0.69 p 0.40
Moderate	24 (21)	14 (16)	6 (25)	X <sup>2</sup> 0.72 p 0.39	X <sup>2</sup> 1.13 p 0.29	X <sup>2</sup> 2.66 p 0.10
Palliative Care	3 (3)	0 (0)	0 (0)	X <sup>2</sup> 2.3 p 0.13	X <sup>2</sup> 0.64 p 0.42	N/A
Seek parents' wishes	72 (63)	62 (72)	19 (79)	X <sup>2</sup> 1.77 p 0.18	X <sup>2</sup> 2.26 p 0.13	X <sup>2</sup> 0.48 p 0.49
Active Care	39 (34)	24 (28)	5 (21)	X <sup>2</sup> 0.9 p 0.342	X <sup>2</sup> 1.63 p 0.2	X <sup>2</sup> 0.18 p 0.67

CASE B (25+0U)	Number (%)			X <sup>2</sup> and P value for difference between groups		
	Consultant	Reg/ Fellow	ANNP	Consultant and Reg/Fellow	Consultant and ANNP	Reg/Fellow and ANNP
Risk and Management Allocation						
Extremely High	15 (14)	18 (20)	10 (43)	X <sup>2</sup> 1.64 p 0.2	<b>X<sup>2</sup> 11 p &lt; 0.001</b>	X <sup>2</sup> 5.12 p 0.02
High	82 (75)	65 (74)	12 (52)	X <sup>2</sup> 0.01 p 0.91	X <sup>2</sup> 4.6 p 0.03	X <sup>2</sup> 4.03 p 0.04
Moderate	13 (12)	5 (6)	1 (4)	X <sup>2</sup> 2.22 p 0.14	X <sup>2</sup> 1.13 p 0.29	X <sup>2</sup> 0.063 p 0.08
Palliative Care	3 (3)	11 (13)	3 (13)	X <sup>2</sup> 7.1 p 0.008	X <sup>2</sup> 4.7 p 0.03	X <sup>2</sup> 0.005 p 0.94
Seek parents' wishes	40 (36)	43 (49)	10 (43)	X <sup>2</sup> 3.13 p 0.08	X <sup>2</sup> 0.41 p 0.52	X <sup>2</sup> 0.21 p 0.65
Active Care	67 (61)	34 (39)	10 (43)	X <sup>2</sup> 9.7 p 0.002	X <sup>2</sup> 2.7 p 0.12	X <sup>2</sup> 0.18 p 0.67



CASE C (22+6F)	Number (%)			X <sup>2</sup> and P value for difference between groups		
	Consultant	Reg/ Fellow	ANNP	Consultant and Reg/Fellow	Consultant and ANNP	Reg/Fellow and ANNP
Extremely High	72 (64)	61 (70)	18 (69)	X <sup>2</sup> 0.9 p 0.34	X <sup>2</sup> 0.28 p 0.60	X <sup>2</sup> 0.007 p 0.93
High	39 (35)	25 (29)	8 (31)	X <sup>2</sup> 0.75 p 0.38	X <sup>2</sup> 0.13 p 0.72	X <sup>2</sup> 0.04 p 0.84
Moderate	2 (2)	1 (1)	0 (0)	X <sup>2</sup> 0.13 p 0.72	X <sup>2</sup> 0.47 p 0.49	X <sup>2</sup> 0.3 p 0.58
Palliative Care	16 (14)	9 (10)	3 (12)	X <sup>2</sup> 0.65 p 0.42	X <sup>2</sup> 0.12 p 0.73	X <sup>2</sup> 0.03 p 0.86
Seek parents' wishes	93 (82)	77 (89)	23 (88)	X <sup>2</sup> 1.38 p 0.22	X <sup>2</sup> 0.58 p 0.45	X <sup>2</sup> <0.001 p 0.99
Active Care	4 (4)	1 (1)	0 (0)	X <sup>2</sup> 1.15 p 0.28	X <sup>2</sup> 0.95 p 0.33	X <sup>2</sup> 0.3 p 0.58

CASE D (22+3F)	Number (%)			X <sup>2</sup> and P value for difference between groups		
	Consultant	Reg/ Fellow	ANNP	Consultant and Reg/Fellow	Consultant and ANNP	Reg/Fellow and ANNP
Extremely High	46 (41)	32 (36)	11 (46)	X <sup>2</sup> 0.64 p 0.42	X <sup>2</sup> 0.18 p 0.67	X <sup>2</sup> 0.85 p 0.36
High	66 (59)	56 (62)	13 (54)	X <sup>2</sup> 0.22 p 0.63	X <sup>2</sup> 0.18 p 0.67	X <sup>2</sup> 0.51 p 0.47
Moderate	0 (0)	2 (2)	0 (0)	X <sup>2</sup> 2.5 p 0.11	N/A	X <sup>2</sup> 0.27 p 0.60
Palliative Care	32 (29)	13 (14)	7 (29)	X <sup>2</sup> 5.7 p 0.02	X <sup>2</sup> 0.003 p 0.95	X <sup>2</sup> 2.84 p 0.09
Seek parents' wishes	80 (71)	76 (84)	16 (67)	X <sup>2</sup> 4.8 p 0.02	X <sup>2</sup> 0.22 p 0.64	X <sup>2</sup> 3.8 p 0.05
Active Care	0 (0)	1 (1)	1 (4)	X <sup>2</sup> 1.25 p 0.26	X <sup>2</sup> 4.7 p 0.03	X <sup>2</sup> 1.03 p 0.31

CASE E (23+4U)	Number (%)			X <sup>2</sup> and P value for difference between groups		
	Consultant	Reg/ Fellow	ANNP	Consultant and Reg/Fellow	Consultant and ANNP	Reg/Fellow and ANNP
Extremely High	103 (93)	77 (91)	22 (88)	X <sup>2</sup> 0.31 p 0.58	X <sup>2</sup> 0.63 p 0.43	X <sup>2</sup> 0.14 p 0.70
High	8 (7)	8 (9)	3 (12)	X <sup>2</sup> 0.31 p 0.58	X <sup>2</sup> 0.63 p 0.43	X <sup>2</sup> 0.14 p 0.70
Moderate	0 (0)	0 (0)	0 (0)	N/A	N/A	N/A
Palliative Care	64 (58)	54 (64)	10 (40)	X <sup>2</sup> 0.69 p 0.41	X <sup>2</sup> 2.56 p 0.11	X <sup>2</sup> 4.4 p 0.04
Seek parents' wishes	45 (41)	31 (36)	15 (60)	X <sup>2</sup> 0.34 p 0.56	X <sup>2</sup> 3.1 p 0.08	X <sup>2</sup> 4.4 p 0.04
Active Care	2 (2)	0 (0)	0 (0)	X <sup>2</sup> 1.55 p 0.21	X <sup>2</sup> 0.46 p 0.5	N/A

**Supplementary Table 4:** For cases A to E the number and percentage of respondents classifying the risk as extremely high, high or moderate and allocating the management as palliative care, seeking parents' wishes or active care by professional group, with the X<sup>2</sup> and P values between these estimates by professional groups, as calculated by Chi Squared Test. Statistical significance was taken as p<0.001.

Each case has been given an abbreviation of the gestational age followed by U or F, depending whether there is a majority of unfavourable (U) or favourable (F) risk factors.

CASE A (23+3F)	Number (%)		X <sup>2</sup> and P value for difference between groups
Risk and Management Allocation	NICU	LNU/SCU	NICU and LNU/SCU
Extremely High	5 (3)	6 (14)	X <sup>2</sup> 7.32 p 0.007
High	122 (77)	30 (68)	X <sup>2</sup> 1.5 p 0.22
Moderate	31 (20)	8 (18)	X <sup>2</sup> 0.04 p 0.83
Palliative Care	0 (0)	3 (7)	<b>X<sup>2</sup> 10.9 p &lt;0.001</b>
Seek parents' wishes	111 (70)	29 (66)	X <sup>2</sup> 0.31 p 0.58
Active Care	47 (30)	12 (27)	X <sup>2</sup> 0.101 p 0.75

CASE B (25+0U)	Number (%)		X <sup>2</sup> and P value for difference between groups
Risk and Management Allocation	NICU	LNU/SCU	NICU and LNU/SCU
Extremely High	29 (18)	8 (18)	X <sup>2</sup> 0.002 p 0.965
High	112 (71)	33 (75)	X <sup>2</sup> 0.23 p 0.63
Moderate	16 (10)	3 (7)	X <sup>2</sup> 0.46 p 0.5
Palliative Care	12 (8)	2 (5)	X <sup>2</sup> 0.51 p 0.48
Seek parents' wishes	60 (38)	26 (59)	X <sup>2</sup> 6.11 p 0.01
Active Care	85 (54)	16 (36)	X <sup>2</sup> 4.3 p 0.04

CASE C (22+6F)	Number (%)		X <sup>2</sup> and P value for difference between groups
Risk and Management Allocation	NICU	LNU/SCU	NICU and LNU/SCU
Extremely High	47 (30)	18 (41)	X <sup>2</sup> 0.89 p 0.17
High	109 (69)	25 (57)	X <sup>2</sup> 2.46 p 0.12
Moderate	1 (1)	1 (2)	X <sup>2</sup> 0.933 p 0.334
Palliative Care	15 (10)	11 (25)	X <sup>2</sup> 7.28 p 0.007
Seek parents' wishes	138 (88)	32 (73)	X <sup>2</sup> 6.1 p 0.01
Active Care	4 (3)	1 (2)	X <sup>2</sup> 0.01 p 0.917


CASE D (22+3F)	Number (%)		X <sup>2</sup> and P value for difference between groups
Risk and Management Allocation	NICU	LNU/SCU	NICU and LNU/SCU
Extremely High	63 (40)	18 (41)	X <sup>2</sup> 0.009 p 0.93
High	93 (59)	26 (59)	X <sup>2</sup> 0.0003 p 0.986
Moderate	1 (1)	0 (0)	X <sup>2</sup> 0.28 p 0.60
Palliative Care	36 (23)	14 (32)	X <sup>2</sup> 1.45 p 0.23
Seek parents' wishes	120 (76)	30 (68)	X <sup>2</sup> 1.23 p 0.26
Active Care	1 (1)	0 (0)	X <sup>2</sup> 0.28 p 0.6

CASE E (23+4U)	Number (%)		X <sup>2</sup> and P value for difference between groups
	NICU	LNU/SCU	NICU and LNU/SCU
<b>Risk and Management Allocation</b>			
<b>Extremely High</b>	147 (94)	38 (86)	$\chi^2$ 2.48 p 0.12
<b>High</b>	10 (6)	6 (14)	$\chi^2$ 2.48 p 0.12
<b>Moderate</b>	0 (0)	0 (0)	N/A
<b>Palliative Care</b>	98 (62)	24 (55)	$\chi^2$ 0.89 p 0.35
<b>Seek parents' wishes</b>	58 (37)	20 (45)	$\chi^2$ 1.04 p 0.31
<b>Active Care</b>	1 (1)	0 (0)	$\chi^2$ 0.29 p 0.60


**Supplementary Table 5:** For cases A to E the number and percentage of respondents classifying the risk as extremely high, high or moderate and allocating the management as palliative care, seeking parents' wishes or active care by centre worked in (neonatal intensive care unit (NICU) or local neonatal unit (LNU)/special care unit (SCU)), with the  $\chi^2$  and P values between these estimates by centre worked in, as calculated by Chi Squared Test. Statistical significance was taken as  $p < 0.001$ .

Each case has been given an abbreviation of the gestational age followed by U or F, depending whether there is a majority of unfavourable (U) or favourable (F) risk factors.


**Case A (23+3, favourable risk factors)**




**Case B (25+0, unfavourable risk factors)**




**Case C (22+6, favourable risk factors)**






**Supplementary figure 1:** Dot plots showing respondents estimates of survival in percentages by professional group. Each dot represents one respondent's estimates. The horizontal lines represent the median and interquartile range. There was no statistical difference found between the different professional group's estimates (using Mann Whitney U test. Statistical significance was taken as  $p < 0.005$ )


**Case A (23+3, favourable risk factors)**




**Case B (25+0, unfavourable risk factors)**




**Case C (22+6, favourable risk factors)**



**Case D (22+3, favourable risk factors)**







**Supplementary figure 2:** Dot plots showing respondents estimates of survival and severe disability in percentages by type of unit worked in. Each dot represents one respondent's estimates. The horizontal lines represent the median and interquartile range. There was no statistical difference found between type of unit worked in for both estimates of survival and severe disability (using Mann Whitney U test. Statistical significance was taken as  $p < 0.005$ )




**Case A (23+3, favourable risk factors)**




**Case B (25+0, unfavourable risk factors)**



**Case C (22+6, favourable risk factors)**





**Supplementary Figure 3:** Dot plots showing respondents estimates of severe disability in percentages by professional group. Each dot represents one respondent's estimates. Horizontal lines represent the median and interquartile range. There was statistically significant difference between estimates of consultants and registrars in cases B and D (case B;  $U=3381$ ,  $p<0.0001$  and case D;  $U=3135$ ,  $p 0.004$ ) and between consultants and ANNPs in cases B, C and D (case B;  $U=564$ ,  $p <0.001$ , case C;  $U 826$ ,  $p<0.001$  and case D;  $U 531$ ,  $p <0.001$ )