

Biointerface topography mediates the interplays between endothelial cells and monocytes

Yan Liu ^a, Wenshuai Deng ^d, Liangliang Yang ^e, Xiuxiu Fu ^c, Zhibin Wang ^c, Patrick
van Rijn ^{*e}, Qihui Zhou ^{*a,b}, Tao Yu ^{*a,c}

Author Affiliations:

^a Institute for Translational Medicine, School of Basic Medicine, Qingdao University, Qingdao, 266021, China.

^b Stomatology Center, The Affiliated Hospital of Qingdao University, Qingdao, 266003, China.

^c Department of Echocardiography, The Affiliated Hospital of Qingdao University, Qingdao, 26603, China.

^d Department of Neurosurgery, The Affiliated Hospital of Qingdao University, Qingdao 266003, China.

^e University of Groningen, W.J. Kolff Institute for Biomedical Engineering and Materials Science, Department of Biomedical Engineering, University Medical Center Groningen, A. Deusinglaan 1, 9713 AV, Groningen, the Netherlands.

*Corresponding author:

Patrick van Rijn, Ph.D., Associate professor of Biomedical Engineering, Email:
p.van.rijn@umcg.nl.

Qihui Zhou, Ph.D., Associate professor of Biomaterials, Phone: +86-17660670299,
Email: qihuizhou@qdu.edu.cn.

Tao Yu, Ph.D., Associate professor of Basic Medicine, Email:
yutao0112@qdu.edu.cn.




Figure S1. Imprinted PDMS wrinkles using the wrinkled PDMS surface as the template.