

**Supplementary Table I (online only).** Upper extremity, lower extremity, and central venous catheter-associated DVT

DVT	Central venous catheter association		Total
	Yes	No	
Upper extremity <sup>a</sup>	12 (0.8)	6 (0.4)	18 (1.2)
Lower extremity <sup>b</sup>	1 (0.1)	18 (1.2)	19 (1.3)
Total	13 (0.9)	24 (1.7)	37 (2.6)

DVT, Deep vein thrombosis.  
Data presented as number (% of total cohort [n = 1449]).  
<sup>a</sup>Included axillary, subclavian, and internal jugular vein DVTs.  
<sup>b</sup>Included popliteal, femoral, and iliac vein DVTs.

**Supplementary Table II (online only).** Logistic regression for association of Caprini score with postoperative and postdischarge VTE<sup>a</sup>

Caprini score, points	Odds ratio (95% CI)			
	VTE	P value	Postdischarge VTE	P value
2-6 (reference; n = 1033)	1.0	NA	NA	NA
7-9 (n = 321)	3.6 (1.8-7.0)	<.01	3.3 (1.1-9.4)	.03
≥10 (n = 95)	10.3 (4.9-21.7)	<.01	6.4 (1.9-22.4)	<.01

CI, Confidence interval; NA, not applicable; VTE, venous thromboembolism.  
<sup>a</sup>C-statistic = 0.71 for all VTE and 0.68 for postdischarge VTE (n = 1449).

**Supplementary Table III (online only).** Cox proportional hazards model for association of VTE with mortality (N = 1449; n = 49 for VTE, including symptomatic and asymptomatic; C-statistic = 0.74)

Variable	HR	95% CI	P value
Postoperative VTE	2.7	1.3-5.9	.01
Age, years			
<60 (reference)	1.0		
61-70	1.1	0.5-2.1	.84
71-80	1.6	0.8-3.0	.18
>80	1.8	0.9-3.6	.11
ESRD	2.7	1.7-4.4	<.01
Prior VTE	0.7	0.3-1.8	.41
Length of stay (per 5 days)	1.1	0.9-1.3	.40
Nonelective admission	2.9	1.7-5.0	<.01
Procedure type			
Open aortic (reference)	1.0		
TEVAR	0.6	0.3-1.2	.13
EVAR	0.3	0.1-0.9	.04
Lower extremity bypass	0.5	0.3-0.9	.02
Femoral endarterectomy	1.0	0.5-2.1	.98

CI, Confidence interval; ESRD, end-stage renal disease; EVAR, endovascular abdominal aortic aneurysm repair; HR, hazard ratio; TEVAR, thoracic endovascular aortic aneurysm repair; VTE, venous thromboembolism.