

Electronic Supporting Information for

Solid-State AIEnh-Circularly Polarised Luminescence of Chiral Perylene Diimide Fluorophores

Ayano Taniguchi,^a Daiki Kaji,^a Nobuyuki Hara,^a Ryosuke Murata,^b Shogo Akiyama,^a Takunori Harada,^c Atsushi Sudo,^a Hiroyuki Nishikawa,^{*b} and Yoshitane Imai^{*a}

^a Department of Applied Chemistry, Faculty of Science and Engineering, Kindai University, 3-4-1 Kowakae, Higashi-Osaka, Osaka 577-8502, Japan. E-mail: (Y.I.) y-imai@apch.kindai.ac.jp.

^b Graduate School of Science and Engineering, Ibaraki University, 2-1-1 Bunkyo, Mito, Ibaraki 310-8512, Japan. E-mail: (H.N.) hiroyuki.nishikawa.sci@vc.ibaraki.ac.jp

^c Department of Integrated Science and Technology, Faculty of Science and Technology, Oita University, Dannoharu, 700, Oita city 870-1192, Japan

Table of Contents

Fig. S1	¹ H-NMR spectrum of (<i>R,R</i>)- BPP in CDCl ₃ (500 MHz).....	S2
Fig. S2	¹³ C-NMR spectrum of (<i>R,R</i>)- BPP in CDCl ₃ (125 MHz).....	S2
Fig. S3	¹ H-NMR spectrum of (<i>S,S</i>)- BPP in CDCl ₃ (500 MHz).....	S3
Fig. S4	¹³ C-NMR spectrum of (<i>S,S</i>)- BPP in CDCl ₃ (125 MHz).....	S3




Fig. S1 ^1H -NMR spectrum of (*R,R*)-**BPP** in CDCl_3 (500 MHz).




Fig. S2 ^{13}C -NMR spectrum of (*R,R*)-**BPP** in CDCl_3 (125 MHz).




Fig. S3 ^1H -NMR spectrum of (*S,S*)-BPP in CDCl_3 (500 MHz).




Fig. S4 ^{13}C -NMR spectrum of (*S,S*)-BPP in CDCl_3 (125 MHz).