**Table S2.** Numbers needed to harm (NNH) for exposure to diclofenac (years 2005-2015).

Outcomes	No analgesic (n=20,544) n (%)	Diclofenac 2005-2015 (n=10,291)	
	()	n (%)	NNH
Gestation at delivery (weeks)			
>=37	19,407 (94.5%)	9,640 (93.7%)	
<37	1,137 (5.5%)	651 (6.3%)	125
Pregnancy outcome			
Livebirth	20,393 (99.3%)	10,227 (99.4%)	
Stillbirth	116 (0.5%)	39 (0.4%)	n/a
Neonatal Death	35 (0.2%)	25 (0.2%)	n/a
Weight of baby (grams)			
NBW	16,869 (82.1%)	8,116 (78.9%)	
LBW	965 (4.7%)	572 (5.6%)	111
HBW	2,707 (13.2%)	1,600 (15.5%)	44
Missing	3 (0.0%)	3 (0.0%)	
Admitted to neonatal unit			
No	18,224 (88.7%)	8,747 (85.0%)	
Yes	2,175 (10.6%)	1,492 (14.5%)	26
Missing	145 (0.7%)	52 (0.5%)	
APGAR score at 1 min			
Normal	18,709 (91.1%)	9,350 (90.9%)	
<7	1,658 (8.1%)	924 (9.0%)	111
Missing	177 (0.9%)	17 (0.2%)	
APGAR score at 5 min			
Normal	20,065 (97.7%)	10,096 (98.1%)	
<7	302 (1.5%)	177 (1.7%)	500
Missing	177 (0.9%)	18 (0.2%)	
Cryptorchidism (only males included)			
No	10,284 (98.7%)	5,314 (98.7%)	_
Yes	133 (1.3%)	70 (1.3%)	n/a
Neural Tube D			1
No	20,527 (99.9%)	10,263 (99.7%)	
Yes	17 (0.1%)	28 (0.3%)	500
Amniotic Band Defects			
No	20,514 (99.9%)	10,277 (99.9%)	
Yes	30 (0.1%)	14 (0.1%)	n/a
Hypospadias (only males included)			
No	10,317 (99.0%)	5,308 (98.6%)	250
Yes	100 (1.0%)	76 (1.4%)	250
Gastroschisis	20 520 (00 00/)	10.204 (20.00()	
No	20,538 (99.9%)	10,284 (99.9%)	
Yes	6 (0.1%)	7 (0.1%)	n/a
At least one o		10.007 (00.10)	1
No	20,258 (98.6%)	10,097 (98.1%)	200
Yes	286 (1.4%)	194 (1.9%)	200

n/a, not applicable