

Procedure	Average OMEs prescribed/patient (SD)	Average OMEs over state-recommended dose	Number of patients prescribed more than state-recommended dose	Percent of patients prescribed more than state-recommended dose
Immediate breast reconstruction with tissue expanders or implants (n=73)	280.2 (153.6)	70.2	34	46.6%
Breast reconstruction revision (n=56)	125.6 (62.4)	-84.4	2	3.6%
Panniculectomy or abdominoplasty (n=43)	193.1 (75.9)	-16.9	9	20.9%
Excision of skin or soft tissue (n=35)	110.8 (98.5)	-99.2	2	5.7%
Breast reduction or mastopexy only (n=33)	142.7 (50.5)	-67.3	1	3.0%
Wound debridement (n=32)	201.3 (199.7)	-8.7	8	25.0%
Lower extremity or upper extremity reconstruction with local flap (n=24)	239.6 (205.5)	29.6	8	33.3%
Delayed breast reconstruction with free flap (n=21)	140.2 (53.4)	-69.8	0	0.0%
Immediate breast reconstruction with free flap (n=18)	165.4 (82.7)	-44.6	2	11.1%
Skin graft only (n=18)	253.9 (302.9)	43.9	4	22.2%
Amputation w/targeted muscle reinnervation (n=16)	595.0 (666.3)	385.0	12	75.0%
Spine reconstruction (n=14)	948.6 (2646.0)	738.6	5	35.7%
Abdominal wall reconstruction (n=13)	369.7 (534.4)	159.7	4	30.8%
Lower extremity or upper extremity reconstruction with free flap (n=11)	442.5 (520.2)	232.5	4	36.4%
Liposuction or fat grafting, non-breast reconstruction (n=11)	130.2 (52.6)	-79.8	0	0.0%
Trunk or pelvic reconstruction with local flaps (n=10)	531.9 (772.7)	321.9	5	50.0%
Delayed breast reconstruction with tissue expanders or implants (n=9)	161.4 (39.7)	-48.6	0	0.0%
Oncoplastic breast reduction or mastopexy (n=9)	168.9 (61.8)	-41.1	1	11.1%
Sternal wound reconstruction (n=9)	123.3 (112.4)	-86.7	1	11.1%
Lymphovenous bypass only (n=7)	87.1 (101.8)	-122.9	1	14.3%
Targeted muscle reinnervation only (n=6)	209.2 (215.3)	-0.8	2	33.3%
Craniofacial reconstruction (n=5)	102.0 (75.9)	-108.0	0	0.0%
Thighplasty or Brachioplasty (n=2)	132.5	-77.5	0	0.0%
Total	236.9	26.9	105	22.1%