

Biophysical Journal, Volume 121

Supplemental information

**High turnover and rescue effect of XRCC1 in response to heavy charged
particle radiation**

**Wenjing Liu, Ruqun Wu, Jinlong Guo, Cheng Shen, Jing Zhao, Guangbo Mao, Hongjin
Mou, Lei Zhang, and Guanghua Du**

Supporting Material

High Turnover and Rescue Effect of XRCC1 in Response to Heavy Charged Particle Radiation

Wenjing Liu^{1,2}, Ruqun Wu¹, Jinlong Guo^{1,2}, Cheng Shen^{1,2}, Jing Zhao^{1,2}, Guangbo Mao^{1,3},
Hongjin Mou^{1,3}, Lei Zhang^{1,2}, Guanghua Du^{1,2,*}

¹ Materials research center, Institute of Modern Physics, Lanzhou, Gansu, China

² Institute of Modern Physics, University of Chinese Academy of Sciences, Beijing, Beijing, China

³ School of Materials and Energy, Lanzhou University, Lanzhou, Gansu, China

* Correspondence: gh_du@impcas.ac.cn




Figure S1. The kinetics of XRCC1 protein in HT1080 (XRCC1-RFP) cells exposed to 5.5 MeV/u Kr ions. The recruitment rate constant k_1 is $2.47 \pm 0.29 \times 10^{-2} \text{ s}^{-1}$, and dissociation rate constant k_2 is $1.28 \pm 0.17 \times 10^{-3} \text{ s}^{-1}$. The data represent the measured mean value and the standard error of 6 cells. Ion irradiated is 2.0 ± 0.5 per cell.