

Title: “Thandi should feel embarrassed”: Describing the validity and reliability of a tool to measure depression-related stigma among patients with depressive symptoms in Malawi

Authors: J.M. Dussault¹, C. Zimba², J. Malava³, H. Akello², M.A. Stockton^{4,5}, M. Udedi⁶, B.N. Gaynes^{1,7}, M.C. Hosseinipour^{2,8}, B.W. Pence¹, J. Masiye⁶

¹UNC Gillings School of Global Public Health, Dept of Epidemiology, Chapel Hill, USA; ²UNC Project - Malawi, Lilongwe, Malawi; ³Malawi Epidemiology and Intervention Research Unit, Karonga, Malawi; ⁴Columbia University Vagelos College of Physicians and Surgeons, Dept of Psychiatry, New York, USA; ⁵New York State Psychiatric Institute, New York, USA; ⁶Ministry of Health Malawi, NCDs & Mental Health Unit, Lilongwe, Malawi; ⁷UNC School of Medicine, Dept of Psychiatry, Chapel Hill, USA; ⁸UNC School of Medicine, Division of Infectious Disease, Chapel Hill, USA

Corresponding Author: Josée Dussault, josee.dussault@unc.edu

Supplementary Table 1. Exploratory factor analysis results excluding stigma item #4.^a

Questionnaire Items	Factor 1: Negative Affect	Factor 2: Treatment Carryover	Factor 3: Disclosure Carryover	Alpha
<i>Overall stigma scale</i>				<i>0.76</i>
<i>Negative affect subscale</i>				<i>0.68</i>
1. Being around Thandi would make me feel uncomfortable.	0.58	-0.02	-0.08	
2. Thandi should feel embarrassed about her situation.	0.50	-0.16	0.21	
6. Being around someone diagnosed with depression like Thandi would make me feel uncomfortable.	0.65	0.12	-0.10	
7. Thandi should feel embarrassed about her diagnosis of depression.	0.53	0.10	0.12	
<i>Disclosure carryover subscale</i>				<i>0.74</i>
3. Members of Thandi’s family would be better off if Thandi’s situation was kept secret.	-0.07	0.02	0.69	
8. Members of Thandi’s family would be better off if her diagnosis of depression was kept secret.	0.05	0.05	0.67	
<i>Treatment carryover subscale</i>				<i>0.87</i>
9. If Thandi were going to a clinic regularly to speak with a trained therapist about her problems to help treat her depression, and people found out she was going, she would lose some of her friends.	0.04	0.81	0.04	
10. If Thandi were taking medication to help treat her depression, and people found out she was taking medication, she would lose some of her friends.	0.01	0.81	0.01	

a. This EFA used the principal-factors method and an oblique (promax) rotation, just as Table 2 in the main text.