

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

The association between depressive symptoms and CBT receipt within a psychosis sample.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-051873
Article Type:	Original research
Date Submitted by the Author:	22-Apr-2021
Complete List of Authors:	Mason, Ava; University College London, Division of Psychiatry ; King's College London, Institute of Psychiatry, Psychology and Neuroscience Irving, Jessica; Institute of Psychiatry Psychology and Neuroscience, Psychosis Studies Pritchard, Megan; King's College London, Institute of Psychiatry, Psychology and Neuroscience Sanyal, Jyoti Colling, Craig; South London and Maudsley NHS Foundation Trust, Biomedical Research Centre Chandran, David; King's College London, Institute of Psychiatry Stewart, Robert; King's College London, Institute of Psychiatry
Keywords:	Schizophrenia & psychotic disorders < PSYCHIATRY, Depression & mood disorders < PSYCHIATRY, Adult psychiatry < PSYCHIATRY, Health informatics < BIOTECHNOLOGY & BIOINFORMATICS

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

The association between depressive symptoms and CBT receipt within a psychosis sample.

Ava Mason¹, Jessica Irving¹, Megan Pritchard^{1,2}, Jyoti Sanyal², Craig Colling^{1,2}, David Chandran¹, Robert Stewart^{1,2}.

¹King's College London Institute of Psychiatry, Psychology and Neuroscience, De Crespigny Park, London, SE5 8AF

²South London and Maudsley NHS Foundation Trust, London, UK

* Address correspondence to: Ava Mason, University College London, Gower St, Bloomsbury, London WC1E 6BT, UK. Email: ava.mason.20@ucl.ac.uk.

Keywords: CBTp, psychosis, cognitive behaviour therapy, depression, mental illness.

Abstract word count: 248

Main body word count: 2918

Abstract

Objectives: To examine whether depressive symptoms predict receipt of cognitive behavioural therapy (CBTp) in individuals with psychosis.

Design: Retrospective cross-sectional analysis of electronic health records (EHRs) of a clinical cohort.

Setting: A secondary NHS mental health care service serving four boroughs of south London, UK.

Participants: 20,078 patients diagnosed with an ICD-10 code between F20-29 extracted from an EHR database.

Primary and secondary outcome measures: Primary: Whether recorded depressive symptoms predicted CBTp session receipt, defined as at least one session of CBT for psychosis (CBTp) identified from structured EHR fields supplemented by a natural language processing algorithm. Secondary: Whether age, gender, ethnicity, psychotic symptoms (negative, manic and disorganisation symptoms), a comorbid diagnosis of depression, anxiety or bipolar disorder, general CBT receipt prior to the primary psychosis diagnosis date, or type of psychosis diagnosis predicted CBTp receipt.

Results: Of patients with a psychotic disorder, 8.2% received CBTp. Individuals with at least one depressive symptom recorded and 12 out of 15 of the individual depressive symptoms independently predicted CBTp receipt. Female gender, White ethnicity and presence of a comorbid affective disorder or primary schizoaffective diagnosis were independently positively associated with CBTp receipt within the whole sample and the top 25% of mentioned depressive symptoms.

Conclusions: Individuals with a psychotic disorder who had recorded depressive symptoms were significantly more likely to receive CBTp sessions, aligning with CBTp guidelines of managing depressive symptoms related to a psychotic experience. However, overall receipt of CBTp needs to increase before targeted approaches can be undertaken.

Strengths and limitations of the study

- To our knowledge, this is the first electronic health record (EHR) study to measure how clinical symptomatology predicts CBTp receipt, providing insight on a large sample into whether individuals who may be more in need of CBTp are more likely to have a session
- We replicate previous findings of inequalities in gender and ethnicity in real-world CBTp treatment receipt in a large heterogeneous sample.
- The natural language processing approach allows automated processing of EHR text at scale and can evaluate larger samples than manually conducted case note audits; this could therefore be used more routinely to monitor CBTp receipt.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- This study was limited to a single service provider; however, the results identified themes consistent with previous CBTp provision research in other services.
- Analysing EHRs in this way can identify CBTp receipt but is less suited to investigate whether CBTp is offered or not, or to quantify the quality or focus of the sessions.

For peer review only

Introduction

There are a variety of cognitive and emotional processes involved in the development of psychotic symptoms,¹ with intense distress emerging early on in the course of the disorder. Content of positive symptoms often mirrors the content of depressive thinking processes,² suggesting therapeutic need for individuals experiencing additional depressive symptoms. Specific depressive symptoms that often accompany psychotic disorders are hopelessness, social avoidance and problems in forming relationships.³ Around 50% of patients with psychosis report having experienced suicidal ideation at least once,⁴ and around 40% of individuals with schizophrenia report clinical levels of depression and low self-esteem.⁵ Importantly, individuals report these emotional difficulties and resulting social exclusion to be more debilitating than their psychotic symptoms.⁶ Consequentially, individuals' negative appraisal of their psychotic experiences may lead to loss of social goals and increased shame, predicting later hopelessness and post-psychotic depression.⁷ This comorbid depression increases the likelihood of having a lower quality of life, function, motivation, poorer social relationships, lower medication adherence and relapse to mental health services.^{8,9} Therefore, treatment should focus on the psychotic symptoms and the broader distress they produce, building self-esteem, confidence and a sense of self control and purpose.¹⁰

It is increasingly recognised that medication alone is inadequate for tackling psychosis symptoms.¹¹ In the UK, the National Institute of Clinical Excellence¹² has recommended that cognitive behavioural therapy for psychosis (CBTp) be offered universally to individuals with psychosis. Based on the stress-vulnerability model,¹³ CBTp focuses on distress reduction through targeting negative beliefs and improving self-esteem.¹⁴ Sessions often focus on goal setting and emotional issues such as rebuilding one's self, positivity and acceptance.¹⁰ While CBTp reductions in depressive symptoms are promising, specifically with long term reductions in suicidal behaviour,¹⁵ service provision of this intervention still falls far short of the universal access recommended.¹¹

Considering the impact of targeting these symptoms in CBTp sessions, it is important to monitor receipt of CBTp within psychosis samples. While CBTp provision shows moderate yearly increases (12.8% in 2013 to 14.8% in 2014), the treatment is still only available to a small proportion of individuals,¹¹ short of NICE universal access recommendations.¹² Previous studies investigating CBTp receipt have conducted time-consuming audits on limited sample sizes; these can be affected by under-reporting. On the other hand, the UK's National Mental Health Minimum Data Set report does not require CBT interventions to be recorded in a given individual's record. Natural language processing techniques (NLP)¹⁶ offer the opportunity to extract this information from free text in electronic health records (EHRs) across large numbers of patients with psychosis, and a recent study developed and applied NLP in this respect, finding higher levels of receipt than reported in previous audit, supported by the high positive predictive value and sensitivity of the technique (95% and 96% respectively).¹¹

We investigated whether depressive symptoms predict CBTp receipt in people with psychosis by applying these previously data extraction techniques to secondary mental health care EHRs for a large South London catchment population. Secondary predictors of receipt were type of psychosis diagnosis (schizophrenia, schizoaffective disorder or other schizophrenia spectrum disorder), psychotic symptoms (negative, manic or disorganisation), general CBT receipt prior to psychosis diagnosis, comorbid depression, anxiety or bipolar diagnosis and socio-demographic factors (ethnicity, gender and age).

Methods

For this study, we extracted data on individuals with a diagnosis of a recognised schizophrenia spectrum diagnosis from the case registry of the South London and Maudsley NHS Foundation Trust (SLaM). This is a large secondary care mental healthcare provider, serving around 1.3 million residents in Croydon, Lambeth, Lewisham and Southwark. EHRs have been used for all SLaM services since 2006, with the Clinical Record Interactive Search system (CRIS) being established in 2008 to facilitate the retrieval of de-identified data from these records of patients previously or currently receiving mental healthcare from SLaM.¹⁷ The source EHR contains unstructured free text fields from correspondence, personal histories, mental health examinations and management plans, as well as structured fields for coding demographic information, like age and ethnicity. Implementing data from all these fields reduces selection bias of utilising only specific sources of information from the EHR. Consequently, a large programme of work has developed a range of NLP algorithms over the last decade, whose detailed descriptions and performance data are contained in an open-access catalogue.¹⁸ CRIS has approval as a data resource for secondary analysis (Oxford Research Ethics Committee C, reference 18/SC/0372), and a service user-led committee considers all proposed research before access to CRIS data is granted.

We extracted data for all individuals receiving SLaM care between January 2007 and June 2020 with a primary diagnosis of an ICD-10-defined schizophrenia spectrum disorder (F20-F29) and above the age of 18 at the time their original referral was accepted. The index date for covariate definitions was the date of the first diagnosis within this grouping. Individuals may have been active within the service before their index date, allowing us to extract data on prior CBT receipt. The sample was restricted to those with data on all variables.

Ethnicity, age at referral and gender were also extracted. Ethnicity was categorised into six groups for analysis: 'White British' (British), 'White other' (Irish or any other white background), 'Black' (Caribbean, African or any other black background), 'Asian' (Indian, Bangladeshi, Pakistani, Chinese or any there Asian background), 'Other/mixed' (white and Asian, white and black Caribbean, white and black African, any other ethnic group) and 'Not stated'.

Diagnosis was categorised into three subgroups of schizophrenia (ICD-10 codes F20.0–F20.9), schizoaffective disorder (F25.0–F25.9) and other schizophrenia spectrum disorder (F21, F22.0–F22.9, F23.0–F23.9, F24, F28 and F29). Within the data collection period, secondary diagnosis of depression (ICD-10: F32 or F33), anxiety (ICD-10: F40 or F41), or bipolar disorder (ICD-10: F31) were also extracted from structured field data.

NLP algorithms for each specific symptom were used to identify recorded depressive and psychotic symptoms within participants. Symptoms were categorised as depressive, positive, negative, manic or disorganisation. These symptoms had been categorised *a priori* by developers of the original independent symptom NLP algorithms. As symptoms could be labelled in more than one category during analysis, multicollinearity tests using the R function `vif()` within the [car package] were undertaken to avoid issues with overlapping predictor variables. Positive symptoms were excluded from the regression analysis due to multicollinearity affecting results, all other variables were included due to their VIF values being well below five. The overall symptom list and subsequent recoding can be found in Table 1. Presence of at least one mention of any symptom in the five categories was computed as a binary variable (0/1). This was prioritised over calculating the frequency of mentions, as the number of symptoms in each of the categories was unevenly distributed (e.g. 15 depressive symptoms vs. 6 manic symptoms).

Patient and public involvement

The Clinical Record Interactive system as a data resource was developed and is run with extensive patient involvement. However, this particular analysis did not involve patients in its design or implementation.

Table 1
Classification of symptom predictors

Symptom	Symptom label
Aggression	Positive
Agitation	Positive
Anergia	Depressive /Negative
Anhedonia	Depressive/Negative
Apathy	Depressive/Negative
Arousal	Manic
Blunted affect	Depressive/Negative
Circumstantiality	Disorganisation
Delusions	Positive
Derailment	Disorganisation
Disturbed sleep	Depressive/Manic
Elation	Manic
Emotional Withdrawal	Negative
Flight of ideas	Disorganisation
Formal thought disorder	Disorganisation
Grandiosity	Manic
Guilt	Depressive
Hallucinations (auditory)	Positive
Helplessness	Depressive
Hopelessness	Depressive
Hostility	Positive
Insomnia	Depressive/Manic
Irritability	Manic
Paranoia	Positive
Persecutory ideation	Positive
Poor appetite	Depressive
Poor concentration	Depressive
Poor motivation	Depressive
Poverty of speech	Negative
Poverty of thought	Negative
Social withdrawal	Negative
Suicidal ideation	Depressive
Tangentiality	Disorganisation
Tearfulness	Depressive
Thought block	Disorganisation
Worthlessness	Depressive

The date of the first and last general CBT session before the index date was extracted. This was coded as a binary variable, with individuals in the 'Prior CBT' receipt group having at least one session date mention prior to their index date. This was included as a predictor to adjust for previous experience of the specific CBT intervention. Mentions were extracted using the same NLP tool as the CBTp outcome measure mentioned subsequently.

1
2
3 The primary outcome was CBTp receipt, identified using a combination of structured fields and NLP.¹⁶ The NLP
4 algorithms for general CBT has high PPV and sensitivity,¹¹ consistent with other NLP algorithms such as
5 medication dose and diagnosis.¹⁹ The date of the first CBTp session on or after the index date was extracted
6 and computed as a binary variable, so that individuals in the 'CBTp receipt' group had at least one CBTp
7 session mention after the index date.
8

9 **Statistical analysis**

10 To avoid overfitting, we followed the 'one in ten' rule, whereby one predictor can be measured for every 10
11 events. As the data included 1647 CBTp events, our study was able to include all 12 predictors within the same
12 regression model.
13

14 All statistical analyses were conducted using R (version 1.3.9). Descriptive statistics for demographic and
15 clinical variables are reported as frequencies for categorical variables and means and standard deviations for
16 the continuous variable (age at referral). Chi square tests were also calculated for categorical variables, and t-
17 test for age to measure between-group differences in those with/without CBT receipt.
18

19 Binary logistic regression was used to examine the association between depressive symptoms and receipt of at
20 least one CBT session in the whole sample. For this, three regression models were analysed. Model 1 was an
21 unadjusted model with only depressive symptoms as the predictor variable. Due to significant provision
22 differences seen in previous CBTp studies,¹¹ model 2 (partially adjusted model), adjusted for sociodemographic
23 variables (age at referral, ethnicity, gender), primary diagnosis group and presence of a comorbid diagnosis
24 (anxiety, depression and bipolar disorder). Model 3 (fully adjusted model) also adjusted for prior CBT receipt
25 before the index date (first psychosis diagnosis date) and psychotic symptoms mention (manic, negative and
26 disorganisation symptoms). Positive psychotic symptoms were not included in these models, as individuals all
27 had at least one mention within their case notes.
28

29 As the primary aim of the study was to investigate depressive symptoms as a predictor of CBTp receipt, we
30 also split the depressive symptoms category into the 15 specific depressive symptoms applications. Model 4
31 was an unadjusted model with the 15 symptoms as predictor variables. Model 5 was a fully adjusted model
32 that adjusted for all the variables in Model 3. We also conducted a sensitivity analysis to investigate how
33 results were affected by overlap of negative or depressive symptom annotations, by removing negative
34 symptoms as a predictor from the logistic regression model.
35

36 Lastly, to compare differences in the general sample with those with the top 25% quantity for depressive
37 symptoms, we conducted two further regression models. Model 6 partially adjusted for socio-demographic
38 factors, diagnostic group and comorbid diagnosis and Model 7 fully adjusted for prior CBT, negative and
39 disorganisation symptoms additionally. This group all had at least one manic and psychotic symptom, so these
40 variables were not included in the model.
41

42 **Results**

43 **Participants**

44 The cohort comprised 20,078 individuals with the inclusion diagnoses, 1647 (8.2%) of whom received at least
45 one session of CBTp after their first diagnosis date. The mean age of the cohort was 42.4 years (SD=16.5).
46 Distribution frequencies for all categorical variables can be found in Table 2. Chi-square test results
47 represented in this table compared those with or without CBTp receipt. All mentioned variables showed
48 significant between-group differences at $p < .001$ apart from gender ($X^2=2.75$, $p=.097$). Additionally, the Welch
49 two sample t-test found significant between-group differences in age ($t=15.34$, $p < .01$). Where those who had
50 received CBTp had a lower mean age ($M=33.12$ $SD= 11.5$) compared to those who did not ($M=35.88$,
51 $SD=13.08$). The significant results confirmed the need for further analysis through the regression models.
52

Table 2*Distribution frequencies on baseline demographics and diagnoses split by CBTp receipt and primary diagnosis group.*

	No CBTp delivery (n = 18431)	CBTp delivery (n=1647)	Chi square tests (χ^2)
Ethnicity			$\chi^2=100.57^{***}$
White British	30% (5516/18431)	32.8% (540/1647)	
White Other	10.4% (1908/18431)	8.5% (140/1647)	
Black	36.5% (6719/18431)	41.7% (687/1647)	
Asian	6.5% (1193/18431)	5.2% (86/1647)	
Other/Mixed	9.8% (1808/18431)	10.5% (173/1647)	
Not stated	7.0% (1287/18431)	1.3% (21/1647)	
Gender			$\chi^2= 2.75$
Female	41.4% (7636/18431)	43.5% (717/1647)	
Male	58.6% (10795/18431)	56.5% (930/1647)	
Bipolar diagnosis	4.4% (810/18431)	9.0% (149/1647)	$\chi^2=71.94^{***}$
No biop diagnosis	95.6% (17621/18431)	91.0% (1498/1647)	
Depression diagnosis	7.4% (1373/18431)	14.0% (230/1647)	$\chi^2=87.36^{***}$
f32 no diagnosis	92.6% (17058/18431)	86.0% (1417/1647)	
Anxiety diagnosis	2.4% (441/18431)	7.0% (115/1647)	$\chi^2=118.28^{***}$
f40/41 no diagnosis	97.6% (17990/18431)	93.0% (1532/1647)	
Prior CBT	3.1% (573/18431)	14.4% (237/1647)	$\chi^2=497^{***}$
No prior cbt	96.9% (17858/18431)	85.6% (1410/1647)	

***p<.001

Positive psychotic symptoms were excluded from chi square and regression analysis, as all patients had at least one positive psychotic symptom.

General depressive symptom regression analysis

Results from the unadjusted (model 1), partially adjusted (model 2) and fully adjusted regression (model 3) are displayed in Table 3. Regression model 1 found that general mention of at least one of 15 potential depressive symptoms significantly predicted CBTp receipt. Regarding model 2 and 3, individuals with at least one depressive, negative or disorganisation symptom mention, being of female gender, white ethnicity, prior CBT receipt and presence of a comorbid affective disorder independently positively associated with CBTp receipt.

Table 3

Unadjusted, partially and fully adjusted logistic regression models for CBTp receipt (Regression model 1,2 and 3).

		Unadjusted	Partially adjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)	OR(95%CI)
Depressive symptoms				
1+ depressive symptom mention	18286(91.1)	44.13(17-118)***	30.7(11.51-82.30)***	7.37(2.66-20.42)***
Bipolar diagnosis				
Has f31 diagnosis	959(4.80)		1.72(1.43-2.09)***	1.38(1.13-1.68)***
Depression diagnosis				
Has f32 diagnosis	603(80)		1.67(1.43-1.96)***	1.39(1.18-1.63)**
Anxiety diagnosis				
Has f40/41 diagnosis	556(2.80)		2.43(1.95-3.03)***	2.07(1.65-2.61)***
Age	N/A		0.034(0.03-0.04)***	0.028(0.024-0.03)***
Gender				
Male			Reference category	
Female	8353(41.60)		1.22(1.10-1.36)***	1.23(1.10-1.38)***
Ethnic group				
White British	6056(30.10)		Reference category	
White Other	2048(10.20)		0.67(0.55-0.81)***	0.69(0.57-0.85)***
Black	7406(36.90)		0.85(0.75-0.96)*	0.79(0.7-0.90)*
Asian	1279(6.40)		0.61(0.48-0.78)***	0.60(0.47-0.77)***
Other/Mixed	1981(9.90)		0.81(0.67-0.97)*	0.83(0.7-1.01)*
Not Stated	1308(6.50)		0.17(0.11-0.27)***	0.22(0.14-0.34)***
Primary diagnosis				
Schizophrenia	9845(49.00)		Reference category	
Schizoaffective disorder	2142(10.70)		1.04(0.88-1.24)	1.01(0.85-0.21)
Other schizophrenia spectrum	8091(40.30)		0.9(0.81-1.02)	0.98(0.87-1.01)
Negative symptoms				
1+ Negative symptom mention	13169(65.60)			2.12(1.80-2.50)***
Manic symptoms				
1+ Manic symptom mention	17945(89.40)			3.46(1.95-6.15)***
Disorganisation symptoms				
1+ Disorganisation symptom mention	11513(57.30)			1.36(1.20-1.55)***
CBT prior				
1+ prior CBT session				3.65(3.08-4.32)***

*p<.05, **p<.01, ***p<.001

Unadjusted (model 1): depressive symptom as a predictor with no adjusted covariates

Partially adjusted (model 2): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, depressive symptoms.

Fully adjusted (model 3): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, depressive symptoms, prior CBT, negative symptoms, disorganisation symptoms, manic symptoms.

Regression analysis with individual depressive symptoms

Results from the unadjusted (model 4) and fully adjusted (model 5) regression analyses for each of the 15 individual depressive symptoms are displayed in Table 4. Each symptom refers to presence of at least one mention in the patients notes. While all variables were significant in the unadjusted model at $p < .001$, the fully adjusted model reduced the significance of suicide ideation ($p < .01$) and disturbed sleep ($p < .01$), with anhedonia, anergia, apathy, and blunted affect becoming non-significant ($p > 0.05$).

Table 4

Unadjusted and fully adjusted logistic regression models for CBTp receipt with individual depressive symptoms as covariates (Regression model 4 and 5).

		Unadjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)
Hopelessness	7345(36.60)	4.81(4.3-5.40)***	1.45(1.26-1.66)
Helplessness	3124(15.60)	4.03(3.62-4.50)***	1.55(1.37-1.76)***
Suicide ideation	9451(47.10)	4.11(3.66-4.63)***	1.25(1.09-1.44)**
Poor appetite	8044(40.10)	3.31(2.97-3.68)***	1.28(1.13-1.45)***
Poor motivation	8630(43.00)	4.34(3.87-4.86)***	1.43(1.24-1.64)***
Insomnia	6870(34.20)	3.74(3.35-4.15)***	1.4(1.24-1.58)***
Disturbed sleep	16667(83.00)	15.3(10.16-22.8)***	2.76(1.5-5.08)**
Poor concentration	12289(61.20)	8.16(6.81-9.77)***	2.33(1.9-2.85)***
Anhedonia	4047(20.20)	2.9(2.61-3.22)***	0.97(0.85-1.10)
Anergia	873(43.50)	2.63(2.20-3.15)***	0.98(0.80-1.20)
Apathy	4149(20.70)	2.21(1.98-2.46)***	0.93(0.82-1.05)
Guilt	8178(40.70)	4.6(4.1-5.15)***	1.49(1.30-1.70)***
Tearfulness	10951(54.50)	3.87(3.41-4.39)***	1.22(1.05-1.42)**
Blunted affect	6889(34.30)	2.66(2.41-2.95)***	0.91(0.80-1.03)
Worthlessness	2921(14.50)	3.94(3.53-4.40)***	1.37(1.21-1.56)***

* $p < .05$, ** $p < .01$, *** $p < .001$

Sensitivity analysis

The non-significant results of certain depressive symptoms (anhedonia and anergia) may have been due to their inclusion within the negative symptom category, causing over-adjustment of the model. To test this, sensitivity analysis was conducted, where the fully adjusted regression (model 3) did not include negative symptoms as a covariate. While all significant variables remained significant, non-significant results for anhedonia and apathy were still found. Therefore, we report the fully adjusted model with negative symptoms as a variable for both grouped and individual depressive symptom associations.

Depressive symptom regression analysis within the top 25% number of depressive symptoms.

This sample comprised individuals with the top 25% number of depressive symptoms (5018 patients), defined to reflect those who might reasonably expect to receive CBT. The sample characteristics and regression analysis can be seen in Table 5. Results from the partially adjusted (model 6) and fully adjusted regression (model 7) are displayed in Table 5. Regression model 6 found that general mention of at least one of 15 potential depressive symptoms significantly predicted CBTp receipt. Regarding table 3, we found that individuals with at least one depressive, negative or disorganisation symptom mention, being of female

gender, white ethnicity, prior CBT receipt and presence of a comorbid affective disorder were positively associated with CBTp receipt.

Table 5

Partially and fully adjusted logistic regression models for CBTp receipt with individual depressive symptoms as covariates within top 25% quantity of depressive symptoms (Regression model 6 and 7).

		Partially adjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)
Bipolar diagnosis			
Has f31 diagnosis	541(10.78)	1.21(0.97-1.51)*	1.16(0.92-1.46)
Depression diagnosis			
Has f32 diagnosis	885(17.63)	1.11(0.93-1.35)	1.08(0.90-1.30)
Anxiety diagnosis			
Has f40/41 diagnosis	270(5.38)	1.70(1.29-2.24)****	1.61(1.21-2.13)***
Age			
M=	36.24(18-93)	0.98(0.97-0.99)****	0.98(0.97-0.99)****
Gender			
Male	2058(41.01)	Reference category	
Female	2960(58.99)	0.82(0.71-0.95)***	0.85(0.73-0.98)**
Ethnic group			
White British	1485(29.59)	Reference category	
White Other	433(8.63)	0.81(0.62-1.06)	0.80(0.61-1.05)
Black	2262(45.08)	0.73(0.62-0.86)****	0.74(0.63-0.87)****
Asian	328(6.53)	0.59(0.43-0.82)***	0.60(0.43-0.83)***
Other/Mixed	467(9.31)	0.93(0.72-1.19)	0.92(0.72-1.19)
Not Stated	43(0.86)	0.24(0.07-0.79)**	0.26(0.80-0.86)**
Primary diagnosis			
Schizophrenia	2219(44.22)	Reference category	
Schizoaffective disorder	740(14.72)	1.05(1.84-1.30)	1.00(0.80-1.25)
Other schizophrenia spectrum	2059(41.03)	1.02(0.87-1.19)	1.05(0.89-1.23)
Negative symptoms			
1+ Negative symptom mention	3744(74.61)		0.41(0.24-0.70)***
Disorganisation symptoms			
1+ Disorganisation symptom mention	3209(63.94)		1.19(0.97-1.46)*
CBT prior			
1+ prior CBT session	360(7.17)		2.45(1.98-3.04)****

*p<.05, **p<.01, ***p<.001, ****p<.000

Partially adjusted (model): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis.

Fully adjusted (model): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, prior CBT, negative symptoms, disorganisation symptoms.

Discussion

We believe that this is the first study to examine the relationship between clinical symptomatology and CBTp receipt within a sample of people with psychosis in a naturalistic community setting. In general, only 8.2% of individuals received CBTp, despite 91% having at least one depressive symptom recorded. Individuals with at least one depressive symptom mention were 7.37 times more likely to have at least one CBTp session in the fully adjusted model (table 3), suggesting that the minority who don't present with any depressive symptoms are very unlikely to receive CBTp. This could possibly be due to clinicians tending to cite a depressive symptom when referring an individual with psychosis to psychotherapy. In the sample of those with the highest number of depressive symptoms (top 25%), relationships between CBTp receipt and comorbid anxiety diagnosis, age, gender, ethnicity, prior CBT, negative and disorganised psychotic symptoms remained (effect size ranging from 0.26-2.45).

Overall there was therefore a low prevalence of CBTp receipt within those with one depressive symptom. The depressive symptom which was the strongest predictor of this intervention in fully adjusted models was disturbed sleep. There is a known high prevalence of sleeping problems in this population,^{20,21} described by some researchers as

1
2
3 an 'intrinsic feature of schizophrenia,²² known to reduce quality of life, decreasing coping and exacerbate positive
4 symptoms.²³ Furthering this, the recommended first line of treatment for sleep problems in this sample is CBT.²⁴ Poor
5 concentration was the next strongest depressive symptom predictor in the fully-adjusted model, supporting previous
6 research of its association with psychosis vulnerability.²⁵ The significance of helplessness, guilt and hopelessness
7 mirrors CBTp research that found significant post-treatment reduction in hopelessness, self-depreciation and guilt using
8 the Calgary Depression Rating Scale for Schizophrenia.²⁶ Other significant depressive symptoms associated with low
9 self-esteem and negative self-evaluation and emotions have been found to significantly affect the development and
10 severity of positive symptoms.²⁷ This may be because positive symptoms develop as a psychological defence against
11 low self-esteem²⁸ and depression-induced guilt.²⁹ Therefore, it can be said that the significance of each of the
12 depressive symptoms is often linked to psychotic symptoms and CBTp effectiveness. However, general results suggest
13 that receipt of this intervention requires an increase for all of this population before individuals with these specific
14 symptoms could be targeted.

15
16
17 Regarding negative symptoms, the contrasting results suggest multiple theories that require specific testing. Further
18 testing was not conducted in the current study due to the primary aim focusing on depressive symptoms. However,
19 from our results on specific depressive symptoms in table 4, symptoms that overlapped with negative symptoms
20 (anhedonia, anergia, apathy and blunted affect) were not associated with CBTp receipt. This raises concerns, suggesting
21 that individuals with these specific negative/depressive symptoms are no more likely to receive CBTp than someone
22 without these symptoms. Possibly, this is due to clinicians not tending to refer these individuals because they don't
23 believe intervention would be effective. This is in line with a CBTp review of randomised control trials, finding non-
24 significant reductions of negative symptoms,³⁰ perhaps due to the narrowing of treatments to specifically target
25 positive symptoms.³¹ However, further work should be undertaken to verify that individuals are not being denied a
26 potentially beneficial intervention because of their symptom profile.

27
28 Prior CBT receipt, comorbid disorder presence and specific psychotic symptoms (manic, disorganised and negative) also
29 emerged as independent predictors of CBTp receipt for the general sample and within those with the top 25%
30 depressive symptom numbers. Within table 3, individuals who had any recorded CBT receipt prior to the index date
31 were nearly four times more likely to have recorded CBTp receipt later on. Also, patients with an additional comorbid
32 disorder were 1.38-2.07 times more likely to have received CBTp compared to those with just a psychosis diagnosis.
33 Within table 5, individuals with prior CBT were 2.45 times more likely to receive CBTp, those with anxiety were 1.61
34 times more likely and those with disorganised or negative symptoms were more 1.19 and 0.41 times more likely
35 respectively. After general CBTp receipt has increased, there could be a method to focus more on patients with
36 different types of psychotic symptoms and comorbid affective diagnosis. Furthering this, future research could
37 investigate whether those who have had prior general CBT would benefit from CBTp, or whether those who have not
38 had any experience developing cognitive behaviour skills in therapy should be targeted.

39
40 Crucially, there were also significant differences in CBTp receipt between different ethnic and gender groups. Male
41 patients were 1.23 times less likely in the general sample and 0.85 times less likely in the top 25% of depressive
42 symptoms to have recorded CBTp receipt. Black, Asian, Other and Mixed ethnic groups were between 0.60 to 0.70
43 times as likely to have a documented CBTp session compared to individuals of white ethnicity within both the general
44 and top 25% depressive symptoms samples. Inequitable access to CBTp has been identified in previous CBTp research
45 within a psychosis sample drawn from the same data resource in 2017, finding female patients to be more likely to
46 have received CBTp and individuals of White ethnicity to have a significantly higher likelihood of CBT receipt than Black
47 or other ethnicity groups.¹¹ This also supports results from a recent CBTp study focusing specifically on ethnic group
48 differences in CBTp provision within SLAM, who found that in comparison to White British individuals, those from Black
49 ethnic groups with psychosis or bipolar disorder were significantly less likely to have a documented CBTp session.³³
50 Inequality in CBTp receipt may be due to ethnic variations in CBTp engagement. Some of these barriers within certain
51 communities may be increased stigma, fear of clinicians by service-users or service users by clinicians, institutional
52 racism within mental health services, or non-culturally appropriate therapy.³³ As differences in documented CBTp
53 receipt between ethnic groups have now been documented by three different papers in this service, it is imperative
54 that further work is conducted to increase provision of CBTp within groups less likely to receive treatment. This may
55 include targeted outreach programs and culturally adapting interventions³⁴ within these minority groups.

56
57 The present study has a number of strengths and limitations. Generally, focusing on patients with more
58 diverse functioning, comorbidity and symptom severity levels helps research identify a larger number of
59 predictors of clinical outcomes. This can be seen through our results, where negative, manic and
60

1
2
3 disorganisation symptoms significantly predicted CBTp receipt, as well as recent heterogenous research,³¹
4 that was the first to identify depression as a significant predictor of positive symptom improvement post-
5 CBTp. This highlights the importance of focusing on a clinically heterogeneous sample to realistically determine
6 significant predictors of CBTp receipt. Secondly, using an NLP approach automates the measurement of what
7 would otherwise require manually conducted audits on records and case notes, increasing the number of
8 cases that can be investigated and providing a method that could be used more routinely to monitor CBT
9 receipt. The large sample size enabled us to identify clinical differences in the real-life administration of CBTp
10 within a psychosis cohort, and we were able to adjust for multiple clinical variables and comorbidity diagnoses
11 to provide a more realistic understanding of the depressive symptom-CBTp receipt relationship. This time
12 frame was broad to allow the inclusion of as many active patients receiving CBTp as possible, additionally
13 circumventing monthly/seasonal variation of CBTp receipt.
14

15 One limitation of the study was the omission of strict time periods for the mention of clinical symptoms prior
16 to CBTp administration. Unfortunately, using this approach would have involved implementing time periods on
17 all of the other clinical symptoms and variables, which would have been difficult considering the number of
18 variables that would need to be controlled. In addition, the NLP symptom algorithms do not currently
19 distinguish between past or present symptoms. Therefore, symptom mentions documented after the CBTp
20 receipt date could refer to mentions of symptoms occurring prior to CBTp receipt, reducing the effectiveness
21 of using time periods. Additionally, a follow-up time period after the index date was not established, meaning
22 that participants included in the cohort at a later date may have been less likely to have had a CBTp session,
23 due to their limited time period within the service.
24

25 While our use of additional querying of text fields allowed us to identify a significantly larger number of CBTp
26 episodes than using structured data alone, we were not able to quantify the gap between CBTp referral and
27 CBTp receipt. This is because the CBTp NLP algorithm detects CBTp receipt rather than CBTp being offered,
28 due to the wide range of subtle wording used for the latter more complex entity. Therefore, the results
29 combine effects on the likelihood of CBTp being offered, with those on session receipt following an offer.
30 While this may have affected our results, previous service audits have suggested that the severity and
31 occurrence of depressive symptoms significantly decreases CBT receipt.³⁴ Therefore, if only receipt was
32 directly measured, we would expect to see similar results.
33

34 **Future directions**

35
36 Initiatives such as the Improving Access to Psychological Therapies programme for serious mental illness, early
37 intervention access and projects to decrease waiting times for referral have been developed to target this
38 clinical population. However, access still falls short of recommendations and is inequitable for specific
39 psychotic diagnoses, age and ethnicity.¹¹ Therefore, given the effect of CBTp on depressive symptoms, perhaps
40 its more pragmatic to focus on patients with additional depressive symptoms. Monitoring CBTp receipt over
41 time could decipher whether these initiatives are effective at increasing general access for those with
42 psychosis, and specific access for different sociodemographic groups and those with additional depressive
43 symptoms (who may benefit the most).
44

45
46 The significant secondary clinical and sociodemographic variables require further analysis in order to fully
47 understand the services' provision. This could involve attention given to the independent psychotic symptoms
48 within the negative, manic and disorganisation categories in a similar manner to the specific depressive
49 symptom regression models analysed. Further research could also explore why the presence of co-morbid
50 anxiety and bipolar disorder in this sample predicted CBTp receipt. Additionally, the results suggest a need to
51 reflect on the steps taken since the previous service study,³³ regarding inequality in CBTp receipt among
52 gender and ethnic groups, due to the consistent significant results seen. Regarding the use of EHR data, future
53 work could involve developing a separate NLP algorithm to ascertain the offering of CBTp or provide another
54 structured field for clinicians to complete for this. However, additional text fields seem an unlikely approach,
55 as clinicians prioritise text field data for communication about CBTp sessions for themselves and their
56 colleagues rather than to collect structure data for the sake of research. Therefore, as previously suggested,¹¹
57 it is important to accept the mixed structured-text field approach that will remain in healthcare record data
58 and perhaps our time is best spent in improving NLP algorithms to detect the subtleties of intervention and
59 clinical outcome data. However, the implications of our results and their consistency three years after the first
60

1
2
3 CBTp service paper suggest the need to use this or future algorithms for service monitoring independent of
4 these improvements.
5

7 **Acknowledgments**

8 This paper would not have been completed without the assistance from Dr Rob Stewart. I would also like to
9 thank Jessica Irving for her assistance in the statistical analysis and general manuscript.
10

11 **Contributorship statement**

12 Ava Mason planned the protocol, analysed the data and wrote the manuscript. Dr Rob Stewart provided
13 access to the data, looked over edits and revisions on the manuscript. The data from this paper was accessed
14 through the assistance of Megan Pritchard and Jyoti Sanyal. The applications used to obtain data specific to
15 CBTp were developed by Craig Colling and David Chandran. Jessica Irving assisted in statistical analysis.
16
17

18 **Competing interests**

19
20
21 The authors report no conflict of interest with respect to the findings described in this manuscript. RS declares
22 research support received in the last 36 months from Janssen, GSK and Takeda.
23

24 **Funding**

25
26
27 The independent research of this paper is funded by the National Institute for Health Research (NIHR)
28 Biomedical Research Centre at the South London and Maudsley NHS Foundation Trust and King's College
29 London. RS is part-funded by: i) the National Institute for Health Research (NIHR) Biomedical Research Centre
30 at the South London and Maudsley NHS Foundation Trust and King's College London; ii) a Medical Research
31 Council (MRC) Mental Health Data Pathfinder Award to King's College London; iii) an NIHR Senior Investigator
32 Award; iv) the National Institute for Health Research (NIHR) Applied Research Collaboration South London
33 (NIHR ARC South London) at King's College Hospital NHS Foundation Trust. The views expressed are those of
34 the authors and not necessarily those of the NIHR or the Department of Health and Social Care.
35
36

37
38 The Corresponding Author (AM) has the right to grant on behalf of all authors and does grant on behalf of all
39 authors, an exclusive licence (or non-exclusive for government employees) on a worldwide basis to the BMJ
40 Publishing Group Ltd to permit this article (if accepted) to be published in BMJ editions and any other BMJPL
41 products and sublicences such use and exploit all subsidiary rights, as set out in our licence.
42

43 **Data Sharing Statement**

44 Due to the terms of Ethics and Information Governance approvals and clinical source of the data, CRIS datasets
45 must remain within the South London and Maudsley NHS Foundation Trust (SLaM) firewall. All data used from
46 this study can be made accessible on request from cris.administrator@slam.nhs.uk, subject to the setting up of
47 an appropriate research passport or SLaM honorary contract.
48
49

50 **Transparency declaration**

51
52 The lead author AM affirms that the manuscript is an honest, accurate, and transparent account of the study
53 being reported; that no important aspects of the study have been omitted; and that any discrepancies from
54 the study as planned (and, if relevant, registered) have been explained.
55
56
57
58
59
60

References

1. Chadwick P, Birchwood M. The omnipotence of voices. A cognitive approach to auditory hallucinations. *Br J Psychiatry* 1994;164:190–201.
2. Fowler D. Cognitive Behavior Therapy for Psychosis: From Understanding to Treatment. *Am J Psychiatr Rehabil* 2000;4(2):199–215.
3. Hazell CM, Hayward M, Cavanagh K et al. Clara Strauss A systematic review and meta-analysis of low intensity CBT for psychosis. *Clinical Psychology Review* 2016;45:183–192.
4. Hawton, K, Sutton, L, Haw, C, Sinclair, J, Deeks, JJ. Schizophrenia and suicide: systematic review of risk factors. *Br. J. Psychiatry* 2005;187:9–20.
5. Sands JR, Harrow M. Depression during the longitudinal course of schizophrenia. *Schizophr Bull* 1999;25:157–171.
6. Greenwood KE, Sweeney A, Williams S, et al. CHOICE of Outcome In Cbt for psychosEs (CHOICE): The Development of a New Service User–Led Outcome Measure of CBT for Psychosis. *Schizophrenia Bulletin* 2010;36(1):126–135.
7. Birchwood M, Iqbal Z, Chadwick P, et al. Cognitive approach to depression and suicidal thinking in psychosis. I: Ontogeny of post-psychotic depression. *Br J Psychiatry* 2000;177: 516–528.
8. Conley RR. The burden of depressive symptoms in people with schizophrenia. *Psychiatr Clin North Am* 2009;32:853–861.
9. Rocca P, Bellino S, Calvarese P, et al. Depressive and negative symptoms in schizophrenia: different effects on clinical features. *Comprehensive Psychiatry* 2005; 46: 304–310.
10. Pitt L, Kilbride M, Nothard S, et al. Researching recovery from psychosis: a user-led project. *Psychiatr Bull R Coll Psychiatr* 2007;31:55–60.
11. Colling C, Evans L, Broadbent M, et al. Identification of the delivery of cognitive behavioural therapy for psychosis (CBTp) using a cross-sectional sample from electronic health records and open-text information in a large UK-based mental health case register. *BMJ Open* 2017;7:e015297.
12. National Institute for Health and Clinical Excellence. Schizophrenia: Core interventions in the treatment and management of schizophrenia in primary and secondary care. London, UK: NICE Clinical Guideline 2009.
13. Rathod S, Phiri P, Kingdon D. Cognitive behavioral therapy for schizophrenia. *Psychiatr Clin North Am* 2010;33:527–536.
14. Gumley A, Schwannauer M. Staying Well after Psychosis: A Cognitive Interpersonal Approach to Recovery and Relapse Prevention. New York, NY: John Wiley & Sons Ltd 2006.
15. Lincoln TM, Rief W, Wetermann S, et al. Who stays, who benefits? Predicting dropout and change in cognitive behaviour therapy for psychosis. *Psychiatry Research* 2014;216(2):198–205.
16. Spyns P. Natural language processing in medicine: an overview. *Methods Inf Med* 1997;35(4-5):285–301.
17. Stewart R, Soremekun M, Perera G, et al. The South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLAM BRC) case register: development and descriptive data. *BMC Psychiatry* 2009;9:51.
18. (<https://www.maudsleybrc.nhr.ac.uk/media/325736/applications-library-v13.pdf>).
19. Hayes RD, Downs J, Chang CK, et al. The effect of clozapine on premature mortality: an assessment of clinical monitoring and other potential confounders. *Schizophr Bull* 2015;41:644–55.
20. Chadwick PDJ, Birchwood MJ, Trower, P. Cognitive Therapy for Delusions, Voices and Paranoia: Towards A Person Model. Chichester: Wiley 1996.
21. Chiu V, Ree M, Janca A, et al. Sleep profiles and CBT-I response in schizophrenia and related psychoses. *Psychiatry Research* 2018;268:279–287.
22. Chouinard S, Poulin J, Stip E, et al. Sleep in untreated patients with schizophrenia: a meta-analysis. *Schizophr Bull* 2004;30:957–967.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
23. Waite F, Evans N, Myers E, et al. The patient experience of sleep problems and their treatment in the context of current delusions and hallucinations. *Psychology and Psychotherapy: Theory, Research and Practice* 2016;89:181–193.
 24. Mitchell MD, Gehrman P, Perlis M, et al. Comparative effectiveness of cognitive behavioral therapy for insomnia: A systematic review. *BMC Family Practice* 2012;13(1):40.
 25. Bogren M, Mattisson C, Tambs K, et al. Predictors of psychosis: a 50-year follow-up of the Lundby population. *Eur Arch Psychiatry Clin Neurosci* 2009;260:113-125.
 26. Muller MJ, Marx-Dannigkeit P, Schlösser R, et al. The Calgary Depression Rating Scale for Schizophrenia: Development and interrater reliability of a German version (CDSS–G). *J Psychiatr Res* 1999;33:433–443.
 27. Smith B, Fowler DG, Freeman D, et al. Emotion and psychosis: links between depression, self-esteem, negative schematic beliefs and delusions and hallucinations. *Schizophr Res* 2006;86(1-3):181-188.
 28. Bentall RP, Kinderman P, Kaney S. The self, attribution processes and abnormal beliefs: Towards a model of persecutory delusions. *Behav Res Ther* 1994;32:331-341.
 29. Lake CR. Hypothesis: Grandiosity and Guilt Cause Paranoia; Paranoid Schizophrenia is a Psychotic Mood Disorder; a Review. *Schizophr Bull* 2008;34(6):1151–1162.
 30. Lincoln TM, Ziegler M, Mehl S, et al. Moving from efficacy to effectiveness in cognitive behavioral therapy for psychosis: a randomized clinical practice trial. *J Consult Clin Psychol* 2012;80(4):674-686.
 31. Penn DL, Meyer PS, Evans E, et al. A randomized controlled trial of group cognitive-behavioral therapy vs. enhanced supportive therapy for auditory hallucinations. *Schizophr Res* 2009;109:52–59.
 32. Morris R, Sellwood W, Edge D, et al. Ethnicity and impact on the receipt of Cognitive Behavioural Therapy in people with psychosis or bipolar disorder: An English cohort study. *BMJ Open* 2020; accepted.
 33. Rathod S, Phiri P, Harris S, et al. Cognitive behaviour therapy for psychosis can be adapted for minority ethnic groups: A randomised controlled trial. *Schizophr Res* 2013;143, 319-326.
<http://dx.doi.org/10.1016/j.schres.2012.11.007>
 34. Binnie J, Boden Z. Non-attendance at psychological therapy appointments. *Mental Health Review Journal* 2016;21(3):231–248.

Reporting checklist for cross sectional study.

Based on the STROBE cross sectional guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the STROBE cross sectional reporting guidelines, and cite them as:

von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies.

		Reporting Item	Page Number
Title and abstract			
Title	#1a	Indicate the study's design with a commonly used term in the title or the abstract	1
Abstract	#1b	Provide in the abstract an informative and balanced summary of what was done and what was found	1
Introduction			
Background / rationale	#2	Explain the scientific background and rationale for the investigation being reported	1,3
Objectives	#3	State specific objectives, including any prespecified hypotheses	3
Methods			
Study design	#4	Present key elements of study design early in the paper	4
Setting	#5	Describe the setting, locations, and relevant dates, including periods of	4

		recruitment, exposure, follow-up, and data collection	
1			
2			
3	Eligibility criteria	#6a Give the eligibility criteria, and the sources and methods of selection of participants.	4
4			
5			
6		#7 Clearly define all outcomes, exposures, predictors, potential	1,4-6,
7		confounders, and effect modifiers. Give diagnostic criteria, if applicable	
8			
9			
10	Data sources /	#8 For each variable of interest give sources of data and details of methods	4-6
11	measurement	of assessment (measurement). Describe comparability of assessment	
12		methods if there is more than one group. Give information separately	
13		for for exposed and unexposed groups if applicable.	
14			
15			
16			
17	Bias	#9 Describe any efforts to address potential sources of bias	4
18			
19	Study size	#10 Explain how the study size was arrived at	4
20			
21	Quantitative	#11 Explain how quantitative variables were handled in the analyses. If	4-6
22	variables	applicable, describe which groupings were chosen, and why	
23			
24			
25	Statistical	#12a Describe all statistical methods, including those used to control for	6
26	methods	confounding	
27			
28			
29	Statistical	#12b Describe any methods used to examine subgroups and interactions	6
30	methods		
31			
32			
33	Statistical	#12c Explain how missing data were addressed	4
34	methods		
35			
36			
37	Statistical	#12d If applicable, describe analytical methods taking account of sampling	6
38	methods	strategy	
39			
40			
41	Statistical	#12e Describe any sensitivity analyses	6
42	methods		
43			
44	Results		
45			
46			
47	Participants	#13a Report numbers of individuals at each stage of study—eg numbers	1,6
48		potentially eligible, examined for eligibility, confirmed eligible,	
49		included in the study, completing follow-up, and analysed. Give	
50		information separately for for exposed and unexposed groups if	
51		applicable.	
52			
53			
54			
55	Participants	#13b Give reasons for non-participation at each stage	6
56			
57	Participants	#13c Consider use of a flow diagram	N/A
58			
59			
60			

1	Descriptive data	#14a	Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders. Give information separately for exposed and unexposed groups if applicable.	6,7
2				
3				
4				
5				
6	Descriptive data	#14b	Indicate number of participants with missing data for each variable of interest	4
7				
8				
9				
10	Outcome data	#15	Report numbers of outcome events or summary measures. Give information separately for exposed and unexposed groups if applicable.	8
11				
12				
13				
14	Main results	#16a	Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	8-10
15				
16				
17				
18				
19	Main results	#16b	Report category boundaries when continuous variables were categorized	8-10
20				
21	Main results	#16c	If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
22				
23				
24				
25	Other analyses	#17	Report other analyses done—e.g., analyses of subgroups and interactions, and sensitivity analyses	10
26				
27				
28				
29	Discussion			
30				
31	Key results	#18	Summarise key results with reference to study objectives	10-11
32				
33				
34	Limitations	#19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias.	12
35				
36				
37				
38				
39	Interpretation	#20	Give a cautious overall interpretation considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence.	10-12
40				
41				
42				
43				
44	Generalisability	#21	Discuss the generalisability (external validity) of the study results	12
45				
46				
47	Other			
48	Information			
49				
50				
51	Funding	#22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	2
52				
53				
54				
55				

The STROBE checklist is distributed under the terms of the Creative Commons Attribution License CC-BY.

This checklist was completed on 11. March 2021 using <https://www.goodreports.org/>, a tool made by the

[EQUATOR Network](#) in collaboration with [Penelope.ai](#)

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

BMJ Open

The association between depressive symptoms and Cognitive Behavioural Therapy receipt within a psychosis sample: A cross-sectional study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-051873.R1
Article Type:	Original research
Date Submitted by the Author:	25-Feb-2022
Complete List of Authors:	Mason, Ava; University College London, Division of Psychiatry ; King's College London, Institute of Psychiatry, Psychology and Neuroscience Irving, Jessica; Institute of Psychiatry Psychology and Neuroscience, Psychosis Studies Pritchard, Megan; King's College London, Institute of Psychiatry, Psychology and Neuroscience Sanyal, Jyoti; South London and Maudsley Mental Health NHS Trust Colling, Craig; South London and Maudsley NHS Foundation Trust, Biomedical Research Centre Chandran, David; King's College London, Institute of Psychiatry Stewart, Robert; King's College London, Institute of Psychiatry
Primary Subject Heading:	Mental health
Secondary Subject Heading:	Mental health
Keywords:	Schizophrenia & psychotic disorders < PSYCHIATRY, Depression & mood disorders < PSYCHIATRY, Adult psychiatry < PSYCHIATRY, Health informatics < BIOTECHNOLOGY & BIOINFORMATICS, PSYCHIATRY

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

The association between depressive symptoms and Cognitive Behavioural Therapy receipt within a psychosis sample: A cross-sectional study.

Ava Mason¹, Jessica Irving¹, Megan Pritchard^{1,2}, Jyoti Sanyal², Craig Colling^{1,2}, David Chandran¹, Robert Stewart^{1,2}.

¹King's College London Institute of Psychiatry, Psychology and Neuroscience, De Crespigny Park, London, SE5 8AF

²South London and Maudsley NHS Foundation Trust, London, UK

* Address correspondence to: Ava Mason, University College London, Gower St, Bloomsbury, London WC1E 6BT, UK. Email: ava.mason.20@ucl.ac.uk.

Keywords: CBTp, psychosis, cognitive behaviour therapy, depression, mental illness.

Abstract word count: 249

Main body word count: 4959

Abstract

Objectives: To examine whether depressive symptoms predict receipt of cognitive behavioural therapy (CBTp) in individuals with psychosis.

Design: Retrospective cross-sectional analysis of electronic health records (EHRs) of a clinical cohort.

Setting: A secondary NHS mental health care service serving four boroughs of south London, UK.

Participants: 20,078 patients diagnosed with an ICD-10 code between F20-29 extracted from an EHR database.

Primary and secondary outcome measures: Primary: Whether recorded depressive symptoms predicted recorded CBTp session receipt (at least one session) identified from structured EHR fields supplemented by a natural language processing algorithm. Secondary: Whether age, gender, ethnicity, symptom profiles (positive, negative, manic and disorganisation symptoms), a comorbid diagnosis of depression, anxiety or bipolar disorder, prior CBT receipt, or type of psychosis diagnosis predicted recorded CBTp receipt.

Results: Of patients with a psychotic disorder, only 8.2% were recorded as receiving CBTp. At least one depressive symptom recorded, depression symptom severity and 12 out of 15 of the individual depressive symptoms independently predicted CBTp receipt. Female gender, White ethnicity and presence of a comorbid affective disorder or primary schizoaffective diagnosis were independently positively associated with CBTp receipt within the whole sample and the top 25% of mentioned depressive symptoms.

Conclusions: Individuals with a psychotic disorder who had recorded depressive symptoms were significantly more likely to have recorded receipt of CBTp sessions, aligning with CBTp guidelines of managing depressive symptoms related to a psychotic experience. However, overall recorded receipt of CBTp is low, unequal between demographic groups, and needs to be increased.

Strengths and limitations of the study

- To our knowledge, this is the first electronic health record (EHR) study to measure how clinical symptomatology predicts recorded CBTp receipt, providing insight on a large sample into whether individuals who may be more in need of CBTp are more likely to have a session
- We replicate previous findings of inequalities in gender and ethnicity in real-world CBTp treatment receipt in a large heterogeneous sample.
- The natural language processing approach allows automated processing of EHR text at scale and can evaluate larger samples than manually conducted case note audits; this could therefore be used more routinely to monitor CBTp receipt.

- This study was limited to a single service provider; however, the results identified themes consistent with previous CBTp provision research in other services.
- Analysing EHRs in this way can identify CBTp receipt but is less suited to investigate whether CBTp is offered or not, or to quantify the quality or focus of the sessions. Furthering this, it cannot be used to examine CBTp completion rates and effectiveness.

Conflict of Interest

The authors report no conflict of interest with respect to the findings described in this manuscript. RS declares research support received in the last 36 months from Janssen, GSK and Takeda.

Financial Support

The independent research of this paper is funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at the South London and Maudsley NHS Foundation Trust and King's College London. RS is part-funded by: i) the National Institute for Health Research (NIHR) Biomedical Research Centre at the South London and Maudsley NHS Foundation Trust and King's College London; ii) an NIHR Senior Investigator Award; iii) the National Institute for Health Research (NIHR) Applied Research Collaboration South London (NIHR ARC South London) at King's College Hospital NHS Foundation Trust; iv) the DATAMIND HDR UK Mental Health Data Hub (MRC grant MR/W014386).

The Corresponding Author (AM) has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non-exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in BMJ editions and any other BMJPL products and sublicences such use and exploit all subsidiary rights, as set out in our licence.

Transparency declaration

The lead author AM affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Patient and public involvement

The Clinical Record Interactive system as a data resource was developed and is run with extensive patient involvement. However, this particular analysis did not involve patients in its design or implementation.

Ethics statement

The CRIS data platform has received research ethics approval as an anonymised data resource for secondary analysis (Oxford Research Ethics Committee C, reference 18/SC/0372)

Introduction

There are a variety of cognitive and emotional processes involved in the development of psychotic symptoms,¹ with intense distress emerging early on in the course of the disorder. Content of positive symptoms often mirrors the content of depressive thinking processes,² suggesting therapeutic need for individuals experiencing additional depressive symptoms. Specific depressive symptoms that often accompany psychotic disorders are hopelessness, social avoidance and problems in forming relationships.³ Around 50% of patients with psychosis report having experienced suicidal ideation at least once,⁴ and around 40% of individuals with schizophrenia report clinical levels of depression and low self-esteem.⁵ Importantly, individuals report these emotional difficulties and resulting social exclusion to be more debilitating than their psychotic symptoms.⁶ Consequentially, individuals' negative appraisal of their psychotic experiences may lead to loss of social goals and increased shame, predicting later hopelessness and post-psychotic depression.⁷ This comorbid depression increases the likelihood of having a lower quality of life, function, motivation, poorer social relationships, lower medication adherence and psychotic relapse (significant increase in psychotic symptoms).^{8,9,10} Therefore, treatment should focus on the psychotic symptoms and the broader distress they produce, building self-esteem, confidence and a sense of self control and purpose.¹¹ Additionally, focusing on mood symptoms such as self-esteem and pessimism can help differentiate depressive symptoms from negative psychotic symptoms, that often show significant clinical overlap.⁵

It is increasingly recognised that medication alone is inadequate for tackling psychosis symptoms.¹² In the UK, the National Institute of Clinical Excellence¹³ has recommended that cognitive behavioural therapy for psychosis (CBTp) be offered universally to individuals with psychosis. Based on the stress-vulnerability model,¹⁴ CBTp focuses on distress reduction related to hallucinations and delusions, through targeting negative beliefs and improving self-esteem.^{10, 15} Sessions often focus on goal setting and emotional issues such as rebuilding one's self, positivity and acceptance.¹¹ While studies examining characteristics of CBTp show strong evidence that CBTp improves depressive symptoms in the context of psychosis, specifically with long term reductions in suicidal behaviour,^{10, 15, 16} service provision of this intervention still falls far short of the universal access recommended.¹²

Considering the impact of targeting these symptoms in CBTp sessions, it is important to monitor receipt of CBTp within psychosis samples. While CBTp provision shows moderate yearly increases (12.8% in 2013 to 14.8% in 2014), the treatment is still only available to a small proportion of individuals,¹² short of NICE universal access recommendations.¹³ Previous studies investigating CBTp receipt have conducted time-consuming audits on limited sample sizes; these can be affected by under-reporting. On the other hand, the UK's National Mental Health Minimum Data Set report does not require CBT interventions to be recorded in a given individual's record. Natural language processing techniques (NLP)¹⁷ offer the opportunity to extract this information from free text in electronic health records (EHRs) across large numbers of patients with psychosis, and a recent study developed and applied NLP in this respect, finding higher levels of receipt than reported in previous audit, supported by the high positive predictive value and sensitivity of the technique (95% and 96% respectively).¹²

While studies have examined general CBTp receipt within patients with psychosis, no study has examined a link between depressive symptoms and CBTp receipt.¹² Therefore, we investigated whether depressive symptoms predict CBTp receipt in people with psychosis by applying these previously data extraction techniques to secondary mental health care EHRs for a large South London catchment population. Secondary predictors of receipt were type of psychosis diagnosis (schizophrenia, schizoaffective disorder, or other schizophrenia spectrum disorder), symptom profiles (negative, manic or disorganisation), general CBT receipt prior to psychosis diagnosis, comorbid depression, anxiety or bipolar diagnosis and socio-demographic factors (ethnicity, gender and age).

Methods

For this study, we extracted data on individuals with a diagnosis of a recognised schizophrenia spectrum diagnosis from the case registry of the South London and Maudsley NHS Foundation Trust (SLaM). This is a large secondary care mental healthcare provider, serving around 1.3 million residents in Croydon, Lambeth, Lewisham, and Southwark. SLaM care covers all specialist mental health care, including early intervention services, liaison and crisis teams and community and inpatient services. EHRs have been used for all SLaM services since 2006, with the Clinical Record Interactive Search system (CRIS) being established in 2008 to facilitate the retrieval of de-identified data from these records of patients previously or currently receiving mental healthcare from SLaM.¹⁸ The source EHR contains unstructured free text fields from correspondence, personal histories, mental health examinations and management plans, as well as structured fields for coding demographic information, like age and ethnicity. Implementing data from all these fields reduces selection bias of utilising only specific sources of information from the EHR. Consequently, a large programme of work has developed a range of NLP algorithms over the last decade, whose detailed descriptions and performance data are contained in an open-access catalogue.¹⁹ CRIS has approval as a data resource for secondary analysis (Oxford Research Ethics Committee C, reference 18/SC/0372), and a service user-led committee considers all proposed research before access to CRIS data is granted.

We extracted data for all individuals receiving SLaM care between January 2007 and June 2020 with a primary diagnosis of an ICD-10-defined schizophrenia spectrum disorder (F20-F29) and above the age of 18 at the time their original referral was accepted. The index date for covariate definitions was the date of the first diagnosis within this grouping. Individuals may have been active within the service before their index date, allowing us to extract data on prior CBT receipt. The sample was restricted to those with data on all variables.

Ethnicity, age at referral and gender were also extracted. Ethnicity was categorised into six groups for analysis: 'White British' (British), 'White other' (Irish or any other white background), 'Black' (Caribbean, African or any other black background), 'Asian' (Indian, Bangladeshi, Pakistani, Chinese or any other Asian background), 'Other/mixed' (white and Asian, white and black Caribbean, white and black African, any other ethnic group) and 'Not stated'.

Diagnosis was categorised into three subgroups of schizophrenia (ICD-10 codes F20.0–F20.9), schizoaffective disorder (F25.0–F25.9) and other schizophrenia spectrum disorder (F21, F22.0–F22.9, F23.0–F23.9, F24, F28 and F29). Within the data collection period, secondary diagnosis of depression (ICD-10: F32 or F33), anxiety (ICD-10: F40 or F41), or bipolar disorder (ICD-10: F31) were also extracted from structured field data.

NLP algorithms for each specific symptom were used to identify recorded symptom profiles within participants. Symptoms were categorised as depressive, positive, negative, manic or disorganisation. These symptoms had been categorised *a priori* by developers of the original independent symptom NLP algorithms. As symptoms could be labelled in more than one category during analysis, multicollinearity tests using the R function `vif()` within the [car package] were undertaken to avoid issues with overlapping predictor variables. All variables were included due to their VIF values being below five. However, positive symptoms were excluded from regression models using categorical symptom variables (having at least one mention within the EHR), as this factor variable only had one level, due to all participants having at least one positive symptom. The overall symptom list and subsequent recoding can be found in Table 1. Presence of at least one mention of any symptom in the five categories was computed as a binary variable (0/1).

Table 1
Classification of symptom predictors

Symptom	Symptom label
Aggression	Positive
Agitation	Positive
Anergia	Depressive /Negative
Anhedonia	Depressive/Negative
Apathy	Depressive/Negative
Arousal	Manic
Blunted affect	Depressive/Negative
Circumstantiality	Disorganisation
Delusions	Positive
Derailment	Disorganisation
Disturbed sleep	Depressive/Manic
Elation	Manic
Emotional Withdrawal	Negative
Flight of ideas	Disorganisation
Formal thought disorder	Disorganisation
Grandiosity	Manic
Guilt	Depressive
Hallucinations (auditory)	Positive
Helplessness	Depressive
Hopelessness	Depressive
Hostility	Positive
Insomnia	Depressive/Manic
Irritability	Manic
Paranoia	Positive
Persecutory ideation	Positive
Poor appetite	Depressive
Poor concentration	Depressive
Poor motivation	Depressive
Poverty of speech	Negative
Poverty of thought	Negative
Social withdrawal	Negative
Suicidal ideation	Depressive
Tangentiality	Disorganisation
Tearfulness	Depressive
Thought block	Disorganisation
Worthlessness	Depressive

The date of the first and last general CBT session before the index date was extracted. This was coded as a binary variable, with individuals in the 'Prior CBT' receipt group having at least one session date mention prior to their index date. This was included as a predictor to adjust for previous experience of the specific CBT intervention. Mentions were extracted using the same NLP tool as the CBTp outcome measure mentioned subsequently.

1
2
3 The primary outcome was recorded CBTp receipt, identified using a combination of structured fields and
4 NLP.¹⁷ The NLP algorithms for general CBT has high PPV and sensitivity,¹² consistent with other NLP algorithms
5 such as medication dose and diagnosis.²⁰ The date of the first CBTp session on or after the index date was
6 extracted and computed as a binary variable, so that individuals in the 'CBTp receipt' group had at least one
7 CBTp session mention after the index date.
8

9 **Statistical analysis**

10
11 To avoid overfitting, we followed the 'one in ten' rule, whereby one predictor can be measured for every 10
12 events. As the data included 1647 recorded CBTp events, our study was able to include all 12 predictors within
13 the same regression model.
14

15 All statistical analyses were conducted using R (version 1.3.9). Descriptive statistics for demographic and
16 clinical variables are reported as frequencies for categorical variables and means and standard deviations for
17 the continuous variable (age at referral). Chi square tests were also calculated for categorical variables, and t-
18 test for age to measure between-group differences in those with/without CBT receipt. Descriptive statistics
19 were also provided for yearly CBT prior to index date and recorded CBTp receipt post index date within the
20 data extraction time period (2007-2020).
21
22

23
24 Binary logistic regression was used to examine the association between depressive symptoms and recorded
25 receipt of at least one CBT session in the whole sample. For this, three regression models were analysed.
26 Model 1 was an unadjusted model with only depressive symptoms as the predictor variable. Due to significant
27 provision differences seen in previous CBTp studies,¹² model 2 (partially adjusted model), adjusted for
28 sociodemographic variables (age at referral, ethnicity, gender), primary diagnosis group and presence of a
29 comorbid diagnosis (anxiety, depression, and bipolar disorder). Model 3 (fully adjusted model) also adjusted
30 for prior CBT receipt before the index date (first psychosis diagnosis date) and symptoms mention (manic,
31 negative and disorganisation symptoms). Positive psychotic symptoms were not included in these models, as
32 individuals all had at least one mention within their case notes.
33
34

35
36 As the primary aim of the study was to investigate depressive symptoms as a predictor of recorded CBTp
37 receipt, we also split the depressive symptoms category into the 15 specific depressive symptoms applications
38 within the whole sample. Model 4 was an unadjusted model with the 15 symptoms as predictor variables.
39 Model 5 was a fully adjusted model that adjusted for all the variables in Model 3. We also conducted a
40 sensitivity analysis to investigate how results were affected by overlap of negative or depressive symptom
41 annotations, by removing negative symptoms as a predictor from the logistic regression model.
42
43

44
45 As well as investigating individual depressive symptoms as predictors of recorded CBTp receipt, we also
46 investigated depression severity predicted CBTp receipt. These logistic regression models involved converting
47 depressive, disorganised, manic, positive, and negative symptoms into continuous variables, whereby severity
48 reflected the number of different individual symptoms mentioned within each symptom construct. This
49 allowed for positive symptoms to also be included within regression models. Model 6 was an unadjusted
50 model, with depressive symptom severity as a predictor of CBTp receipt. Model 7 and model 8 were partially
51 and fully adjusted models, controlling for the same variables as model 2 and 3, except categorising symptoms
52 as the continuous rather than categorical variable.
53

54
55 Lastly, to compare differences in the general sample with those with the top 25% quantity for depressive
56 symptoms, we conducted two further regression models. This subsample analysis was conducted to examine
57 predictors of CBTp receipt where a clear clinical indication was present, supplementing the overall findings.
58 Model 9 partially adjusted for socio-demographic factors, diagnostic group and comorbid diagnosis and Model
59 10 fully adjusted for prior CBT, negative and disorganisation symptoms additionally. This group all had at least
60 one manic and psychotic symptom, so these variables were not included in the model.

Table 2

Distribution frequencies on baseline demographics and diagnoses split by recorded CBTp receipt and primary diagnosis group.

	No CBTp recorded (n = 18431)	Recorded CBTp (n=1647)	Chi square tests (χ^2)
Ethnicity			$\chi^2=100.57^{***}$
White British	30% (5516/18431)	32.8% (540/1647)	
White Other	10.4% (1908/18431)	8.5% (140/1647)	
Black	36.5% (6719/18431)	41.7% (687/1647)	
Asian	6.5% (1193/18431)	5.2% (86/1647)	
Other/Mixed	9.8% (1808/18431)	10.5% (173/1647)	
Not stated	7.0% (1287/18431)	1.3% (21/1647)	
Gender			$\chi^2= 2.75$
Female	41.4% (7636/18431)	43.5% (717/1647)	
Male	58.6% (10795/18431)	56.5% (930/1647)	
Bipolar diagnosis	4.4% (810/18431)	9.0% (149/1647)	$\chi^2=71.94^{***}$
No diagnosis	95.6% (17621/18431)	91.0% (1498/1647)	
Depression diagnosis	7.4% (1373/18431)	14.0% (230/1647)	$\chi^2=87.36^{***}$
No diagnosis	92.6% (17058/18431)	86.0% (1417/1647)	
Anxiety diagnosis	2.4% (441/18431)	7.0% (115/1647)	$\chi^2=118.28^{***}$
No diagnosis	97.6% (17990/18431)	93.0% (1532/1647)	
Prior CBT	3.1% (573/18431)	14.4% (237/1647)	$\chi^2=497^{***}$
No prior CBT	96.9% (17858/18431)	85.6% (1410/1647)	

*** $p<.001$

Results

Participants

The cohort comprised 20,078 individuals with the inclusion diagnoses, 1647 (8.2%) of whom received at least one recorded session of CBTp after their first diagnosis date. The mean age of the cohort was 42.4 years (SD=16.5). Distribution frequencies for all categorical variables can be found in Table 2. Chi-square test results represented in this table compared those with or without CBTp receipt. All mentioned variables showed significant between-group differences at $p<.001$ apart from gender (No CBTp group females=41.4%, CBTp delivery group females= 43.5%; $\chi^2=2.75$, $p=.097$). These significant variables include depression diagnosis ($\chi^2=87.36$), bipolar diagnosis ($\chi^2=71.94$), anxiety diagnosis ($\chi^2=118.28$) and prior CBT receipt ($\chi^2=497$). Additionally, the Welch two sample t-test found significant between-group differences in age ($t=15.34$, $p<.01$). Where those who had received CBTp had a lower mean age ($M=33.12$ SD= 11.5) compared to those who did not ($M=35.88$, SD=13.08). The significant results confirmed the need for further analysis through the regression models. Positive psychotic symptoms were excluded from chi square and regression analysis, as all patients had at least one positive psychotic symptom.

CBT receipt.

The descriptive results shown suggest that there is a low prevalence of both prior CBT and recorded CBTp post diagnosis across the years (Table 3, supplementary figure). The decrease in recorded receipt in 2020 can be explained by COVID-19, while 2019 receipt is comparable to previous years.

Table 3

Distribution frequencies on recorded CBT receipt (prior to diagnosis) and recorded CBTp receipt (post diagnosis) per year of data extraction.

Year	CBT prior	CBT post	All CBT
2007	130	81	211
2008	89	146	235
2009	59	111	170
2010	48	107	155
2011	37	105	142
2012	39	96	135
2013	32	128	160
2014	25	143	168
2015	24	150	174
2016	29	115	144
2017	16	127	143
2018	16	114	130
2019	15	153	168
2020	2	71	73
Total	561	1647	2208

General depressive symptom mention regression analysis

Results from the unadjusted (model 1), partially adjusted (model 2) and fully adjusted regression (model 3) are displayed in Table 4. Regression model 1 found that general mention of at least one of 15 potential depressive symptoms significantly predicted recorded CBTp receipt. Regarding model 2 and 3, individuals with at least one depressive, negative or disorganisation symptom mention, being of female gender, white ethnicity, prior CBT receipt and presence of a comorbid affective disorder independently positively associated with recorded CBTp receipt.

Table 4

Unadjusted, partially and fully adjusted logistic regression models for recorded CBTp receipt (Regression model 1, 2 and 3) with categorical symptom measures.

		Unadjusted	Partially adjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)	OR(95%CI)
Depressive symptoms				
1+ depressive symptom mention	18286(91.1)	3.78(2.94-4.96)***	3.42 (2.58-4.60)***	2.00(1.10-3.20)***
Bipolar diagnosis				
Has f31 diagnosis	959(4.80)		0.52(0.33-0.71)***	0.32(0.12-0.52)***
Depression diagnosis				
Has f32 diagnosis	603(80)		0.52(0.36-0.67)***	0.33(0.16-0.49)**
Anxiety diagnosis				
Has f40/41 diagnosis	556(2.80)		0.89(0.66-1.11)***	0.73(0.49-0.97)***
Age	N/A		-0.03(-0.04- -0.03)***	-0.03(-0.03- -0.02)***
Gender				
Male			Reference category	
Female	8353(41.60)		0.20(0.09-0.31)***	0.20 (0.10-0.32)***
Ethnic group				
White British	6056(30.10)		Reference category	
White Other	2048(10.20)		-0.40(-0.60 - -0.21)***	-0.37(-0.57- -0.17)***
Black	7406(36.90)		-0.16(-0.28- -0.04)**	-0.24 (-0.36- -0.11)***
Asian	1279(6.40)		-0.49(-0.74- -0.26)***	-0.50 (-0.75- -0.27)***
Other/Mixed	1981(9.90)		-0.21(-0.40- -0.02)**	-0.18(-0.37- -0.01)*
Not Stated	1308(6.50)		-1.75(2.23- -1.22)***	-1.52(-2.00- -1.10)***
Primary diagnosis				
Schizophrenia	9845(49.00)		Reference category	
Schizoaffective disorder	2142(10.70)		0.04(-0.13-0.21)	0.01(-0.17-0.19)
Other schizophrenia spectrum	8091(40.30)		-0.10(-0.22-0.01)*	-0.02(-0.14 -0.10)
Negative symptoms				
1+ Negative symptom mention	13169(65.60)			0.75(0.59-0.92)***
Manic symptoms				
1+ Manic symptom mention	17945(89.40)			1.24(0.70-1.87)***
Disorganisation symptoms				
1+ Disorganisation symptom mention	11513(57.30)			0.31(0.18-0.44)***
CBT prior				
1+ prior CBT session				1.29(1.12-1.46)***

*p<.05, **p<.01, ***p<.001

Unadjusted (model 1): depressive symptom as a predictor with no adjusted covariates

Partially adjusted (model 2): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis

Fully adjusted (model 3): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, depressive symptoms, prior CBT, negative symptoms, disorganisation symptoms, manic symptoms.

Regression analysis with individual depressive symptoms

Results from the unadjusted (model 4) and fully adjusted (model 5) regression analyses for each of the 15 individual depressive symptoms are displayed in Table 5 (N=20,078). Each symptom refers to presence of at least one mention in the patients notes compared to no mention. While all variables were significant in the unadjusted model at $p < .001$, the fully adjusted model reduced the significance of suicide ideation ($p < .01$) and disturbed sleep ($p < .01$), with anhedonia, anergia, apathy, and blunted affect becoming non-significant ($p > 0.05$).

Table 5

Unadjusted and fully adjusted logistic regression models for recorded CBTp receipt with individual depressive symptoms as covariates (Regression model 4 and 5) for the overall sample.

		Unadjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)
Hopelessness	7345(36.60)	4.81(4.3-5.40)***	1.45(1.26-1.66)
Helplessness	3124(15.60)	4.03(3.62-4.50)***	1.55(1.37-1.76)***
Suicide ideation	9451(47.10)	4.11(3.66-4.63)***	1.25(1.09-1.44)**
Poor appetite	8044(40.10)	3.31(2.97-3.68)***	1.28(1.13-1.45)***
Poor motivation	8630(43.00)	4.34(3.87-4.86)***	1.43(1.24-1.64)***
Insomnia	6870(34.20)	3.74(3.35-4.15)***	1.4(1.24-1.58)***
Disturbed sleep	16667(83.00)	15.3(10.16-22.8)***	2.76(1.5-5.08)**
Poor concentration	12289(61.20)	8.16(6.81-9.77)***	2.33(1.9-2.85)***
Anhedonia	4047(20.20)	2.9(2.61-3.22)***	0.97(0.85-1.10)
Anergia	873(43.50)	2.63(2.20-3.15)***	0.98(0.80-1.20)
Apathy	4149(20.70)	2.21(1.98-2.46)***	0.93(0.82-1.05)
Guilt	8178(40.70)	4.6(4.1-5.15)***	1.49(1.30-1.70)***
Tearfulness	10951(54.50)	3.87(3.41-4.39)***	1.22(1.05-1.42)**
Blunted affect	6889(34.30)	2.66(2.41-2.95)***	0.91(0.80-1.03)
Worthlessness	2921(14.50)	3.94(3.53-4.40)***	1.37(1.21-1.56)***

* $p < .05$, ** $p < .01$, *** $p < .001$

Sensitivity analysis

The non-significant results of certain depressive symptoms (anhedonia and anergia) may have been due to their inclusion within the negative symptom category, causing over-adjustment of the model. To test this, sensitivity analysis was conducted, where the fully adjusted regression (model 3) did not include negative symptoms as a covariate. While all significant variables remained significant, non-significant results for anhedonia and apathy were also still found. Therefore, we report the fully adjusted model with negative symptoms as a variable for both grouped and individual depressive symptom associations.

General depressive symptom severity regression analysis

Results from the unadjusted (model 6), partially adjusted (model 7) and fully adjusted regression (model 8) are displayed in Table 6. Regression model 6 found that depression symptom severity significantly predicted CBTp receipt. Regarding model 7 and 8, depression symptom severity, positive symptom severity, anxiety diagnosis, and being of older age or being of white ethnicity independently positive predicted CBTp receipt. Within model 7., being female also positively increased likelihood of CBTp receipt. Within model 8, negative symptom severity and prior CBT significantly predicted CBTp receipt additionally.

Table 6

Unadjusted, partially and fully adjusted logistic regression models for recorded CBTp receipt (Regression model 1, 2 and 3) with continuous symptom measures.

		Unadjusted	Partially adjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)	OR(95%CI)
Depressive symptoms				
Severity	18286(91.1)	0.29(1.31-1.35)***	0.27(1.20-1.44)***	0.23(0.13-0.33)***
Positive symptoms				
Severity	20078(100)		-0.18(0.75-0.92)***	-0.21(-0.31- -0.09)***
Bipolar diagnosis				
Has f31 diagnosis	959(4.80)		0.21(0.94-1.63)	0.15(-0.13-0.43)
Depression diagnosis				
Has f32 diagnosis	603(80)		-0.09(0.72-1.15)	-0.09(-0.33-0.15)
Anxiety diagnosis				
Has f40/41 diagnosis	556(2.80)		0.49(1.15-2.29)***	0.46(0.11-0.80)***
Age	N/A		-0.02(0.97-0.99)***	-0.02(-0.03- -0.01)***
Gender				
Male			Reference category	
Female	8353(41.60)		0.17(0.97-1.44)*	0.17(-0.03-0.36)
Ethnic group				
White British	6056(30.10)		Reference category	
White Other	2048(10.20)		-0.41(0.45-0.96)**	-0.44(-0.93- -0.07)***
Black	7406(36.90)		-0.25(0.63-0.97)**	-0.29(-0.51- -0.07)**
Asian	1279(6.40)		-0.66(0.32-0.80)***	-0.67(-1.13 - -0.23)***
Other/Mixed	1981(9.90)		-0.14(0.62-1.22)	-0.16(-0.50-0.18)
Not Stated	1308(6.50)		-0.92(0.02-2.42)	-0.79(-3.74-1.01)
Primary diagnosis				
Schizophrenia	9845(49.00)		Reference category	
Schizoaffective disorder	2142(10.70)		-0.08(0.69-1.21)	-0.11(-0.40-0.18)
Other schizophrenia spectrum	8091(40.30)		0.02(0.82-1.26)	-0.06(-0.15-0.28)
Negative symptoms				
Severity	13169(65.60)			0.06(-0.01-0.123)*
Manic symptoms				
Severity	17945(89.40)			-0.01(-0.13-0.12)
Disorganisation symptoms				
Severity	11513(57.30)			0.10(-0.05-0.25)
CBT prior				
1+ prior CBT session	1647(8.20)			0.62(0.34-0.89)***

*p<.05, **p<.01, ***p<.001

Unadjusted (model 1): depressive symptom severity as a predictor with no adjusted covariates

Partially adjusted (model 2): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, positive symptom severity

Fully adjusted (model 3): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, positive symptom severity, prior CBT, negative symptom severity, disorganisation

symptom severity and manic symptom severity.

Depressive symptom regression analysis within the top 25% number of depressive symptoms.

This sample comprised individuals with the top 25% number of depressive symptoms (5018 patients), defined to reflect a sub-group who might reasonably expect to receive CBT on this basis. The sample characteristics and regression analysis can be seen in Table 7. Results from the partially adjusted (model 9) and fully adjusted regression (model 10) are displayed in Table 7. Table 4 found that general mention of at least one of 15 potential depressive symptoms significantly predicted CBTp receipt. Regarding model 9, we found that individuals with at least one depressive, negative or disorganisation symptom mention, being of female gender, white ethnicity, prior CBT receipt and presence of comorbid bipolar disorder were positively associated with recorded CBTp receipt.

Table 7

Partially and fully adjusted logistic regression models for CBTp receipt with individual depressive symptoms as covariates within top 25% quantity of depressive symptoms (Regression model 9 and 10).

	N(%)	Partially adjusted OR(95%CI)	Fully adjusted OR(95%CI)
Bipolar diagnosis			
Has f31 diagnosis	541(10.78)	0.19(-0.03-0.41)*	0.15(-0.08-0.38)
Depression diagnosis			
Has f32 diagnosis	885(17.63)	0.11(-0.08-0.30)	0.08(-0.90-1.30)
Anxiety diagnosis			
Has f40/41 diagnosis	270(5.38)	0.53 (0.25-0.80)***	0.47(0.19-0.75)***
Age			
M= 36.24(18-93)		-0.02(-0.02- -0.01)***	-0.02(-0.03- -0.01)***
Gender			
Male	2059(41.01)	Reference category	
Female	2960(58.99)	0.20 (0.05-0.34)***	0.17(0.02-0.32)**
Ethnic group			
White British	1486(29.59)	Reference category	
White Other	433(8.63)	-0.21(-0.48-0.05)	-0.22(0.50-0.05)
Black	2262(45.08)	-0.32(-0.49- -0.16)***	-0.31(0.47- -0.14)***
Asian	328(6.53)	-0.53(0.86- -0.21)***	-0.52(0.85- -0.20)***
Other/Mixed	467(9.31)	-0.08(-0.34-0.17)	-0.08(0.34-0.17)
Not Stated	43(0.86)	-1.42 (-2.85 -0.40)**	-1.34(-2.8 - -0.31)**
Primary diagnosis			
Schizophrenia	2219(44.22)	Reference category	
Schizoaffective disorder	740(14.72)	0.04(-0.18-0.26)	0.003(-0.22-0.22)
Other schizophrenia spectrum	2060(41.03)	0.02(-0.14 -0.17)	0.05(-0.11-0.21)
Negative symptoms			
1+ Negative symptom mention	4956(98.7)		-0.88(1.41- -0.33)***
Disorganisation symptoms			
1+ Disorganisation symptom mention	4199(83.66)		1.18(-0.02-0.38)*
CBT prior			
1+ prior CBT session	436(8.7)		-0.90(0.68-1.11)***

*p<.05, **p<.01, ***p<.001,

Partially adjusted (model 9): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis.

Fully adjusted (model 10): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, prior CBT, negative symptoms, disorganisation symptoms

Discussion

1
2
3
4 We believe that this is the first study to examine the relationship between clinical symptomatology and CBTp receipt
5 within a sample of people with psychosis in a naturalistic community setting. In general, only 8.2% of individuals were
6 ascertained as having received CBTp within the 13-year timeframe of the study, showing the low prevalence of
7 recorded receipt despite current clinical guidelines. This finding shows a lower overall level of recorded CBTp provision
8 compared to previous studies in 2013 (12.8%) and 2014 (14.8%)¹². This requires further examination, considering the
9 importance of CBTp mentioned within NICE universal access recommendations.¹³ Additionally, the significant decrease
10 of CBTp receipt in 2020 can be explained by the COVID pandemic and therefore, it is important to consider how we can
11 improve receipt despite this.
12

13 In the analysed sample, 91% of patients had at least one recorded depressive symptom mention, and these individuals
14 were 2 times more likely to have at least one recorded CBTp session in the fully adjusted model (table 4), suggesting
15 that the minority who don't present with any depressive symptoms are very unlikely to receive CBTp. This could
16 possibly be due to clinicians tending to cite a depressive symptom when referring an individual with psychosis to
17 psychotherapy. Additionally, the severity of depressive symptoms, as well as having at least one recorded mention
18 significantly increased likelihood of having at least one recorded CBTp session. In the sample of those with the highest
19 number of depressive symptoms (top 25%), relationships between recorded CBTp receipt and comorbid anxiety
20 diagnosis, age, gender, ethnicity, prior CBT, negative and disorganised psychotic symptoms remained. This suggests the
21 importance of these predictors in a reasonable sample of patients with higher clinical need for CBTp receipt.
22

23 Considering individual depressive symptom, the symptom which was the strongest predictor of this
24 intervention in fully adjusted models was disturbed sleep. There is a known high prevalence of sleeping
25 problems in this population,^{21,22} described by some researchers as an 'intrinsic feature of schizophrenia,²³
26 known to reduce quality of life, decreasing coping and exacerbate positive symptoms.²⁴ The significant
27 association between insomnia and psychotic-like symptoms, such as paranoia, has also been seen in non-
28 clinical populations.²⁵ Further to this, the recommended first line of treatment for sleep problems in this
29 clinical population is CBT.²⁶ Poor concentration was the next strongest depressive symptom predictor in the
30 fully-adjusted model, possibly reflecting previous findings of its association with psychosis vulnerability.²⁷ The
31 significance of helplessness, guilt and hopelessness is relevant to previous CBTp research that found significant
32 post-treatment reduction in hopelessness, self-depreciation and guilt using the Calgary Depression Rating
33 Scale for Schizophrenia.²⁸ Other significant depressive symptoms associated with low self-esteem and negative
34 self-evaluation and emotions have been found to significantly affect the development and severity of positive
35 symptoms.²⁹ This may be because positive symptoms develop as a psychological defence against low self-
36 esteem³⁰ and depression-induced guilt.³¹ Therefore, it could be suggested that the significance of each of the
37 depressive symptoms is often linked to psychotic symptoms and CBTp effectiveness. However, while there is
38 evidence of the clinical impact of depressive symptoms in schizophrenia, the associations with choice of
39 therapy must be viewed as exploratory and in need of independent replication. While a possibility may be that
40 clinicians are assuming that certain depressive symptoms are likely to be more responsive to CBTp than others,
41 there may be other unknown reasons for therapy choice that requires further investigation. General results
42 suggest that receipt of this intervention requires an increase for all of this clinical population before individuals
43 with specific symptoms can be targeted.
44
45

46 Regarding negative symptoms, the non-significant associations between specific negative symptoms (that overlapped
47 with depressive symptoms) and CBTp receipt requires further evaluation and confirmation. This was not conducted in
48 the current study due to the primary aim focusing on depressive symptoms. However, from our results on specific
49 depressive symptoms in table 5, symptoms that overlapped with negative symptoms (anhedonia, anergia, apathy and
50 blunted affect) were not associated with CBTp receipt. Additionally, negative symptoms were associated with a
51 significantly decreased likelihood of recorded CBTp receipt within the group with highest numbers of depressive
52 symptoms mentioned. Overall, this raises concerns that individuals with these specific negative/depressive symptoms
53 are no more likely and perhaps less likely to receive CBTp than someone without these symptoms. Possibly, this is due
54 to clinicians not referring these individuals because they don't believe intervention will be effective. This is in line with a
55 CBTp review of randomised control trials, finding non-significant reductions of negative symptoms,³² perhaps due to
56 the narrowing of treatments to specifically target positive symptoms.³³ However, further work should be undertaken to
57 verify that individuals are not being denied a potentially beneficial intervention because of their symptom profile.
58
59
60

1
2
3 Prior CBT receipt, comorbid disorder presence and specific symptoms (manic, disorganised, and negative) also emerged
4 as independent predictors of CBTp receipt for the general sample and within those with the top 25% depressive
5 symptom numbers. Within table 4, individuals who had any recorded CBT receipt prior to the index date were 1.29
6 times more likely to have recorded CBTp receipt later on. Also, patients with an additional comorbid disorder were
7 0.32-0.73 times more likely to have received CBTp compared to those with just a psychosis diagnosis. However, in the
8 top 25% of individuals within table 7, individuals with prior CBT were 0.40 times less likely to receive CBTp. This finding
9 requires further research to understand the effects of prior CBT and negative symptoms on CBTp receipt within
10 different psychosis subsamples. Additionally, those with anxiety were 0.47 times more likely to receive CBTp and those
11 with disorganised symptoms were 1.18 times more likely respectively. After general CBTp receipt has increased, there
12 could be a method to focus more on patients with different types of psychotic symptoms and comorbid affective
13 diagnosis. Furthering this, future research could investigate whether those who have had prior general CBT would
14 benefit from CBTp, or whether those who have not had any experience developing cognitive behaviour skills in therapy
15 should be targeted.³³

16
17 Crucially, there were also significant differences in recorded CBTp receipt between different ethnic and gender groups.
18 Male patients were 0.20 times as likely in the general sample and top 25% of depressive symptoms to have recorded
19 CBTp receipt. Black, Asian, Other and Mixed ethnic groups were between 0.21 to 0.49 times as likely to have a
20 documented CBTp session compared to individuals of white ethnicity within both the general and top 25% depressive
21 symptoms samples. Inequitable access to CBTp has been identified in previous CBTp research within a psychosis sample
22 drawn from the same data resource in 2017, finding female patients to be more likely to have received CBTp and
23 individuals of White ethnicity to have a significantly higher likelihood of CBT receipt than Black or other ethnicity
24 groups.¹² This also supports results from a recent CBTp study focusing specifically on ethnic group differences in CBTp
25 provision within SLAM, who found that in comparison to White British individuals, those from Black ethnic groups with
26 psychosis or bipolar disorder were significantly less likely to have a documented CBTp session. This is especially
27 important when considering the high prevalence of psychosis within UK Black and minority ethnic group populations.³⁴
28 Inequality in CBTp receipt may be due to ethnic variations in CBTp engagement. Some of these barriers within certain
29 communities may be increased stigma, fear of clinicians by service-users or service users by clinicians, institutional
30 racism within mental health services, or non-culturally appropriate therapy.³⁵ As differences in documented CBTp
31 receipt between ethnic groups have now been documented by three different papers in this service, it is imperative
32 that further work is conducted to increase provision of CBTp within groups less likely to receive treatment. This may
33 include targeted outreach programs and culturally adapting interventions³⁵ within these minority groups.

34
35 The present study has a number of strengths and limitations. Generally, focusing on patients with more
36 diverse functioning, comorbidity and symptom severity levels helps research identify a larger number of
37 predictors of clinical outcomes. This can be seen through our results, where negative, manic and
38 disorganisation symptoms significantly predicted recorded CBTp receipt, as well as recent research,³³ that was
39 the first to identify depression as a significant predictor of positive symptom improvement post-CBTp. This
40 highlights the importance of focusing on a clinically heterogeneous sample to realistically determine significant
41 predictors of CBTp receipt. Secondly, using an NLP approach automates the measurement of what would
42 otherwise require manually conducted audits on records and case notes, increasing the number of cases that
43 can be investigated and providing a method that could be used more routinely to monitor CBT receipt. The
44 large sample size enabled us to identify potential clinical differences in the real-life administration of CBTp
45 within a psychosis cohort, and we were able to adjust for multiple clinical variables and comorbidity diagnoses
46 to provide a more realistic understanding of the depressive symptom-CBTp receipt relationship. This time
47 frame was broad to allow the inclusion of as many active patients receiving CBTp as possible, additionally
48 circumventing monthly/seasonal variation of CBTp receipt.

49
50 One limitation of the study was the omission of strict time periods for the mention of clinical symptoms prior
51 to CBTp administration. Unfortunately, using this approach would have involved implementing time periods on
52 all of the other clinical symptoms and variables, which would have been difficult considering the number of
53 variables that would need to be controlled. In addition, the NLP symptom algorithms do not currently
54 distinguish between past or present symptoms. Therefore, symptom mentions documented after the CBTp
55 receipt date could refer to mentions of symptoms occurring prior to CBTp receipt, reducing the effectiveness
56 of using time periods. A follow-up time period after the index date was also not established, meaning that
57 participants included in the cohort at a later date may have been less likely to have had a CBTp session, due to
58 their limited time period within the service. Additionally, we did not have data regarding which type of service
59 was providing CBTp for each patient (for example, early intervention services compared to other community
60

1
2
3 services). Future studies should examine whether CBTp receipt differs depending on the service, especially
4 considering how effective CBTp provision may be in those at ultra-high risk.
5

6 While our use of additional querying of text fields allowed us to identify a significantly larger number of CBTp
7 episodes than using structured data alone, we were not able to quantify the gap between CBTp referral and
8 CBTp receipt. This is because the CBTp NLP algorithm detects CBTp receipt rather than CBTp being offered,
9 due to the wide range of subtle wording used for the latter more complex entity. The results combine effects
10 on the likelihood of CBTp being offered, with those on session receipt following an offer. While this may have
11 affected our results, previous service audits have suggested that the severity and occurrence of depressive
12 symptoms significantly decreases CBT receipt.³⁶ Therefore, if only receipt was directly measured, we would
13 expect to see similar results. It is also important to consider that we are only ascertaining recorded CBTp
14 receipt, which may result in failing to pick up all CBTp receipt instances. Considering that previous research
15 describing the app development suggests high precision and recall performance of CBTp instances (PPV= 96%,
16 sensitivity= 96%),¹² it could be suggested that low prevalence within the results is due to lack of recording
17 within the clinical health records, rather than lack of app identification. Therefore, stricter regulations are
18 required for CBTp to be reported within clinical health records. Additionally, completion rates and
19 effectiveness of the CBTp was not measured, meaning we were unable to quantify the quality or focus of the
20 sessions. Lastly, analysis was limited to patients above 18 years old, reducing the generalisability of results to
21 those who develop a schizophrenia-spectrum disorder after this age. However, the outcome of interest was
22 CBTp receipt within a relatively homogenous service structure of working age services, rather than young
23 people treated within Child and Adolescent services. Future studies should examine whether CBTp receipt
24 differs in these services.
25

26 **Future directions**

27 Initiatives such as the Improving Access to Psychological Therapies programme for serious mental illness, early
28 intervention access and projects to decrease waiting times for referral have been developed to target this
29 clinical population. However, access still falls short of recommendations and is inequitable for specific
30 psychotic diagnoses, age, and ethnicity.¹² Therefore, given the effect of CBTp on depressive symptoms³⁷,
31 perhaps its more pragmatic to focus on patients with additional depressive symptoms. Monitoring CBTp
32 receipt over time could decipher whether these initiatives are effective at increasing general access for those
33 with psychosis, and specific access for different sociodemographic groups and those with additional depressive
34 symptoms (who may benefit the most).
35
36

37
38 The significant secondary clinical and sociodemographic variables require further analysis in order to fully
39 understand the services' provision. This could involve attention given to the independent symptoms within the
40 negative, manic and disorganisation categories in a similar manner to the specific depressive symptom
41 regression models analysed. Further research could also explore why the presence of co-morbid anxiety and
42 bipolar disorder in this sample predicted CBTp receipt. Additionally, the results suggest a need to reflect on the
43 steps taken since the previous service study,³⁵ regarding inequality in CBTp receipt among gender and ethnic
44 groups, due to the consistent significant results seen. Regarding the use of EHR data, future work could involve
45 developing a separate NLP algorithm to ascertain the offering of CBTp or provide another structured field for
46 clinicians to complete for this. However, clinicians prioritise text field data for communication about CBTp
47 sessions for themselves and their colleagues rather than to collect structure data for the sake of research.
48 Therefore, as previously suggested,¹² it is important to accept the mixed structured-text field approach that
49 will remain in healthcare record data and perhaps our time is best spent in improving NLP algorithms to detect
50 the subtleties of intervention and clinical outcome data. However, the implications of our results and their
51 consistency three years after the first CBTp service paper suggest the need to use this or future algorithms for
52 service monitoring independent of these improvements.
53
54
55
56
57
58
59
60

References

1. Chadwick P, Birchwood M. The omnipotence of voices. A cognitive approach to auditory hallucinations. *Br J Psychiatry* 1994;164:190–201.
2. Fowler D. Cognitive Behavior Therapy for Psychosis: From Understanding to Treatment. *Am J Psychiatr Rehabil* 2000;4(2):199–215.
3. Hazell CM, Hayward M, Cavanagh K et al. Clara Strauss A systematic review and meta-analysis of low intensity CBT for psychosis. *Clinical Psychology Review* 2016;45:183–192.
4. Hawton, K, Sutton, L, Haw, C, Sinclair, J, Deeks, JJ. Schizophrenia and suicide: systematic review of risk factors. *Br. J. Psychiatry* 2005;187:9–20.
5. Sands JR, Harrow M. Depression during the longitudinal course of schizophrenia. *Schizophr Bull* 1999;25:157–171.
6. Greenwood KE, Sweeney A, Williams S, et al. CHOICE of Outcome In Cbt for psychosEs (CHOICE): The Development of a New Service User–Led Outcome Measure of CBT for Psychosis. *Schizophrenia Bulletin* 2010;36(1):126–135.
7. Birchwood M, Iqbal Z, Chadwick P, et al. Cognitive approach to depression and suicidal thinking in psychosis. I: Ontogeny of post-psychotic depression. *Br J Psychiatry* 2000;177: 516–528.
8. Conley RR. The burden of depressive symptoms in people with schizophrenia. *Psychiatr Clin North Am* 2009;32:853–861.
9. Rocca P, Bellino S, Calvarese P, et al. Depressive and negative symptoms in schizophrenia: different effects on clinical features. *Comprehensive Psychiatry* 2005; 46: 304–310.
10. Richardson T, Dasyam B, Courtney H, et al. Predictors of disengagement from cognitive behavioural therapy for psychosis in a National Health Service setting: A retrospective evaluation. *British journal of clinical psychology* 2019; 58(4): 440–451.
11. Pitt L, Kilbride M, Nothard S, et al. Researching recovery from psychosis: a user-led project. *Psychiatr Bull R Coll Psychiatr* 2007;31:55–60.
12. Colling C, Evans L, Broadbent M, et al. Identification of the delivery of cognitive behavioural therapy for psychosis (CBTp) using a cross-sectional sample from electronic health records and open-text information in a large UK-based mental health case register. *BMJ Open* 2017;7:e015297.
13. National Institute for Health and Clinical Excellence. Schizophrenia: Core interventions in the treatment and management of schizophrenia in primary and secondary care. London, UK: NICE Clinical Guideline 2009.
14. Rathod S, Phiri P, Kingdon D. Cognitive behavioral therapy for schizophrenia. *Psychiatr Clin North Am* 2010;33:527–536.
15. Gumley A, Schwannauer M. Staying Well after Psychosis: A Cognitive Interpersonal Approach to Recovery and Relapse Prevention. New York, NY: John Wiley & Sons Ltd 2006.
16. Lincoln TM, Rief W, Wetermann S, et al. Who stays, who benefits? Predicting dropout and change in cognitive behaviour therapy for psychosis. *Psychiatry Research* 2014;216(2):198–205.
17. Spyns P. Natural language processing in medicine: an overview. *Methods Inf Med* 1997;35(4–5):285–301.
18. Stewart R, Soremekun M, Perera G, et al. The South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLAM BRC) case register: development and descriptive data. *BMC Psychiatry* 2009;9:51. (<https://www.maudsleybrc.nhr.ac.uk/media/325736/applications-library-v13.pdf>).
19. Hayes RD, Downs J, Chang CK, et al. The effect of clozapine on premature mortality: an assessment of clinical monitoring and other potential confounders. *Schizophr Bull* 2015;41:644–55.
20. Chadwick PDJ, Birchwood MJ, Trower, P. Cognitive Therapy for Delusions, Voices and Paranoia: Towards A Person Model. Chichester: Wiley 1996.
21. Chiu V, Ree M, Janca A, et al. Sleep profiles and CBT-I response in schizophrenia and related psychoses. *Psychiatry Research* 2018;268:279–287.

23. Chouinard S, Poulin J, Stip E, et al. Sleep in untreated patients with schizophrenia: a meta-analysis. *Schizophr Bull* 2004;30:957-967.
24. Waite F, Evans N, Myers E, et al. The patient experience of sleep problems and their treatment in the context of current delusions and hallucinations. *Psychology and Psychotherapy: Theory, Research and Practice* 2016;89:181-193.
25. Freeman D, Sheaves B, Goodwin G, et al. The effects of improving sleep on mental health (OASIS): a randomised controlled trial with mediation analysis. *The Lancet Psychiatry* 2017;4(10):749-758.
26. Mitchell MD, Gehrman P, Perlis M, et al. Comparative effectiveness of cognitive behavioral therapy for insomnia: A systematic review. *BMC Family Practice* 2012;13(1):40.
27. Bogren M, Mattisson C, Tambs K, et al. Predictors of psychosis: a 50-year follow-up of the Lundby population. *Eur Arch Psychiatry Clin Neurosci* 2009;260:113-125.
28. Muller MJ, Marx-Dannigkeit P, Schlösser R, et al. The Calgary Depression Rating Scale for Schizophrenia: Development and interrater reliability of a German version (CDSS-G). *J Psychiatr Res* 1999;33:433-443.
29. Smith B, Fowler DG, Freeman D, et al. Emotion and psychosis: links between depression, self-esteem, negative schematic beliefs and delusions and hallucinations. *Schizophr Res* 2006;86(1-3):181-188.
30. Bentall RP, Kinderman P, Kaney S. The self, attribution processes and abnormal beliefs: Towards a model of persecutory delusions. *Behav Res Ther* 1994;32:331-341.
31. Lake CR. Hypothesis: Grandiosity and Guilt Cause Paranoia; Paranoid Schizophrenia is a Psychotic Mood Disorder; a Review. *Schizophr Bull* 2008;34(6):1151-1162.
32. Lincoln TM, Ziegler M, Mehl S, et al. Moving from efficacy to effectiveness in cognitive behavioral therapy for psychosis: a randomized clinical practice trial. *J Consult Clin Psychol* 2012;80(4):674-686.
33. Penn DL, Meyer PS, Evans E, et al. A randomized controlled trial of group cognitive-behavioral therapy vs. enhanced supportive therapy for auditory hallucinations. *Schizophr Res* 2009;109:52-59.
34. Morris R, Sellwood W, Edge D, et al. Ethnicity and impact on the receipt of Cognitive Behavioural Therapy in people with psychosis or bipolar disorder: An English cohort study. *BMJ Open* 2020; accepted.
35. Rathod S, Phiri P, Harris S, et al. Cognitive behaviour therapy for psychosis can be adapted for minority ethnic groups: A randomised controlled trial. *Schizophr Res* 2013;143, 319-326.
<http://dx.doi.org/10.1016/j.schres.2012.11.007>
36. Binnie J, Boden Z. Non-attendance at psychological therapy appointments. *Mental Health Review Journal* 2016;21(3):231-248.
37. Taalman H, Goldberg DM, Ayub M, et al. Effect of cognitive behaviour therapy for psychosis (Cbtp) on depressive symptoms: a review of literature. *J Schizophr Res* 2015;2(3):1019.

Acknowledgments

This paper would not have been completed without the assistance from Dr Rob Stewart. I would also like to thank Jessica Irving for her assistance in the statistical analysis and general manuscript.

Author statement

Ava Mason planned the protocol, analysed the data and wrote the manuscript. Dr Rob Stewart provided access to the data, looked over edits and revisions on the manuscript. The data from this paper was accessed through the assistance of Megan Pritchard and Jyoti Sanyal. The applications used to obtain data specific to CBTp were developed by Craig Colling and David Chandran. Jessica Irving assisted in statistical analysis.


Data Sharing Statement

Due to the terms of Ethics and Information Governance approvals and clinical source of the data, CRIS datasets must remain within the South London and Maudsley NHS Foundation Trust (SLaM) firewall. All data used from this study can be made accessible on request from cris.administrator@slam.nhs.uk, subject to the setting up of an appropriate research passport or SLaM honorary contract.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary figure
Graph demonstrating the frequency of general CBT receipt (CBT receipt prior to diagnosis and recorded CBTp receipt post diagnosis) per year of extraction period.

For peer review only



20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary figure

Graph demonstrating the frequency of general CBT receipt (CBT receipt prior to diagnosis and recorded CBTp receipt post diagnosis) per year of extraction period.

Reporting checklist for cross sectional study.

Based on the STROBE cross sectional guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the STROBE cross sectional reporting guidelines, and cite them as:

von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies.

		Reporting Item	Page Number
Title and abstract			
Title	#1a	Indicate the study's design with a commonly used term in the title or the abstract	1
Abstract	#1b	Provide in the abstract an informative and balanced summary of what was done and what was found	1
Introduction			
Background / rationale	#2	Explain the scientific background and rationale for the investigation being reported	1,3
Objectives	#3	State specific objectives, including any prespecified hypotheses	3
Methods			
Study design	#4	Present key elements of study design early in the paper	4

1	Setting	#5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	4
2				
3				
4	Eligibility criteria	#6a	Give the eligibility criteria, and the sources and methods of selection of participants.	4
5				
6				
7				
8		#7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	1,4-6,
9				
10				
11				
12				
13				
14	Data sources /	#8	For each variable of interest give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group. Give information separately for for exposed and unexposed groups if applicable.	4-6
15	measurement			
16				
17				
18				
19				
20				
21	Bias	#9	Describe any efforts to address potential sources of bias	4
22				
23	Study size	#10	Explain how the study size was arrived at	4
24				
25	Quantitative	#11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen, and why	4-6
26	variables			
27				
28				
29	Statistical	#12a	Describe all statistical methods, including those used to control for confounding	6
30	methods			
31				
32				
33	Statistical	#12b	Describe any methods used to examine subgroups and interactions	6
34	methods			
35				
36				
37	Statistical	#12c	Explain how missing data were addressed	4
38	methods			
39				
40				
41	Statistical	#12d	If applicable, describe analytical methods taking account of sampling strategy	6
42	methods			
43				
44	Statistical	#12e	Describe any sensitivity analyses	6
45	methods			
46				
47				
48	Results			
49				
50				
51	Participants	#13a	Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed. Give information separately for for exposed and unexposed groups if applicable.	1,6
52				
53				
54				
55				
56				
57				
58				
59				
60				

1	Participants	#13b	Give reasons for non-participation at each stage	6
2				
3	Participants	#13c	Consider use of a flow diagram	N/A
4				
5	Descriptive data	#14a	Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders. Give information separately for exposed and unexposed groups if applicable.	4-7
6				
7				
8				
9				
10				
11				
12	Descriptive data	#14b	Indicate number of participants with missing data for each variable of interest	4
13				
14				
15				
16	Outcome data	#15	Report numbers of outcome events or summary measures. Give information separately for exposed and unexposed groups if applicable.	8
17				
18				
19				
20				
21	Main results	#16a	Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	7-12
22				
23				
24				
25				
26				
27				
28	Main results	#16b	Report category boundaries when continuous variables were categorized	7-12
29				
30				
31				
32	Main results	#16c	If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
33				
34				
35				
36	Other analyses	#17	Report other analyses done—e.g., analyses of subgroups and interactions, and sensitivity analyses	12
37				
38				
39				
40	Discussion			
41				
42	Key results	#18	Summarise key results with reference to study objectives	13-15
43				
44	Limitations	#19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias.	14-15
45				
46				
47				
48				
49				
50	Interpretation	#20	Give a cautious overall interpretation considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence.	13-15
51				
52				
53				
54				
55	Generalisability	#21	Discuss the generalisability (external validity) of the study results	13-15
56				
57				
58				
59				
60				

1 **Other**
2 **Information**

3
4
5 Funding [#22](#) Give the source of funding and the role of the funders for the present 2
6 study and, if applicable, for the original study on which the present
7 article is based
8
9

10 The STROBE checklist is distributed under the terms of the Creative Commons Attribution License CC-BY.

11 This checklist was completed on 11. March 2021 using <https://www.goodreports.org/>, a tool made by the

12 [EQUATOR Network](#) in collaboration with [Penelope.ai](#)
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

BMJ Open

The association between depressive symptoms and Cognitive Behavioural Therapy receipt within a psychosis sample: A cross-sectional study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-051873.R2
Article Type:	Original research
Date Submitted by the Author:	18-Apr-2022
Complete List of Authors:	Mason, Ava; University College London, Division of Psychiatry ; King's College London, Institute of Psychiatry, Psychology and Neuroscience Irving, Jessica; Institute of Psychiatry Psychology and Neuroscience, Psychosis Studies Pritchard, Megan; King's College London, Institute of Psychiatry, Psychology and Neuroscience Sanyal, Jyoti; South London and Maudsley Mental Health NHS Trust Colling, Craig; South London and Maudsley NHS Foundation Trust, Biomedical Research Centre Chandran, David; King's College London, Institute of Psychiatry Stewart, Robert; King's College London, Institute of Psychiatry
Primary Subject Heading:	Mental health
Secondary Subject Heading:	Mental health
Keywords:	Schizophrenia & psychotic disorders < PSYCHIATRY, Depression & mood disorders < PSYCHIATRY, Adult psychiatry < PSYCHIATRY, Health informatics < BIOTECHNOLOGY & BIOINFORMATICS, PSYCHIATRY

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

The association between depressive symptoms and Cognitive Behavioural Therapy receipt within a psychosis sample: A cross-sectional study.

Ava Mason¹, Jessica Irving¹, Megan Pritchard^{1,2}, Jyoti Sanyal², Craig Colling^{1,2}, David Chandran¹, Robert Stewart^{1,2}.

¹King's College London Institute of Psychiatry, Psychology and Neuroscience, De Crespigny Park, London, SE5 8AF

²South London and Maudsley NHS Foundation Trust, London, UK

* Address correspondence to: Ava Mason, University College London, Gower St, Bloomsbury, London WC1E 6BT, UK. Email: ava.mason.20@ucl.ac.uk.

Keywords: CBTp, psychosis, cognitive behaviour therapy, depression, mental illness.

Abstract word count: 248

Main body word count: 2918

Abstract

Objectives: To examine whether depressive symptoms predict receipt of cognitive behavioural therapy (CBTp) in individuals with psychosis.

Design: Retrospective cross-sectional analysis of electronic health records (EHRs) of a clinical cohort.

Setting: A secondary NHS mental health care service serving four boroughs of south London, UK.

Participants: 20,078 patients diagnosed with an ICD-10 code between F20-29 extracted from an EHR database.

Primary and secondary outcome measures: Primary: Whether recorded depressive symptoms predicted CBTp session receipt, defined as at least one session of CBT for psychosis (CBTp) identified from structured EHR fields supplemented by a natural language processing algorithm. Secondary: Whether age, gender, ethnicity, symptom profiles (positive, negative, manic and disorganisation symptoms), a comorbid diagnosis of depression, anxiety or bipolar disorder, general CBT receipt prior to the primary psychosis diagnosis date, or type of psychosis diagnosis predicted CBTp receipt.

Results: Of patients with a psychotic disorder, only 8.2% received CBTp. Individuals with at least one depressive symptom recorded, depression symptom severity and 12 out of 15 of the individual depressive symptoms independently predicted CBTp receipt. Female gender, White ethnicity and presence of a comorbid affective disorder or primary schizoaffective diagnosis were independently positively associated with CBTp receipt within the whole sample and the top 25% of mentioned depressive symptoms.

Conclusions: Individuals with a psychotic disorder who had recorded depressive symptoms were significantly more likely to receive CBTp sessions, aligning with CBTp guidelines of managing depressive symptoms related to a psychotic experience. However, overall receipt of CBTp is low and more common in certain demographic groups, and needs to be increased.

Strengths and limitations of the study

- To our knowledge, this is the first electronic health record (EHR) study to measure how clinical symptomatology predicts CBTp receipt, providing insight on a large sample into whether individuals who may be more in need of CBTp are more likely to have a session
- We replicate previous findings of inequalities in gender and ethnicity in real-world CBTp treatment receipt in a large heterogeneous sample.

- The natural language processing approach allows automated processing of EHR text at scale and can evaluate larger samples than manually conducted case note audits; this could therefore be used more routinely to monitor CBTp receipt.
- This study was limited to a single service provider; however, the results identified themes consistent with previous CBTp provision research in other services.
- Analysing EHRs in this way can identify CBTp receipt but is less suited to investigate whether CBTp is offered or not, or to quantify the quality or focus of the sessions. Furthering this, it cannot be used to examine CBTp completion rates and effectiveness.

Conflict of Interest

The authors report no conflict of interest with respect to the findings described in this manuscript. RS declares research support received in the last 36 months from Janssen, GSK and Takeda.

Financial Support

The independent research of this paper is funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at the South London and Maudsley NHS Foundation Trust and King's College London. RS is part-funded by: i) the National Institute for Health Research (NIHR) Biomedical Research Centre at the South London and Maudsley NHS Foundation Trust and King's College London; ii) a Medical Research Council (MRC) Mental Health Data Pathfinder Award to King's College London; iii) an NIHR Senior Investigator Award; iv) the National Institute for Health Research (NIHR) Applied Research Collaboration South London (NIHR ARC South London) at King's College Hospital NHS Foundation Trust. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

The Corresponding Author (AM) has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non-exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in BMJ editions and any other BMJ PGL products and sublicences such use and exploit all subsidiary rights, as set out in our licence.

Transparency declaration

The lead author AM affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Patient and public involvement

The Clinical Record Interactive system as a data resource was developed and is run with extensive patient involvement. However, this particular analysis did not involve patients in its design or implementation.

Ethics statement

The CRIS data platform has received research ethics approval as an anonymised data resource for secondary analysis (Oxford Research Ethics Committee C, reference 18/SC/0372)

Introduction

There are a variety of cognitive and emotional processes involved in the development of psychotic symptoms,¹ with intense distress emerging early on in the course of the disorder. Content of positive symptoms often mirrors the content of depressive thinking processes,² suggesting therapeutic need for individuals experiencing additional depressive symptoms. Specific depressive symptoms that often accompany psychotic disorders are hopelessness, social avoidance and problems in forming relationships.³ Around 50% of patients with psychosis report having experienced suicidal ideation at least once,⁴ and around 40% of individuals with schizophrenia report clinical levels of depression and low self-esteem.⁵ Importantly, individuals report these emotional difficulties and resulting social exclusion to be more debilitating than their psychotic symptoms.⁶ Consequentially, individuals' negative appraisal of their psychotic experiences may lead to loss of social goals and increased shame, predicting later hopelessness and post-psychotic depression.⁷ This comorbid depression increases the likelihood of having a lower quality of life, function, motivation, poorer social relationships, lower medication adherence and psychotic relapse.^{8,9} Therefore, treatment should focus on the psychotic symptoms and the broader distress they produce, building self-esteem, confidence and a sense of self control and purpose.¹⁰ Additionally, focusing on mood symptoms such as self-esteem and pessimism can help differentiate depressive symptoms from negative psychotic symptoms, that often show significant clinical overlap.⁵

It is increasingly recognised that medication alone is inadequate for tackling psychosis symptoms.¹¹ In the UK, the National Institute of Clinical Excellence¹² has recommended that cognitive behavioural therapy for psychosis (CBTp) be offered universally to individuals with psychosis. Based on the stress-vulnerability model,¹³ CBTp focuses on distress reduction related to hallucinations and delusions, through targeting negative beliefs and improving self-esteem.¹⁴ Sessions often focus on goal setting and emotional issues such as rebuilding one's self, positivity and acceptance.¹¹ While studies examining characteristics of CBTp show strong evidence that CBTp improves depressive symptoms in the context of psychosis, specifically with long term reductions in suicidal behaviour,^{14,15} service provision of this intervention still falls far short of the universal access recommended.¹¹

Considering the impact of targeting these symptoms in CBTp sessions, it is important to monitor receipt of CBTp within psychosis samples. While CBTp provision shows moderate yearly increases (12.8% in 2013 to 14.8% in 2014), the treatment is still only available to a small proportion of individuals,¹¹ short of NICE universal access recommendations.¹² Previous studies investigating CBTp receipt have conducted time-consuming audits on limited sample sizes; these can be affected by under-reporting. On the other hand, the UK's National Mental Health Minimum Data Set report does not require CBT interventions to be recorded in a given individual's record. Natural language processing techniques (NLP)¹⁶ offer the opportunity to extract this information from free text in electronic health records (EHRs) across large numbers of patients with psychosis, and a recent study developed and applied NLP in this respect, finding higher levels of receipt than reported in previous audit, supported by the high positive predictive value and sensitivity of the technique (95% and 96% respectively).¹¹

While studies have examined general CBTp receipt within patients with psychosis, no study has examined a link between depressive symptoms and CBTp receipt.¹¹ Therefore, we investigated whether depressive symptoms predict CBTp receipt in people with psychosis by applying these previously data extraction techniques to secondary mental health care EHRs for a large South London catchment population. Secondary predictors of receipt were type of psychosis diagnosis (schizophrenia, schizoaffective disorder or other schizophrenia spectrum disorder), symptom profiles (negative, manic or disorganisation), general CBT receipt prior to psychosis diagnosis, comorbid depression, anxiety or bipolar diagnosis and socio-demographic factors (ethnicity, gender and age).

Methods

For this study, we extracted data on individuals with a diagnosis of a recognised schizophrenia spectrum diagnosis from the case registry of the South London and Maudsley NHS Foundation Trust (SLaM). This is a large secondary care mental healthcare provider, serving around 1.3 million residents in Croydon, Lambeth, Lewisham and Southwark. SLaM care covers all specialist mental health care, including early intervention services, liaison and crisis teams and community and inpatient services. EHRs have been used for all SLaM services since 2006, with the Clinical Record Interactive Search system (CRIS) being established in 2008 to facilitate the retrieval of de-identified data from these records of patients previously or currently receiving mental healthcare from SLaM.¹⁷ The source EHR contains unstructured free text fields from correspondence, personal histories, mental health examinations and management plans, as well as structured fields for coding demographic information, like age and ethnicity. Implementing data from all these fields reduces selection bias of utilising only specific sources of information from the EHR. Consequently, a large programme of work has developed a range of NLP algorithms over the last decade, whose detailed descriptions and performance data are contained in an open-access catalogue.¹⁸ CRIS has approval as a data resource for secondary analysis (Oxford Research Ethics Committee C, reference 18/SC/0372), and a service user-led committee considers all proposed research before access to CRIS data is granted.

We extracted data for all individuals receiving SLaM care between January 2007 and June 2020 with a primary diagnosis of an ICD-10-defined schizophrenia spectrum disorder (F20-F29) and above the age of 18 at the time their original referral was accepted. The index date for covariate definitions was the date of the first diagnosis within this grouping. Individuals may have been active within the service before their index date, allowing us to extract data on prior CBT receipt. The sample was restricted to those with data on all variables.

Ethnicity, age at referral and gender were also extracted. Ethnicity was categorised into six groups for analysis: 'White British' (British), 'White other' (Irish or any other white background), 'Black' (Caribbean, African or any other black background), 'Asian' (Indian, Bangladeshi, Pakistani, Chinese or any other Asian background), 'Other/mixed' (white and Asian, white and black Caribbean, white and black African, any other ethnic group) and 'Not stated'.

Diagnosis was categorised into three subgroups of schizophrenia (ICD-10 codes F20.0–F20.9), schizoaffective disorder (F25.0–F25.9) and other schizophrenia spectrum disorder (F21, F22.0–F22.9, F23.0–F23.9, F24, F28 and F29). Within the data collection period, secondary diagnosis of depression (ICD-10: F32 or F33), anxiety (ICD-10: F40 or F41), or bipolar disorder (ICD-10: F31) were also extracted from structured field data.

NLP algorithms for each specific symptom were used to identify recorded symptom profiles within participants. Symptoms were categorised as depressive, positive, negative, manic or disorganisation. These symptoms had been categorised *a priori* by developers of the original independent symptom NLP algorithms. As symptoms could be labelled in more than one category during analysis, multicollinearity tests using the R function `vif()` within the [car package] were undertaken to avoid issues with overlapping predictor variables. All variables were included due to their VIF values being below five. However, positive symptoms were excluded from regression models using categorical symptom variables (having at least one mention within HER), as this factor variable only had one level, due to all participants having at least one positive symptom. The overall symptom list and subsequent recoding can be found in Table 1. Presence of at least one mention of any symptom in the five categories was computed as a binary variable (0/1).

Table 1
Classification of symptom predictors

Symptom	Symptom label
Aggression	Positive
Agitation	Positive
Anergia	Depressive /Negative
Anhedonia	Depressive/Negative
Apathy	Depressive/Negative
Arousal	Manic
Blunted affect	Depressive/Negative
Circumstantiality	Disorganisation
Delusions	Positive
Derailment	Disorganisation
Disturbed sleep	Depressive/Manic
Elation	Manic
Emotional Withdrawal	Negative
Flight of ideas	Disorganisation
Formal thought disorder	Disorganisation
Grandiosity	Manic
Guilt	Depressive
Hallucinations (auditory)	Positive
Helplessness	Depressive
Hopelessness	Depressive
Hostility	Positive
Insomnia	Depressive/Manic
Irritability	Manic
Paranoia	Positive
Persecutory ideation	Positive
Poor appetite	Depressive
Poor concentration	Depressive
Poor motivation	Depressive
Poverty of speech	Negative
Poverty of thought	Negative
Social withdrawal	Negative
Suicidal ideation	Depressive
Tangentiality	Disorganisation
Tearfulness	Depressive
Thought block	Disorganisation
Worthlessness	Depressive

The date of the first and last general CBT session before the index date was extracted. This was coded as a binary variable, with individuals in the 'Prior CBT' receipt group having at least one session date mention prior to their index date. This was included as a predictor to adjust for previous experience of the specific CBT intervention. Mentions were extracted using the same NLP tool as the CBTp outcome measure mentioned subsequently.

1
2
3 The primary outcome was CBTp receipt, identified using a combination of structured fields and NLP.¹⁶ The NLP
4 algorithms for general CBT has high PPV and sensitivity,¹² consistent with other NLP algorithms such as
5 medication dose and diagnosis.¹⁹ The date of the first CBTp session on or after the index date was extracted
6 and computed as a binary variable, so that individuals in the 'CBTp receipt' group had at least one CBTp
7 session mention after the index date.
8

9 10 **Statistical analysis**

11 To avoid overfitting, we followed the 'one in ten' rule, whereby one predictor can be measured for every 10
12 events. As the data included 1647 CBTp events, our study was able to include all 12 predictors within the same
13 regression model.
14

15 All statistical analyses were conducted using R (version 1.3.9). Descriptive statistics for demographic and
16 clinical variables are reported as frequencies for categorical variables and means and standard deviations for
17 the continuous variable (age at referral). Chi square tests were also calculated for categorical variables, and t-
18 test for age to measure between-group differences in those with/without CBT receipt. Descriptive statistics
19 were also provided for yearly CBT prior to index date and CBTp receipt post index date within the data
20 extraction time period (2007-2020).
21
22

23 Binary logistic regression was used to examine the association between depressive symptoms and receipt of at
24 least one CBT session in the whole sample. For this, three regression models were analysed. Model 1 was an
25 unadjusted model with only depressive symptoms as the predictor variable. Due to significant provision
26 differences seen in previous CBTp studies,¹¹ model 2 (partially adjusted model), adjusted for sociodemographic
27 variables (age at referral, ethnicity, gender), primary diagnosis group and presence of a comorbid diagnosis
28 (anxiety, depression and bipolar disorder). Model 3 (fully adjusted model) also adjusted for prior CBT receipt
29 before the index date (first psychosis diagnosis date) and symptoms mention (manic, negative and
30 disorganisation symptoms). Positive psychotic symptoms were not included in these models, as individuals all
31 had at least one mention within their case notes.
32
33

34 As the primary aim of the study was to investigate depressive symptoms as a predictor of CBTp receipt, we
35 also split the depressive symptoms category into the 15 specific depressive symptoms applications within the
36 whole sample. Model 4 was an unadjusted model with the 15 symptoms as predictor variables. Model 5 was a
37 fully adjusted model that adjusted for all the variables in Model 3. We also conducted a sensitivity analysis to
38 investigate how results were affected by overlap of negative or depressive symptom annotations, by removing
39 negative symptoms as a predictor from the logistic regression model.
40
41
42

43 Additionally to measuring whether individual depressive symptoms could predict CBTp receipt, we also also
44 measured whether overall depression severity predicted CBTp receipt. These logistic regression models
45 involved converting depressive, disorganised, manic, positive and negative symptoms into a continuous
46 variable, whereby severity reflected the number of different individual symptoms mentioned within each
47 symptom construct. This allowed for positive symptoms to also be included within regression models. Model 6
48 was an unadjusted model, with depressive symptom severity as a predictor of CBTp receipt. Model 7 and
49 model 8 were partially and fully adjusted models, controlling for the same variables as model 2 and 3, except
50 categorising symptoms as the continuous rather than categorical variable.
51
52
53

54 Lastly, to compare differences in the general sample with those with the top 25% quantity for depressive
55 symptoms, we conducted two further regression models. This subsample analysis was conducted to examine
56 predictors of CBTp receipt where a clear clinical indication was present, supplementing the overall findings.
57 Model 9 partially adjusted for socio-demographic factors, diagnostic group and comorbid diagnosis and Model
58 Model 10 fully adjusted for prior CBT, negative and disorganisation symptoms additionally. This group all had at least
59 one manic and psychotic symptom, so these variables were not included in the model.
60

Table 2*Distribution frequencies on baseline demographics and diagnoses split by CBTp receipt and primary diagnosis group.*

	No CBTp delivery (n = 18431)	CBTp delivery (n=1647)	Chi square tests (χ^2)
Ethnicity			$\chi^2=100.57^{***}$
White British	30% (5516/18431)	32.8% (540/1647)	
White Other	10.4% (1908/18431)	8.5% (140/1647)	
Black	36.5% (6719/18431)	41.7% (687/1647)	
Asian	6.5% (1193/18431)	5.2% (86/1647)	
Other/Mixed	9.8% (1808/18431)	10.5% (173/1647)	
Not stated	7.0% (1287/18431)	1.3% (21/1647)	
Gender			$\chi^2= 2.75$
Female	41.4% (7636/18431)	43.5% (717/1647)	
Male	58.6% (10795/18431)	56.5% (930/1647)	
Bipolar diagnosis	4.4% (810/18431)	9.0% (149/1647)	$\chi^2=71.94^{***}$
No diagnosis	95.6% (17621/18431)	91.0% (1498/1647)	
Depression diagnosis	7.4% (1373/18431)	14.0% (230/1647)	$\chi^2=87.36^{***}$
No diagnosis	92.6% (17058/18431)	86.0% (1417/1647)	
Anxiety diagnosis	2.4% (441/18431)	7.0% (115/1647)	$\chi^2=118.28^{***}$
No diagnosis	97.6% (17990/18431)	93.0% (1532/1647)	
Prior CBT	3.1% (573/18431)	14.4% (237/1647)	$\chi^2=497^{***}$
No prior cbt	96.9% (17858/18431)	85.6% (1410/1647)	

***p<.001

Results

Participants

The cohort comprised 20,078 individuals with the inclusion diagnoses, 1647 (8.2%) of whom received at least one session of CBTp after their first diagnosis date. The mean age of the cohort was 42.4 years (SD=16.5). Distribution frequencies for all categorical variables can be found in Table 2. Chi-square test results represented in this table compared those with or without CBTp receipt. All mentioned variables showed significant between-group differences at p<.001 apart from gender (No CBTp group females=41.4%, CBTp delivery group females= 43.5%; $\chi^2=2.75$, p=.097). These significant variables include depression diagnosis ($\chi^2=87.36$), bipolar diagnosis ($\chi^2=71.94$), anxiety diagnosis ($\chi^2=118.28$) and prior CBT receipt ($\chi^2=497$). Additionally, the Welch two sample t-test found significant between-group differences in age (t=15.34, p<.01). Where those who had received CBTp had a lower mean age (M=33.12 SD= 11.5) compared to those who did not (M=35.88, SD=13.08). The significant results confirmed the need for further analysis through the regression models. Positive psychotic symptoms were excluded from chi square and regression analysis, as all patients had at least one positive psychotic symptom.

CBT receipt.

The descriptive results shown in table 3 and figure 1 (found in supplementary materials), suggest that there is a low prevalence of both prior CBT and CBTp post diagnosis across the years, with receipt reducing in recent years (2019-2020) compared to earlier years (2007) of the data extraction period.

Table 3

Distribution frequencies on CBT receipt (prior to diagnosis) and CBTp receipt (post diagnosis) per year of data extraction.

Year	CBT prior	CBT post	All CBT
2007	130	81	211
2008	89	146	235
2009	59	111	170
2010	48	107	155
2011	37	105	142
2012	39	96	135
2013	32	128	160
2014	25	143	168
2015	24	150	174
2016	29	115	144
2017	16	127	143
2018	16	114	130
2019	15	153	168
2020	2	71	73
Total	561	1647	2208

General depressive symptom mention regression analysis

Results from the unadjusted (model 1), partially adjusted (model 2) and fully adjusted regression (model 3) are displayed in Table 4. Regression model 1 found that general mention of at least one of 15 potential depressive symptoms significantly predicted CBTp receipt. Regarding model 2 and 3, individuals with at least one depressive, negative or disorganisation symptom mention, being of female gender, white ethnicity, prior CBT receipt and presence of a comorbid affective disorder independently positively associated with CBTp receipt.

Table 4

Unadjusted, partially and fully adjusted logistic regression models for CBTp receipt (Regression model 1,2 and 3) with categorical symptom measures.

		Unadjusted	Partially adjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)	OR(95%CI)
Depressive symptoms				
1+ depressive symptom mention	18286(91.1)	3.78(2.94-4.96)***	3.42 (2.58-4.60)***	2.00(1.10-3.20)***
Bipolar diagnosis				
Has f31 diagnosis	959(4.80)		0.52(0.33-0.71)***	0.32(0.12-0.52)***
Depression diagnosis				
Has f32 diagnosis	603(80)		0.52(0.36-0.67)***	0.33(0.16-0.49)**
Anxiety diagnosis				
Has f40/41 diagnosis	556(2.80)		0.89(0.66-1.11)***	0.73(0.49-0.97)***
Age	N/A		-0.03(-0.04- -0.03)***	-0.03(-0.03- -0.02)***
Gender				
Male			Reference category	
Female	8353(41.60)		0.20(0.09-0.31)***	0.20 (0.10-0.32)***
Ethnic group				
White British	6056(30.10)		Reference category	
White Other	2048(10.20)		-0.40(-0.60 - -0.21)***	-0.37(-0.57- -0.17)***
Black	7406(36.90)		-0.16(-0.28- -0.04)**	-0.24 (-0.36- -0.11)***
Asian	1279(6.40)		-0.49(-0.74- -0.26)***	-0.50 (-0.75- -0.27)***
Other/Mixed	1981(9.90)		-0.21(-0.40- -0.02)**	-0.18(-0.37- -0.01)*
Not Stated	1308(6.50)		-1.75(2.23- -1.22)***	-1.52(-2.00- -1.10)***
Primary diagnosis				
Schizophrenia	9845(49.00)		Reference category	
Schizoaffective disorder	2142(10.70)		0.04(-0.13-0.21)	0.01(-0.17-0.19)
Other schizophrenia spectrum	8091(40.30)		-0.10(-0.22-0.01)*	-0.02(-0.14 -0.10)
Negative symptoms				
1+ Negative symptom mention	13169(65.60)			0.75(0.59-0.92)***
Manic symptoms				
1+ Manic symptom mention	17945(89.40)			1.24(0.70-1.87)***
Disorganisation symptoms				
1+ Disorganisation symptom mention	11513(57.30)			0.31(0.18-0.44)***
CBT prior				
1+ prior CBT session				1.29(1.12-1.46)***

*p<.05, **p<.01, ***p<.001

Unadjusted (model 1): depressive symptom as a predictor with no adjusted covariates

Partially adjusted (model 2): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis

Fully adjusted (model 3): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, depressive symptoms, prior CBT, negative symptoms, disorganisation symptoms, manic symptoms.

Regression analysis with individual depressive symptoms

Results from the unadjusted (model 4) and fully adjusted (model 5) regression analyses for each of the 15 individual depressive symptoms are displayed in Table 5 (N=20078). Each symptom refers to presence of at least one mention in the patients notes compared to no mention. While all variables were significant in the unadjusted model at $p < .001$, the fully adjusted model reduced the significance of suicide ideation ($p < .01$) and disturbed sleep ($p < .01$), with anhedonia, anergia, apathy, and blunted affect becoming non-significant ($p > 0.05$).

Table 5

Unadjusted and fully adjusted logistic regression models for CBTp receipt with individual depressive symptoms as covariates (Regression model 4 and 5) for the overall sample.

		Unadjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)
Hopelessness	7345(36.60)	4.81(4.3-5.40)***	1.45(1.26-1.66)
Helplessness	3124(15.60)	4.03(3.62-4.50)***	1.55(1.37-1.76)***
Suicide ideation	9451(47.10)	4.11(3.66-4.63)***	1.25(1.09-1.44)**
Poor appetite	8044(40.10)	3.31(2.97-3.68)***	1.28(1.13-1.45)***
Poor motivation	8630(43.00)	4.34(3.87-4.86)***	1.43(1.24-1.64)***
Insomnia	6870(34.20)	3.74(3.35-4.15)***	1.4(1.24-1.58)***
Disturbed sleep	16667(83.00)	15.3(10.16-22.8)***	2.76(1.5-5.08)**
Poor concentration	12289(61.20)	8.16(6.81-9.77)***	2.33(1.9-2.85)***
Anhedonia	4047(20.20)	2.9(2.61-3.22)***	0.97(0.85-1.10)
Anergia	873(43.50)	2.63(2.20-3.15)***	0.98(0.80-1.20)
Apathy	4149(20.70)	2.21(1.98-2.46)***	0.93(0.82-1.05)
Guilt	8178(40.70)	4.6(4.1-5.15)***	1.49(1.30-1.70)***
Tearfulness	10951(54.50)	3.87(3.41-4.39)***	1.22(1.05-1.42)**
Blunted affect	6889(34.30)	2.66(2.41-2.95)***	0.91(0.80-1.03)
Worthlessness	2921(14.50)	3.94(3.53-4.40)***	1.37(1.21-1.56)***

* $p < .05$, ** $p < .01$, *** $p < .001$

Sensitivity analysis

The non-significant results of certain depressive symptoms (anhedonia and anergia) may have been due to their inclusion within the negative symptom category, causing over-adjustment of the model. To test this, sensitivity analysis was conducted, where the fully adjusted regression (model 3) did not include negative symptoms as a covariate. While all significant variables remained significant, non-significant results for anhedonia and apathy were still found. Therefore, we report the fully adjusted model with negative symptoms as a variable for both grouped and individual depressive symptom associations.

General depressive symptom severity regression analysis

Results from the unadjusted (model 6), partially adjusted (model 7) and fully adjusted regression (model 8) are displayed in Table 6. Regression model 6 found that depression symptom severity significantly predicted CBTp receipt. Regarding model 7 and 8, depression symptom severity, positive symptom severity, anxiety diagnosis, and being of older age or being of white ethnicity independently positive predicted CBTp receipt. Within model 7., being female also positively increased likelihood of CBTp receipt. Within model 8, negative symptom severity and prior CBT significantly predicted CBTp receipt additionally.

Table 6
Unadjusted, partially and fully adjusted logistic regression models for CBTp receipt (Regression model 1,2 and 3) with continuous symptom measures.

		Unadjusted	Partially adjusted	Fully adjusted
	N(%)	OR(95%CI)	OR(95%CI)	OR(95%CI)
Depressive symptoms				
Severity	18286(91.1)	0.29(1.31-1.35)***	0.27(1.20-1.44)***	0.23(0.13-0.33)***
Positive symptoms				
Severity	20078(100)		-0.18(0.75-0.92)***	-0.21(-0.31- -0.09)***
Bipolar diagnosis				
Has f31 diagnosis	959(4.80)		0.21(0.94-1.63)	0.15(-0.13-0.43)
Depression diagnosis				
Has f32 diagnosis	603(80)		-0.09(0.72-1.15)	-0.09(-0.33-0.15)
Anxiety diagnosis				
Has f40/41 diagnosis	556(2.80)		0.49(1.15-2.29)***	0.46(0.11-0.80)***
Age	N/A		-0.02(0.97-0.99)***	-0.02(-0.03- -0.01)***
Gender				
Male			Reference category	
Female	8353(41.60)		0.17(0.97-1.44)*	0.17(-0.03-0.36)
Ethnic group				
White British	6056(30.10)		Reference category	
White Other	2048(10.20)		-0.41(0.45-0.96)**	-0.44(-0.93- -0.07)***
Black	7406(36.90)		-0.25(0.63-0.97)**	-0.29(-0.51- -0.07)**
Asian	1279(6.40)		-0.66(0.32-0.80)***	-0.67(-1.13 - -0.23)***
Other/Mixed	1981(9.90)		-0.14(0.62-1.22)	-0.16(-0.50-0.18)
Not Stated	1308(6.50)		-0.92(0.02-2.42)	-0.79(-3.74-1.01)
Primary diagnosis				
Schizophrenia	9845(49.00)		Reference category	
Schizoaffective disorder	2142(10.70)		-0.08(0.69-1.21)	-0.11(-0.40-0.18)
Other schizophrenia spectrum	8091(40.30)		0.02(0.82-1.26)	-0.06(-0.15-0.28)
Negative symptoms				
Severity	13169(65.60)			0.06(-0.01-0.123)*
Manic symptoms				
Severity	17945(89.40)			-0.01(-0.13-0.12)
Disorganisation symptoms				
Severity	11513(57.30)			0.10(-0.05-0.25)
CBT prior				
1+ prior CBT session	1647(8.20)			0.62(0.34-0.89)***

*p<.05, **p<.01, ***p<.001

Unadjusted (model 1): depressive symptom severity as a predictor with no adjusted covariates

Partially adjusted (model 2): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, positive symptom severity

Fully adjusted (model 3): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, positive symptom severity, prior CBT, negative symptom severity, disorganisation

symptom severity and manic symptom severity.

Depressive symptom regression analysis within the top 25% number of depressive symptoms.

This sample comprised individuals with the top 25% number of depressive symptoms (5018 patients), defined to reflect those who might reasonably expect to receive CBT. The sample characteristics and regression analysis can be seen in Table 7. Results from the partially adjusted (model 9) and fully adjusted regression (model 10) are displayed in Table 7. Table 4 found that general mention of at least one of 15 potential depressive symptoms significantly predicted CBTp receipt. Regarding model 9, we found that individuals with at least one depressive, negative or disorganisation symptom mention, being of female gender, white ethnicity, prior CBT receipt and presence of comorbid bipolar disorder were positively associated with CBTp receipt.

Table 7

Partially and fully adjusted logistic regression models for CBTp receipt with individual depressive symptoms as covariates within top 25% quantity of depressive symptoms (Regression model 9 and 10).

	N(%)	Partially adjusted OR(95%CI)	Fully adjusted OR(95%CI)
Bipolar diagnosis			
Has f31 diagnosis	541(10.78)	0.19(-0.03-0.41)*	0.15(-0.08-0.38)
Depression diagnosis			
Has f32 diagnosis	885(17.63)	0.11(-0.08-0.30)	0.08(-0.90-1.30)
Anxiety diagnosis			
Has f40/41 diagnosis	270(5.38)	0.53 (0.25-0.80)***	0.47(0.19-0.75)***
Age			
M= 36.24(18-93)		-0.02(-0.02- -0.01)***	-0.02(-0.03- -0.01)***
Gender			
Male	2059(41.01)	Reference category	
Female	2960(58.99)	0.20 (0.05-0.34)***	0.17(0.02-0.32)**
Ethnic group			
White British	1486(29.59)	Reference category	
White Other	433(8.63)	-0.21(-0.48-0.05)	-0.22(0.50-0.05)
Black	2262(45.08)	-0.32(-0.49- -0.16)***	-0.31(0.47- -0.14)***
Asian	328(6.53)	-0.53(0.86- -0.21)***	-0.52(0.85- -0.20)***
Other/Mixed	467(9.31)	-0.08(-0.34-0.17)	-0.08(0.34-0.17)
Not Stated	43(0.86)	-1.42 (-2.85 -0.40)**	-1.34(-2.8 - -0.31)**
Primary diagnosis			
Schizophrenia	2219(44.22)	Reference category	
Schizoaffective disorder	740(14.72)	0.04(-0.18-0.26)	0.003(-0.22-0.22)
Other schizophrenia spectrum	2060(41.03)	0.02(-0.14 -0.17)	0.05(-0.11-0.21)
Negative symptoms			
1+ Negative symptom mention	4956(98.7)		-0.88(1.41- -0.33)***
Disorganisation symptoms			
1+ Disorganisation symptom mention	4199(83.66)		1.18(-0.02-0.38)*
CBT prior			
1+ prior CBT session	436(8.7)		-0.90(0.68-1.11)***

*p<.05, **p<.01, ***p<.001,

Partially adjusted (model 9): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis.

Fully adjusted (model 10): results were adjusted for age, ethnicity, gender, diagnostic group, f31 diagnosis, f32 diagnosis, f40/41 diagnosis, prior CBT, negative symptoms, disorganisation symptoms

Discussion

1
2
3
4 We believe that this is the first study to examine the relationship between clinical symptomatology and CBTp receipt
5 within a sample of people with psychosis in a naturalistic community setting. In general, only 8.2% of individuals
6 received CBTp within the 13-year timeframe of the study, showing the low prevalence of receipt despite current clinical
7 guidelines. This finding shows a reduction in CBTp provision compared to previous studies in 2013 (12.8%) and 2014
8 (14.8%)¹¹, which was further supported by the descriptive frequency results, showing a drop in both CBT and CBTp
9 receipt in recent years. This requires further examination, as it is unclear why receipt is decreasing considering the
10 importance of CBTp mentioned within NICE universal access recommendations.¹²
11

12 91% of patients had at least one recorded depressive symptom mention. Individuals with at least one depressive
13 symptom mention were 2 times more likely to have at least one CBTp session in the fully adjusted model (table 4),
14 suggesting that the minority who don't present with any depressive symptoms are very unlikely to receive CBTp. This
15 could possibly be due to clinicians tending to cite a depressive symptom when referring an individual with psychosis to
16 psychotherapy. Additionally, the severity of depressive symptoms, as well as having at least one recorded mention
17 significantly increased likelihood of having at least one CBTp session. In the sample of those with the highest number of
18 depressive symptoms (top 25%), relationships between CBTp receipt and comorbid anxiety diagnosis, age, gender,
19 ethnicity, prior CBT, negative and disorganised psychotic symptoms remained (effect size ranging from 0.08-1.34). This
20 suggests the importance of these predictors in a reasonable sample of patients with higher clinical need for CBTp
21 receipt.
22

23 Overall there was therefore a low prevalence of CBTp receipt within those with one depressive symptom. The
24 depressive symptom which was the strongest predictor of this intervention in fully adjusted models was
25 disturbed sleep. There is a known high prevalence of sleeping problems in this population,^{20,21} described by
26 some researchers as an 'intrinsic feature of schizophrenia,²² known to reduce quality of life, decreasing coping
27 and exacerbate positive symptoms.²³ *The significant association between insomnia and psychotic-like*
28 *symptoms, such as paranoia, has also been seen in non-clinical populations.*²⁴ Furthering this, the
29 recommended first line of treatment for sleep problems in this sample is CBT.²⁵ Poor concentration was the
30 next strongest depressive symptom predictor in the fully-adjusted model, supporting previous research of its
31 association with psychosis vulnerability.²⁶ The significance of helplessness, guilt and hopelessness mirrors CBTp
32 research that found significant post-treatment reduction in hopelessness, self-depreciation and guilt using the
33 Calgary Depression Rating Scale for Schizophrenia.²⁷ Other significant depressive symptoms associated with
34 low self-esteem and negative self-evaluation and emotions have been found to significantly affect the
35 development and severity of positive symptoms.²⁸ This may be because positive symptoms develop as a
36 psychological defence against low self-esteem²⁹ and depression-induced guilt.³⁰ Therefore, it could be
37 suggested that the significance of each of the depressive symptoms is often linked to psychotic symptoms and
38 CBTp effectiveness. However, while there is evidence on the clinical impact of depressive symptoms in
39 schizophrenia, the associations with choice of therapy must be viewed as exploratory and in need of
40 independent replication. While a possibility may be that clinicians are assuming that certain depressive
41 symptoms are likely to be more responsive to CBTp than others, there may be other unknown reasons for
42 therapy choice that requires further investigation. General results suggest that receipt of this intervention
43 requires an increase for all of this population before individuals with these specific symptoms could be
44 targeted.
45
46

47 Regarding negative symptoms, the non-significant associations between specific negative symptoms (that overlapped
48 with depressive symptoms) and CBTp receipt requires specific further testing. This was not conducted in the current
49 study due to the primary aim focusing on depressive symptoms. However, from our results on specific depressive
50 symptoms in table 5, symptoms that overlapped with negative symptoms (anhedonia, anergia, apathy and blunted
51 affect) were not associated with CBTp receipt. Additionally, negative symptoms significantly decreased likelihood of
52 CBTp receipt within the top 25% of individuals with a depressive symptom mention. Overall, this raises concerns that
53 individuals with these specific negative/depressive symptoms are no more likely and perhaps less likely to receive CBTp
54 than someone without these symptoms. Possibly, this is due to clinicians not tending to refer these individuals because
55 they don't believe intervention would be effective. This is in line with a CBTp review of randomised control trials,
56 finding non-significant reductions of negative symptoms,³¹ perhaps due to the narrowing of treatments to specifically
57 target positive symptoms.³² However, further work should be undertaken to verify that individuals are not being
58 denied a potentially beneficial intervention because of their symptom profile.
59
60

1
2
3 Prior CBT receipt, comorbid disorder presence and specific symptoms (manic, disorganised and negative) also emerged
4 as independent predictors of CBTp receipt for the general sample and within those with the top 25% depressive
5 symptom numbers. Within table 4, individuals who had any recorded CBT receipt prior to the index date were 1.29
6 times more likely to have recorded CBTp receipt later on. Also, patients with an additional comorbid disorder were
7 0.32-0.73 times more likely to have received CBTp compared to those with just a psychosis diagnosis. However, in the
8 top 25% of individuals within table 7, individuals with prior CBT were 0.40 times less likely to receive CBTp. This finding
9 requires further research to understand the effects of prior CBT and negative symptoms on CBTp receipt within
10 different psychosis subsamples. Additionally, those with anxiety were 0.47 times more likely to receive CBTp and those
11 with disorganised symptoms were 1.18 times more likely respectively. After general CBTp receipt has increased, there
12 could be a method to focus more on patients with different types of psychotic symptoms and comorbid affective
13 diagnosis. Furthering this, future research could investigate whether those who have had prior general CBT would
14 benefit from CBTp, or whether those who have not had any experience developing cognitive behaviour skills in therapy
15 should be targeted.³³

17 Crucially, there were also significant differences in CBTp receipt between different ethnic and gender groups. Male
18 patients were 0.20 times less likely in the general sample and top 25% of depressive symptoms to have recorded CBTp
19 receipt. Black, Asian, Other and Mixed ethnic groups were between 0.21 to 0.49 times less likely to have a documented
20 CBTp session compared to individuals of white ethnicity within both the general and top 25% depressive symptoms
21 samples. Inequitable access to CBTp has been identified in previous CBTp research within a psychosis sample drawn
22 from the same data resource in 2017, finding female patients to be more likely to have received CBTp and individuals of
23 White ethnicity to have a significantly higher likelihood of CBT receipt than Black or other ethnicity groups.¹¹ This also
24 supports results from a recent CBTp study focusing specifically on ethnic group differences in CBTp provision within
25 SLaM, who found that in comparison to White British individuals, those from Black ethnic groups with psychosis or
26 bipolar disorder were significantly less likely to have a documented CBTp session. This is especially important when
27 considering the high prevalence of psychosis within UK BAME populations.³³ Inequality in CBTp receipt may be due to
28 ethnic variations in CBTp engagement. Some of these barriers within certain communities may be increased stigma,
29 fear of clinicians by service-users or service users by clinicians, institutional racism within mental health services, or
30 non-culturally appropriate therapy.³⁴ As differences in documented CBTp receipt between ethnic groups have now
31 been documented by three different papers in this service, it is imperative that further work is conducted to increase
32 provision of CBTp within groups less likely to receive treatment. This may include targeted outreach programs and
33 culturally adapting interventions³⁴ within these minority groups.

35 The present study has a number of strengths and limitations. Generally, focusing on patients with more
36 diverse functioning, comorbidity and symptom severity levels helps research identify a larger number of
37 predictors of clinical outcomes. This can be seen through our results, where negative, manic and
38 disorganisation symptoms significantly predicted CBTp receipt, as well as recent heterogenous research,³²
39 that was the first to identify depression as a significant predictor of positive symptom improvement post-
40 CBTp. This highlights the importance of focusing on a clinically heterogeneous sample to realistically determine
41 significant predictors of CBTp receipt. Secondly, using an NLP approach automates the measurement of what
42 would otherwise require manually conducted audits on records and case notes, increasing the number of
43 cases that can be investigated and providing a method that could be used more routinely to monitor CBT
44 receipt. The large sample size enabled us to identify clinical differences in the real-life administration of CBTp
45 within a psychosis cohort, and we were able to adjust for multiple clinical variables and comorbidity diagnoses
46 to provide a more realistic understanding of the depressive symptom-CBTp receipt relationship. This time
47 frame was broad to allow the inclusion of as many active patients receiving CBTp as possible, additionally
48 circumventing monthly/seasonal variation of CBTp receipt.

50 One limitation of the study was the omission of strict time periods for the mention of clinical symptoms prior
51 to CBTp administration. Unfortunately, using this approach would have involved implementing time periods on
52 all of the other clinical symptoms and variables, which would have been difficult considering the number of
53 variables that would need to be controlled. In addition, the NLP symptom algorithms do not currently
54 distinguish between past or present symptoms. Therefore, symptom mentions documented after the CBTp
55 receipt date could refer to mentions of symptoms occurring prior to CBTp receipt, reducing the effectiveness
56 of using time periods. A follow-up time period after the index date was also not established, meaning that
57 participants included in the cohort at a later date may have been less likely to have had a CBTp session, due to
58 their limited time period within the service. Additionally, we did not have data regarding which type of service
59 was providing CBTp for each patient (for example, early intervention services compared to other community
60

1
2
3 services). Future studies should examine whether CBTp receipt differs depending on the service, especially
4 considering how effective CBTp provision may be in those at ultra-high risk.
5

6 While our use of additional querying of text fields allowed us to identify a significantly larger number of CBTp
7 episodes than using structured data alone, we were not able to quantify the gap between CBTp referral and
8 CBTp receipt. This is because the CBTp NLP algorithm detects CBTp receipt rather than CBTp being offered,
9 due to the wide range of subtle wording used for the latter more complex entity. The results combine effects
10 on the likelihood of CBTp being offered, with those on session receipt following an offer. While this may have
11 affected our results, previous service audits have suggested that the severity and occurrence of depressive
12 symptoms significantly decreases CBT receipt.³⁵ Therefore, if only receipt was directly measured, we would
13 expect to see similar results. Additionally, completion rates and effectiveness of the CBTp was not measured,
14 meaning we were unable to quantify the quality or focus of the sessions. Lastly, analysis was limited to
15 patients above 18 years old, reducing the generalisability of results to those who develop a schizophrenia-
16 spectrum disorder after this age. However, the outcome of interest was CBTp receipt within a relatively
17 homogenous service structure of working age services, rather than young people treated within Child and
18 Adolescent services. Future studies should examine whether CBTp receipt differs in these services.
19

20 **Future directions**

21
22 Initiatives such as the Improving Access to Psychological Therapies programme for serious mental illness, early
23 intervention access and projects to decrease waiting times for referral have been developed to target this
24 clinical population. However, access still falls short of recommendations and is inequitable for specific
25 psychotic diagnoses, age and ethnicity.¹¹ Therefore, given the effect of CBTp on depressive symptoms³⁶,
26 perhaps its more pragmatic to focus on patients with additional depressive symptoms. Monitoring CBTp
27 receipt over time could decipher whether these initiatives are effective at increasing general access for those
28 with psychosis, and specific access for different sociodemographic groups and those with additional depressive
29 symptoms (who may benefit the most).
30

31
32 The significant secondary clinical and sociodemographic variables require further analysis in order to fully
33 understand the services' provision. This could involve attention given to the independent symptoms within the
34 negative, manic and disorganisation categories in a similar manner to the specific depressive symptom
35 regression models analysed. Further research could also explore why the presence of co-morbid anxiety and
36 bipolar disorder in this sample predicted CBTp receipt. Additionally, the results suggest a need to reflect on the
37 steps taken since the previous service study,³⁴ regarding inequality in CBTp receipt among gender and ethnic
38 groups, due to the consistent significant results seen. Regarding the use of EHR data, future work could involve
39 developing a separate NLP algorithm to ascertain the offering of CBTp or provide another structured field for
40 clinicians to complete for this. However, additional text fields seem an unlikely approach, as clinicians prioritise
41 text field data for communication about CBTp sessions for themselves and their colleagues rather than to
42 collect structure data for the sake of research. Therefore, as previously suggested,¹¹ it is important to accept
43 the mixed structured-text field approach that will remain in healthcare record data and perhaps our time is
44 best spent in improving NLP algorithms to detect the subtleties of intervention and clinical outcome data.
45 However, the implications of our results and their consistency three years after the first CBTp service paper
46 suggest the need to use this or future algorithms for service monitoring independent of these improvements.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Chadwick P, Birchwood M. The omnipotence of voices. A cognitive approach to auditory hallucinations. *Br J Psychiatry* 1994;164:190–201.
2. Fowler D. Cognitive Behavior Therapy for Psychosis: From Understanding to Treatment. *Am J Psychiatr Rehabil* 2000;4(2):199–215.
3. Hazell CM, Hayward M, Cavanagh K et al. Clara Strauss A systematic review and meta-analysis of low intensity CBT for psychosis. *Clinical Psychology Review* 2016;45:183–192.
4. Hawton, K, Sutton, L, Haw, C, Sinclair, J, Deeks, JJ. Schizophrenia and suicide: systematic review of risk factors. *Br. J. Psychiatry* 2005;187:9–20.
5. Sands JR, Harrow M. Depression during the longitudinal course of schizophrenia. *Schizophr Bull* 1999;25:157–171.
6. Greenwood KE, Sweeney A, Williams S, et al. CHOICE of Outcome In Cbt for psychosEs (CHOICE): The Development of a New Service User–Led Outcome Measure of CBT for Psychosis. *Schizophrenia Bulletin* 2010;36(1):126–135.
7. Birchwood M, Iqbal Z, Chadwick P, et al. Cognitive approach to depression and suicidal thinking in psychosis. I: Ontogeny of post-psychotic depression. *Br J Psychiatry* 2000;177: 516–528.
8. Conley RR. The burden of depressive symptoms in people with schizophrenia. *Psychiatr Clin North Am* 2009;32:853–861.
9. Rocca P, Bellino S, Calvarese P, et al. Depressive and negative symptoms in schizophrenia: different effects on clinical features. *Comprehensive Psychiatry* 2005; 46: 304–310.
10. Pitt L, Kilbride M, Nothard S, et al. Researching recovery from psychosis: a user-led project. *Psychiatr Bull R Coll Psychiatr* 2007;31:55–60.
11. Colling C, Evans L, Broadbent M, et al. Identification of the delivery of cognitive behavioural therapy for psychosis (CBTp) using a cross-sectional sample from electronic health records and open-text information in a large UK-based mental health case register. *BMJ Open* 2017;7:e015297.
12. National Institute for Health and Clinical Excellence. Schizophrenia: Core interventions in the treatment and management of schizophrenia in primary and secondary care. London, UK: NICE Clinical Guideline 2009.
13. Rathod S, Phiri P, Kingdon D. Cognitive behavioral therapy for schizophrenia. *Psychiatr Clin North Am* 2010;33:527–536.
14. Gumley A, Schwannauer M. Staying Well after Psychosis: A Cognitive Interpersonal Approach to Recovery and Relapse Prevention. New York, NY: John Wiley & Sons Ltd 2006.
15. Lincoln TM, Rief W, Wetermann S, et al. Who stays, who benefits? Predicting dropout and change in cognitive behaviour therapy for psychosis. *Psychiatry Research* 2014;216(2):198–205.
16. Spyns P. Natural language processing in medicine: an overview. *Methods Inf Med* 1997;35(4-5):285–301.
17. Stewart R, Soremekun M, Perera G, et al. The South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLAM BRC) case register: development and descriptive data. *BMC Psychiatry* 2009;9:51.
18. (<https://www.maudsleybrc.nihr.ac.uk/media/325736/applications-library-v13.pdf>).
19. Hayes RD, Downs J, Chang CK, et al. The effect of clozapine on premature mortality: an assessment of clinical monitoring and other potential confounders. *Schizophr Bull* 2015;41:644–55.
20. Chadwick PDJ, Birchwood MJ, Trower, P. Cognitive Therapy for Delusions, Voices and Paranoia: Towards A Person Model. Chichester: Wiley 1996.
21. Chiu V, Ree M, Janca A, et al. Sleep profiles and CBT-I response in schizophrenia and related psychoses. *Psychiatry Research* 2018;268:279–287.
22. Chouinard S, Poulin J, Stip E, et al. Sleep in untreated patients with schizophrenia: a meta-analysis. *Schizophr Bull* 2004;30:957–967.

23. Waite F, Evans N, Myers E, et al. The patient experience of sleep problems and their treatment in the context of current delusions and hallucinations. *Psychology and Psychotherapy: Theory, Research and Practice* 2016;89:181–193.
24. Freeman D, Sheaves B, Goodwin G, et al. The effects of improving sleep on mental health (OASIS): a randomised controlled trial with mediation analysis. *The Lancet Psychiatry* 2017;4(10):749–758.
25. Mitchell MD, Gehrman P, Perlis M, et al. Comparative effectiveness of cognitive behavioral therapy for insomnia: A systematic review. *BMC Family Practice* 2012;13(1):40.
26. Bogren M, Mattisson C, Tambs K, et al. Predictors of psychosis: a 50-year follow-up of the Lundby population. *Eur Arch Psychiatry Clin Neurosci* 2009;260:113–125.
27. Muller MJ, Marx-Dannigkeit P, Schlösser R, et al. The Calgary Depression Rating Scale for Schizophrenia: Development and interrater reliability of a German version (CDSS–G). *J Psychiatr Res* 1999;33:433–443.
28. Smith B, Fowler DG, Freeman D, et al. Emotion and psychosis: links between depression, self-esteem, negative schematic beliefs and delusions and hallucinations. *Schizophr Res* 2006;86(1-3):181–188.
29. Bentall RP, Kinderman P, Kaney S. The self, attribution processes and abnormal beliefs: Towards a model of persecutory delusions. *Behav Res Ther* 1994;32:331–341.
30. Lake CR. Hypothesis: Grandiosity and Guilt Cause Paranoia; Paranoid Schizophrenia is a Psychotic Mood Disorder; a Review. *Schizophr Bull* 2008;34(6):1151–1162.
31. Lincoln TM, Ziegler M, Mehl S, et al. Moving from efficacy to effectiveness in cognitive behavioral therapy for psychosis: a randomized clinical practice trial. *J Consult Clin Psychol* 2012;80(4):674–686.
32. Penn DL, Meyer PS, Evans E, et al. A randomized controlled trial of group cognitive-behavioral therapy vs. enhanced supportive therapy for auditory hallucinations. *Schizophr Res* 2009;109:52–59.
33. Morris R, Sellwood W, Edge D, et al. Ethnicity and impact on the receipt of Cognitive Behavioural Therapy in people with psychosis or bipolar disorder: An English cohort study. *BMJ Open* 2020; accepted.
34. Rathod S, Phiri P, Harris S, et al. Cognitive behaviour therapy for psychosis can be adapted for minority ethnic groups: A randomised controlled trial. *Schizophr Res* 2013;143, 319–326.
<http://dx.doi.org/10.1016/j.schres.2012.11.007>
35. Binnie J, Boden Z. Non-attendance at psychological therapy appointments. *Mental Health Review Journal* 2016;21(3):231–248.
36. Taalman H, Goldberg DM, Ayub M, et al. Effect of cognitive behaviour therapy for psychosis (Cbtp) on depressive symptoms: a review of literature. *J Schizophr Res* 2015;2(3):1019.

Acknowledgments

This paper would not have been completed without the assistance from Dr Rob Stewart. I would also like to thank Jessica Irving for her assistance in the statistical analysis and general manuscript.

Author statement

Ava Mason planned the protocol, analysed the data and wrote the manuscript. Dr Rob Stewart provided access to the data, looked over edits and revisions on the manuscript. The data from this paper was accessed through the assistance of Megan Pritchard and Jyoti Sanyal. The applications used to obtain data specific to CBTp were developed by Craig Colling and David Chandran. Jessica Irving assisted in statistical analysis.


Data Sharing Statement

Due to the terms of Ethics and Information Governance approvals and clinical source of the data, CRIS datasets must remain within the South London and Maudsley NHS Foundation Trust (SLaM) firewall. All data used from this study can be made accessible on request from cris.administrator@slam.nhs.uk, subject to the setting up of an appropriate research passport or SLaM honorary contract.

Figure 1: Graph demonstrating the frequency of general CBT receipt (CBT receipt prior to diagnosis and recorded CBTp receipt post diagnosis) per year of extraction period.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only



20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary figure

Graph demonstrating the frequency of general CBT receipt (CBT receipt prior to diagnosis and recorded CBTp receipt post diagnosis) per year of extraction period.

Reporting checklist for cross sectional study.

Based on the STROBE cross sectional guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the STROBE cross sectional reporting guidelines, and cite them as:

von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies.

		Reporting Item	Page Number
Title and abstract			
Title	#1a	Indicate the study's design with a commonly used term in the title or the abstract	1
Abstract	#1b	Provide in the abstract an informative and balanced summary of what was done and what was found	1
Introduction			
Background / rationale	#2	Explain the scientific background and rationale for the investigation being reported	1,3
Objectives	#3	State specific objectives, including any prespecified hypotheses	3
Methods			
Study design	#4	Present key elements of study design early in the paper	4

1	Setting	#5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	4
2				
3				
4	Eligibility criteria	#6a	Give the eligibility criteria, and the sources and methods of selection of participants.	4
5				
6				
7				
8		#7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	1,4-6,
9				
10				
11				
12				
13				
14	Data sources /	#8	For each variable of interest give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group. Give information separately for for exposed and unexposed groups if applicable.	4-6
15	measurement			
16				
17				
18				
19				
20				
21	Bias	#9	Describe any efforts to address potential sources of bias	4
22				
23	Study size	#10	Explain how the study size was arrived at	4
24				
25	Quantitative	#11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen, and why	4-6
26	variables			
27				
28				
29	Statistical	#12a	Describe all statistical methods, including those used to control for confounding	6
30	methods			
31				
32				
33	Statistical	#12b	Describe any methods used to examine subgroups and interactions	6
34	methods			
35				
36				
37	Statistical	#12c	Explain how missing data were addressed	4
38	methods			
39				
40				
41	Statistical	#12d	If applicable, describe analytical methods taking account of sampling strategy	6
42	methods			
43				
44	Statistical	#12e	Describe any sensitivity analyses	6
45	methods			
46				
47				
48	Results			
49				
50				
51	Participants	#13a	Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed. Give information separately for for exposed and unexposed groups if applicable.	1,6
52				
53				
54				
55				
56				
57				
58				
59				
60				

1	Participants	#13b	Give reasons for non-participation at each stage	6
2				
3	Participants	#13c	Consider use of a flow diagram	N/A
4				
5	Descriptive data	#14a	Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders. Give information separately for exposed and unexposed groups if applicable.	4-7
6				
7				
8				
9				
10				
11				
12	Descriptive data	#14b	Indicate number of participants with missing data for each variable of interest	4
13				
14				
15				
16	Outcome data	#15	Report numbers of outcome events or summary measures. Give information separately for exposed and unexposed groups if applicable.	8
17				
18				
19				
20				
21	Main results	#16a	Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	7-12
22				
23				
24				
25				
26				
27				
28	Main results	#16b	Report category boundaries when continuous variables were categorized	7-12
29				
30				
31				
32	Main results	#16c	If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
33				
34				
35				
36	Other analyses	#17	Report other analyses done—e.g., analyses of subgroups and interactions, and sensitivity analyses	12
37				
38				
39				
40	Discussion			
41				
42	Key results	#18	Summarise key results with reference to study objectives	13-15
43				
44	Limitations	#19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias.	14-15
45				
46				
47				
48				
49				
50	Interpretation	#20	Give a cautious overall interpretation considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence.	13-15
51				
52				
53				
54				
55	Generalisability	#21	Discuss the generalisability (external validity) of the study results	13-15
56				
57				
58				
59				
60				

1 **Other**
2 **Information**

3
4
5 Funding [#22](#) Give the source of funding and the role of the funders for the present 2
6 study and, if applicable, for the original study on which the present
7 article is based
8
9

10 The STROBE checklist is distributed under the terms of the Creative Commons Attribution License CC-BY.
11 This checklist was completed on 11. March 2021 using <https://www.goodreports.org/>, a tool made by the
12 [EQUATOR Network](#) in collaboration with [Penelope.ai](#)
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60