

**Supplemental Table 1:** Pain scores on NPRS at each time point (n=10).

<b>Baseline</b>	<b>5 min</b>	<b>10 min</b>	<b>15 min</b>	<b>30 min</b>	<b>45 min</b>	<b>60 min</b>	<b>120 min</b>	<b>180 min</b>	<b>240 min</b>
<b>Visit one</b>									
7.5 (6-8)	6 (5 - 8)	6 (5 - 8)	5 (4 - 7)	4.5 (2 - 7)	5 (3 - 7)	4.5 (3 - 6)	4.5 (3 - 7)	5 (3 - 6)	5 (3 - 7)
Change from baseline	0.5 (0 - 2) p=0.063	0.5 (0 - 3) p=0.062	<b>1 (1 - 2)</b> p=0.004	<b>1 (1 - 5)</b> p=0.004	<b>1 (1 - 4)</b> p=0.008	<b>1.5 (1 - 4)</b> p=0.002	<b>1.5 (0 - 3)</b> p=0.016	<b>2 (0 - 3)</b> p=0.016	<b>1.5 (1 - 2)</b> p=0.008
<b>Visit two</b>									
6 (5 - 8)	<b>1 (0 - 3)</b>	<b>0 (0 - 3)</b>	<b>1.5 (0 - 3)</b>	<b>2.5 (0 - 5)</b>	<b>3 (0 - 5)</b>	<b>3.5 (1 - 5)</b>	<b>3.5 (1 - 4)</b>	<b>4 (1 - 5)</b>	<b>3 (1 - 4)</b>
Change from baseline	<b>4 (3 - 7)</b> p=0.002	<b>5.5 (3 - 7)</b> p=0.002	<b>4.5 (3 - 6)</b> p=0.002	<b>2.5 (1 - 6)</b> p=0.004	<b>2.5 (1 - 5)</b> p=0.004	<b>2 (1 - 4)</b> p=0.004	<b>2 (2 - 4)</b> p=0.008	<b>1.5 (1 - 3)</b> p=0.008	<b>2 (2 - 3)</b> p=0.008
<b>Visit three</b>									
5.5 (5 - 7)	5 (4 - 6)	<b>3 (3 - 6)</b>	<b>3.5 (3 - 6)</b>	<b>3.5 (2 - 6)</b>	<b>3 (2 - 4)</b>	<b>2.5 (2 - 3)</b>	<b>3 (2 - 3)</b>	<b>3 (2 - 3)</b>	<b>3 (3 - 4)</b>
Change from baseline	0.5 (0 - 1) p=0.06	<b>2 (2 - 3)</b> p=0.04	<b>2 (1 - 3)</b> p=0.004	<b>2 (1 - 3)</b> p=0.004	<b>2 (2 - 3)</b> p=0.004	<b>3 (2 - 3)</b> p=0.002	<b>2 (2 - 4)</b> p=0.002	<b>2 (1 - 5)</b> p=0.004	<b>2 (1 - 2)</b> p=0.008
<b>Visit four</b>									
6 (6 - 8)	<b>5 (3 - 5)</b>	<b>4 (2 - 5)</b>	<b>3 (2 - 5)</b>	<b>2 (2 - 5)</b>	<b>2 (2 - 4)</b>	<b>2 (1 - 5)</b>	<b>2 (1 - 5)</b>	<b>2.5 (1 - 6)</b>	<b>5 (2 - 6)</b>
Change from baseline	<b>1 (0 - 2)</b> p=0.047	<b>2 (1 - 3)</b> p=0.008	<b>4 (3 - 4)</b> p=0.004	<b>4 (2 - 4)</b> p=0.004	<b>4 (4 - 4)</b> p=0.008	<b>4 (3 - 5)</b> p=0.008	<b>4 (2 - 5)</b> p=0.008	<b>3 (1 - 5)</b> p=0.008	<b>1 (1 - 4)</b> p=0.008

Data presented as median (interquartile range). Shapiro-Wilk test used to test for normality and more than 80 percent of the time point values were found to have not normal distribution. Therefore, all P-values presented here are for Wilcoxon Signed Rank test for not normally distributed data. Data points with p-values <0.05 are bolded.