

SUPPLEMENTARY INFORMATION

Multistage Antiplasmodial Activity of Hydroxyethylamine Compounds, *In vitro* and *In vivo* Evaluations

Neha Sharma¹, Yash Gupta² Meenakshi Bansal¹, Snigdha Singh¹, Prateek Pathak³, Mohd Shahbaaz^{3,4}, Raman Mathur², Jyoti Singh⁵, Mohammad Kashif⁵, Maria Grishina³, Vladimir Potemkin³, Vinoth Rajendran⁶, Poonam^{3,7}, Prakasha Kempaiah², Agam P Singh^{5,*}, Brijesh Rathi^{1,3,*}

¹Laboratory for Translational Chemistry and Drug Discovery, Department of Chemistry, Hansraj College University Enclave, University of Delhi, Delhi 110007 India

²Department of Medicine, Loyola University Stritch School of Medicine, 2160 South First Avenue, Chicago, Illinois 60153, United States

³South Ural State University, Laboratory of Computational Modelling of Drugs, 454080, Russia

⁴South African National Bioinformatics Institute, University of the Western Cape, Private Bag X17, Bellville, Cape Town 7535, South Africa

⁵Infectious Diseases Laboratory, National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi 110067, India

⁶Department of Microbiology, School of Life Sciences, Pondicherry University, Puducherry 605014, India

⁷Department of Chemistry, Miranda House, University of Delhi, Delhi 110007 India.




Fig S1. ¹H NMR spectra of compound 1 in CDCl₃.




Fig S2. ¹³C NMR spectra of compound 1 in CDCl₃.




Fig S3. 2D COSY NMR spectra of compound 1 in CDCl₃.




Fig S4 ¹H NMR spectra of compound 2 in CDCl₃.




Fig S5. 2D COSY NMR of Compound 2 in CDCl₃.