

**Supplementary Table 2.** Assessment of Difficulties in Doing Certain Activities among the 170 Respondents at 12 Months after Acute COVID-19 Infection

Characteristics	Symptom persistent group (n=129)	Non-persistent group (n=41)	Total (n=170)	p value
Difficulty of seeing even if wearing glasses				0.027
No-no difficulty	90 (69.8)	37 (90.2)	127 (74.7)	
Yes-some difficulty	35 (27.1)	3 (7.3)	38 (22.4)	
Yes-a lot of difficulty	4 (3.1)	1 (2.4)	5 (2.9)	
Cannot do at all	0 (0.0)	0 (0.0)	0 (0.0)	
Difficulty in hearing, even if using a hearing aid				0.178
No-no difficulty	113 (87.6)	40 (97.6)	153 (90.0)	
Yes-some difficulty	15 (11.6)	1 (2.4)	16 (9.4)	
Yes-a lot of difficulty	1 (0.8)	0 (0.0)	1 (0.6)	
Cannot do at all	0 (0.0)	0 (0.0)	0 (0.0)	
Difficulty in walking or climbing steps				0.031
No-no difficulty	97 (75.2)	38 (92.7)	135 (79.4)	
Yes-some difficulty	30 (23.3)	2 (4.9)	32 (18.8)	
Yes-a lot of difficulty	2 (1.6)	1 (2.4)	3 (1.8)	
Cannot do at all	0 (0.0)	0 (0.0)	0 (0.0)	
Difficulty in remembering or concentrating				<0.001
No-no difficulty	65 (50.4)	35 (85.4)	100 (58.8)	
Yes-some difficulty	60 (46.5)	6 (14.6)	66 (38.8)	
Yes-a lot of difficulty	4 (3.1)	0 (0.0)	4 (2.4)	
Cannot do at all	0 (0.0)	0 (0.0)	0 (0.0)	
Difficulty (with self-care such as) washing all over or dressing				>0.999
No-no difficulty	122 (94.6)	39 (95.1)	161 (94.7)	
Yes-some difficulty	7 (5.4)	2 (4.9)	9 (5.3)	
Yes-a lot of difficulty	0 (0.0)	0 (0.0)	0 (0.0)	
Cannot do at all	0 (0.0)	0 (0.0)	0 (0.0)	
Using usual (customary) language, having difficulty in communicating, for example understanding or being understood				0.148
No-no difficulty	114 (88.4)	40 (97.6)	154 (90.6)	
Yes-some difficulty	15 (11.6)	1 (2.4)	16 (9.4)	
Yes-a lot of difficulty	0 (0.0)	0 (0.0)	0 (0.0)	
Cannot do at all	0 (0.0)	0 (0.0)	0 (0.0)	

COVID-19, coronavirus disease.

Data are presented as n (%).