


Supplementary Table 1. Comparison of baseline characteristics for patients with available vs. missing outcomes

Characteristic	mRS available	mRS missing	<i>P</i>	Recurrent events available	Recurrent events missing	<i>P</i>
Number	2,250	400		2,259	391	
Age	72 (14)	71 (14)	0.21	72 (14)	71 (14)	0.31
NIHSS	8 (3–16)	6 (2–13)	<0.01	8 (3–15)	8 (2–14)	0.16
GCS	14 (12–15)	15 (14–15)	<0.01	14 (12–15)	15 (13–15)	<0.01
Systolic blood pressure	165 (144–186)	166 (145–186)	0.78	165 (143–186)	167 (147–189)	0.10
Diastolic blood pressure	90 (78–101)	89 (78–100)	0.55	90 (78–100)	90 (80–104)	0.11
Onset-to-admission time in hours	3.7 (1.4–13.1)	3.8 (1.3–16.9)	0.71	3.9 (1.4–13.5)	3.3 (1.3–12.0)	0.25
Female sex	1,049 (46.7)	178 (45.1)	0.54	1,044 (46.3)	183 (47.3)	0.73
Antiplatelet therapy	617 (28.6)	112 (28.9)	0.91	615 (28.4)	114 (29.9)	0.54
Anticoagulants	508 (23.5)	60 (15.5)	<0.01	516 (23.8)	52 (13.6)	<0.01
Hypertension	1,754 (80.2)	299 (78.1)	0.35	1,745 (79.6)	308 (81.3)	0.46
Diabetes	338 (15.5)	46 (12.0)	0.08	343 (15.7)	41 (10.8)	0.02
Hyperlipidemia	942 (43.4)	154 (41.5)	0.50	943 (43.4)	153 (41.6)	0.52
Smoking	264 (12.1)	42 (11.0)	0.54	261 (12.0)	45 (12.0)	0.99
Atrial fibrillation	417 (19.1)	50 (13.1)	<0.01	423 (19.3)	44 (11.7)	<0.01
History of ischemic stroke	260 (11.9)	30 (7.8)	0.02	260 (11.9)	30 (7.9)	0.03
History of ICH	241 (11.0)	39 (10.2)	0.61	248 (11.3)	32 (8.4)	0.10
Treatment at stroke center	1,799 (80.0)	253 (63.2)	<0.01	1,791 (79.3)	261 (66.8)	<0.01
Discharged to			<0.01			<0.01
Home	371 (21.3)	89 (24.2)		381 (21.8)	79 (21.7)	
Nursing home	118 (6.8)	25 (6.8)		117 (6.7)	26 (7.1)	
Other acute hospital	291 (16.7)	89 (24.2)		284 (16.2)	96 (26.4)	
Rehabilitation	964 (55.3)	165 (44.8)		966 (55.3)	163 (44.8)	

Values are presented as number (%) or median (interquartile range). NIHSS, National Institutes of Health Stroke Scale; GCS, Glasgow coma scale; ICH, intracerebral hemorrhage.



Supplementary Figure 1. Study flowchart. mRS, modified Rankin Scale.