
Supplementary information

A multisystem, cardio-renal investigation of post-COVID-19 illness

In the format provided by the
authors and unedited

1

Supplement

2 **Title: Post-COVID-19 illness trajectory: a multisystem investigation.**

3 **Authors: CISCO-19 investigators.**

4 **Reference: NMED-A116731**

5 **The CISCO-19 Investigators**

6 Chief Investigator: Colin Berry BSc/MBChB/PhD

7 **Grant applicants**

8 Kenneth Mangion MD/PhD, Catherine Bagot MBBS/MD, Colin Church MBChB/PhD,
9 Christian Delles MD, Antonia Ho MBChB/PhD, David J. Lowe MBChB/MD, Alex
10 McConnachie PhD, Giles Roditi MBChB, Rhian M. Touyz MD/PhD, Naveed Sattar
11 MBChB/PhD, Ryan Wereski MBChB, Sylvia Wright MBChB, Colin Berry BSc/MBChB/PhD.

12 **Site investigators**

13 *Queen Elizabeth University Hospital*

14 Kenneth Mangion MBChB/PhD, Pauline Hall Barrientos PhD, Kevin G. Blyth MBChB/PhD,
15 Michael Briscoe MBChB, Colin Church MBChB/PhD, Christian Delles MBChB/PhD, Stephen
16 Dobbin MBChB, Keith Gillis MBChB/PhD, Antonia Ho MBChB/PhD, Anna Kamdar
17 BMedSci, David J. Lowe MBChB/MD, Kaitlin J. Mayne MBChB, Patrick B. Mark
18 MBChB/PhD, Christopher McGinley MBChB, Connor McKee BMedSci/ MBChB, Andrew
19 Morrow MBChB, Oliver Peck MBChB, Alastair J Rankin BSc/MBChB, Giles Roditi MBChB,
20 Claire Rooney MBChB/PhD, Sarah A. Spiers PhD, David Stobo MBChB, Ryan Wereski
21 MBChB, Sylvia Wright MBChB/PhD, Colin Berry BSc/MBChB/PhD.

22 *Royal Alexandra Hospital*

23 Lynn Abel BN, Douglas Grieve MBChB/PhD.

24 *Glasgow Royal Infirmary*

25 Catherine Bagot MBBS/MD, Hannah Bayes MBChB/PhD, Jaclyn Carberry

- 26 MBChB/BMedSci, Daniel Doherty MBChB, Ian Fergus BSc, Vivienne B. Gibson PhD, Fraser
27 Goldie MBChB, Laura Knox MBChB, Giles Roditi MBChB, Katherine Scott MBChB, David
28 Stobo MBChB, Varun Sharma MBChB, Robert Sykes MBChB/BMedSci.

29 **Glasgow Clinical Research Facility and Study Management Group**

- 30 Ammani Brown BN, Andrew Dougherty BN, Kirsty Fallon BN, Lesley Gilmour BN.
31 Chloe Cowan BN, Lynsey Gillespie PhD, Sharon Kean, Jurgen Van-Melckebeke BSc, Kim
32 Moran-Jones PhD, Debra Stuart PhD, Maureen Travers PhD.

33 **Glasgow Clinical Research Imaging Facility, Institute of Clinical Excellence, Queen Elizabeth University Hospital**

- 35 Tracey Hopkins DCR/BSc(Hons)/PGd, Laura Dymock BSc(Hons), Evonne McLennan
36 BSc(Hons)/PGc, Rosemary Woodward BSc/PGc, Fiona Savage BSc(Hons)/PGc, Nicola
37 Tynan BSc(Hons)/PGc.

38 **Radiology, NHS Greater Glasgow and Clyde Health Board**

- 39 Sau Lee Chang MBChB, Mhairi Dupre MBChB, Lindsey Norton MBChB, Liam Peng
40 MBChB, Giles Roditi MBChB, David Stobo MBChB.

41 **Laboratory Medicine and Biorepository, NHS Greater Glasgow and Clyde Health Board**

- 42 Clare Orange BSc, Rory Gunson PhD.

43 **Medical Physics, NHS Greater Glasgow and Clyde**

- 44 Sarah Allwood-Spiers PhD, George Bruce BSc, Rosario Gonzalez-Lopez PhD, Pauline Hall

45 Barrientos PhD, Rebecca Stace BSc.

46 **University of Glasgow**

47 *Institute of Cardiovascular and Medical Sciences*

48 Colin Berry MBChB/PhD, Elaine Butler BSc, Christian Delles MBChB/PhD, Jennifer S Lees
49 MBChB/PhD, Kenneth Mangion MD/PhD, Patrick Mark MBChB/PhD, Andrew Morrow
50 MBChB, Naveed Sattar MBChB/PhD, Robert Sykes MBChB/BMedSci, Rhian M. Touyz
51 MD/PhD, Paul Welsh PhD.

52 *Institute of Infection, Immunity and Inflammation*

53 *MRC Centre for Virology Research*

54 Antonia Ho MBChB/PhD, Massimo Palmarini PhD.

55 *Institute of Health and Wellbeing*

56 Robertson Centre for Biostatistics

57 John G.F. Cleland MBChB/PhD, Sharon Kean, Bernard Kelly BSc, Alex McConnachie PhD,
58 Alasdair McIntosh PhD, Dionne Russell BSc, Sarah Weeden PhD.

59 *Electrocardiology Core Laboratory*

60 Peter W. Macfarlane DSc, Louise Inglis BSc(Hons), Jean Watt, Kathryn McLaren, Shahid Latif
61 MAppSci (now deceased).

62 *Departments of Mathematics and Statistics*

63 Nick Hill PhD, Dirk Husmeier PhD, Xiaoyu Luo PhD.

64 **National Institutes of Health**

65 Peter Kellman PhD, Hui Xue PhD.

66 **Heartflow**

67 Amy Collinsworth MS, Sarah Mullen MBT, Campbell Rogers MD.

68 **Clinical Event Committee**

69 Heerajnarain Bulluck MBChB/PhD, David Carrick MBChB/PhD, David Corcoran
70 MBChB/PhD, Iain Findlay MBChB/MD, Ninian N. Lang, MBChB/PhD, Vera Lennie
71 MBChB, Ross McGeoch MD, Sabrina Nordin MBChB/MD, Alexander Payne MBChB, Keith
72 E. Robertson MBChB/PhD, Nicola Ryan MBChB, Gruschen Veldtman MBChB/PhD, Robin
73 P. Weir MBChB/PhD, Stuart Watkins MBChB/MD.

74 Coordinators – Andrew Morrow MBChB, Robert Sykes MBChB/BMedSci.

75 **Steering Committee**

76 College of Medical, Veterinary and Life Sciences, University of Glasgow - Neil Basu
77 MD/PhD, Iain McInnes MBChB/PhD, Stefan Siebert MD/PhD.

Table S1. Expanded - Clinical characteristics of the population by control vs. COVID-19 and likelihood of adjudicated myocarditis post-COVID-19.

		COVID-19		Control	p-value ^(a)	p-value ^(b)	Adjudicated Myocarditis Diagnosis				p-value
		Without Primary Outcome	With Primary Outcome				Not present	Unlikely	Probable	Very likely	
		n = 104	n = 159				n = 17	n = 56	n = 65	n = 21	
Typical of COVID-19	72 (69.9%)	109 (74.7%)	-	0.848	-	12 (75.0%)	40 (78.4%)	41 (68.3%)	16 (84.2%)	0.024	
Atypical of COVID-19	10 (9.7%)	11 (7.5%)	-				2 (12.5%)	3 (5.9%)	4 (6.7%)	2 (10.5%)	
Unlikely	3 (2.9%)	4 (2.7%)	-				2 (12.5%)	0 (0.0%)	1 (1.7%)	1 (5.3%)	
Normal	18 (17.5%)	22 (15.1%)	-				0 (0.0%)	8 (15.7%)	14 (23.3%)	0 (0.0%)	
<i>COVID-19 treatment</i>											
Oxygen	75 (72.1%)	109 (68.6%)	-	0.584	-	14 (82.4%)	39 (69.6%)	41 (63.1%)	15 (71.4%)	0.509	
Non-invasive respiratory support	18 (17.3%)	31 (19.5%)	-	0.747	-	5 (29.4%)	9 (16.1%)	11 (16.9%)	6 (28.6%)	0.386	
Invasive ventilation		9 (8.7%)	14 (8.8%)	-	1.000	-	2 (11.8%)	1 (1.8%)	5 (7.7%)	6 (28.6%)	0.004
IV inotrope	4 (3.8%)	7 (4.4%)	-	1.000	-	1 (5.9%)	2 (3.6%)	1 (1.5%)	3 (14.3%)	0.092	
Antiviral	24 (23.1%)	42 (26.4%)	-	0.564	-	9 (52.9%)	15 (26.8%)	14 (21.5%)	4 (19.0%)	0.075	
Steroid	45 (43.3%)	89 (56.0%)	-	0.058	-	12 (70.6%)	31 (55.4%)	36 (55.4%)	10 (47.6%)	0.557	
Intensive Care Unit	9 (8.7%)	24 (15.1%)	-	0.133	-	5 (29.4%)	5 (8.9%)	8 (12.3%)	6 (28.6%)	0.048	
<i>Cardiovascular History</i>											
Hypertension	30 (28.8%)	56 (35.2%)	10 (34.5%)	0.347	1.000	8 (47.1%)	21 (37.5%)	18 (27.7%)	9 (42.9%)	0.326	
Renal impairment	11 (10.6%)	7 (4.4%)	0 (0.0%)	0.078	0.598	1 (5.9%)	1 (1.8%)	4 (6.2%)	1 (4.8%)	0.562	
Diabetes mellitus	25 (24.0%)	35 (22.0%)	2 (6.9%)	0.764	0.075	2 (11.8%)	17 (30.4%)	13 (20.0%)	3 (14.3%)	0.294	
Hypercholesterolaemia	47 (45.2%)	76 (47.8%)	13 (44.8%)	0.706	0.841	10 (58.8%)	30 (53.6%)	28 (43.1%)	8 (38.1%)	0.399	
Smoking	Never	62 (59.6%)	106 (66.7%)	18 (62.1%)	0.458	0.737	12 (70.6%)	38 (67.9%)	44 (67.7%)	12 (57.1%)	0.530
	Former	36 (34.6%)	44 (27.7%)	9 (31.0%)			4 (23.5%)	17 (30.4%)	17 (26.2%)	6 (28.6%)	
	Current	6 (5.8%)	9 (5.7%)	2 (6.9%)			1 (5.9%)	1 (1.8%)	4 (6.2%)	3 (14.3%)	
Percutaneous coronary intervention		1 (1.0%)	10 (6.3%)	0 (0.0%)	0.054	0.365	3 (17.6%)	2 (3.6%)	4 (6.2%)	1 (4.8%)	0.215
Coronary artery bypass graft surgery		2 (1.9%)	2 (1.3%)	0 (0.0%)	0.649	1.000	0 (0.0%)	1 (1.8%)	1 (1.5%)	0 (0.0%)	1.000
CCS Angina Class	No Angina	96 (92.3%)	154 (96.9%)	29 (100.0%)	0.144	1.000	16 (94.1%)	55 (98.2%)	64 (98.5%)	19 (90.5%)	0.180
	Angina Class I-IV	8 (7.7%)	5 (3.1%)	0 (0.0%)			1 (5.9%)	1 (1.8%)	1 (1.5%)	2 (9.5%)	
Heart failure		1 (1.0%)	6 (3.8%)	0 (0.0%)	0.250	0.593	0 (0.0%)	2 (3.6%)	3 (4.6%)	1 (4.8%)	1.000
Myocardial infarction		7 (6.7%)	17 (10.7%)	0 (0.0%)	0.382	0.079	3 (17.6%)	6 (10.7%)	6 (9.2%)	2 (9.5%)	0.758
Stroke or transient ischemic attack		2 (1.9%)	5 (3.1%)	3 (10.3%)	0.707	0.108	1 (5.9%)	1 (1.8%)	3 (4.6%)	0 (0.0%)	0.625
Peripheral vascular disease		2 (1.9%)	1 (0.6%)	0 (0.0%)	0.564	1.000	1 (5.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0.107
Cardiovascular disease and/or treatment		52 (50.0%)	74 (46.5%)	14 (48.3%)	0.615	1.000	8 (47.1%)	29 (51.8%)	26 (40.0%)	11 (52.4%)	0.560
<i>Risk Scores</i>											
ISARIC-4c Mortality Score	Mean (SD)	6.76 (3.53)	6.36 (3.33)	4.38 (3.30)	0.368	0.003	7.00 (3.26)	6.71 (3.42)	5.88 (3.06)	6.38 (3.94)	0.426
ISARIC-4c Mortality Risk, in %	Mean (SD)	13.5 (11.5)	12.1 (10.6)	6.9 (8.4)	0.368	0.003	14.0 (10.7)	13.2 (11.4)	10.4 (9.4)	12.8 (11.7)	0.426
QRisk3 10y CVD Risk, in %	Mean (SD)	16.0 (14.0)	13.5 (11.1)	13.1 (10.0)	0.401	0.984	12.5 (7.9)	15.5 (12.8)	12.0 (9.7)	14.3 (13.1)	0.724

		COVID-19		Control	p-value ^(a)	p-value ^(b)	Adjudicated Myocarditis Diagnosis				p-value
		Without Primary Outcome	With Primary Outcome				Not present	Unlikely	Probable	Very likely	
		n = 104	n = 159	n = 29			n = 17	n = 56	n = 65	n = 21	
Charlson Comorbidity Index	Mean (SD)	2.10 (2.21)	1.89 (1.82)	1.45 (1.21)	0.747	0.412	1.71 (1.93)	2.07 (2.01)	1.86 (1.79)	1.62 (1.24)	0.793
<i>Pre-existing maintenance medication</i>											
Aspirin		8 (7.7%)	12 (7.5%)	0 (0.0%)	1.000	0.218	3 (17.6%)	4 (7.1%)	4 (6.2%)	1 (4.8%)	0.419
Statin		25 (24.0%)	46 (28.9%)	10 (34.5%)	0.398	0.659	7 (41.2%)	20 (35.7%)	13 (20.0%)	6 (28.6%)	0.159
Beta-blocker		15 (14.4%)	20 (12.6%)	2 (6.9%)	0.712	0.537	3 (17.6%)	7 (12.5%)	5 (7.7%)	5 (23.8%)	0.208
ACE inhibitor		20 (19.2%)	35 (22.0%)	4 (13.8%)	0.644	0.456	6 (35.3%)	11 (19.6%)	13 (20.0%)	5 (23.8%)	0.542
ARB		9 (8.7%)	10 (6.3%)	2 (6.9%)	0.476	1.000	0 (0.0%)	6 (10.7%)	3 (4.6%)	1 (4.8%)	0.447
Oral anticoagulation		6 (5.8%)	8 (5.0%)	1 (3.4%)	0.786	1.000	1 (5.9%)	3 (5.4%)	3 (4.6%)	1 (4.8%)	1.000
<i>Blood results, index admission</i>											
Haemoglobin, g/l	Mean (SD)	141 (17)	141 (16)	143 (12)	0.970	0.655	142 (15)	140 (17)	140 (15)	143 (16)	0.624
Platelet count, 10⁹/l	Mean (SD)	238 (84)	237 (94)	259 (58)	0.435	0.006	264 (137)	217 (75)	244 (93)	248 (95)	0.344
White cell count, 10 ⁹ /l	Mean (SD)	7.56 (3.34)	7.29 (5.52)	7.51 (3.11)	0.055	0.172	7.32 (4.82)	6.68 (2.65)	7.73 (7.76)	7.50 (2.98)	0.676
Lymphocyte count, 10⁹/l	Mean (SD)	1.21 (0.67)	1.53 (4.66)	1.88 (0.64)	0.823	<0.001	1.02 (0.48)	1.08 (0.52)	2.11 (7.26)	1.35 (0.65)	0.215
Peak D-Dimer, ng/ml	Mean (SD)	1511 (4463)	1740 (5493)	311 (303)	0.722	0.026	2022 (4159)	916 (2132)	1754 (6648)	3127 (7431)	0.881
Peak HbA1c, mmol/mol	Mean (SD)	47.6 (16.9)	48.0 (18.4)	44.2 (22.1)	0.864	0.020	57.3 (32.3)	50.5 (18.1)	44.8 (13.5)	44.9 (19.2)	0.100
Initial creatinine, mmol/l	Mean (SD)	84.3 (29.6)	81.1 (37.1)	75.9 (24.2)	0.069	0.564	96.9 (55.3)	73.8 (21.9)	78.3 (28.2)	96.6 (62.7)	0.316
Initial eGFR, ml/min/1.73m²	Mean (SD)	83.7 (21.8)	88.5 (23.4)	89.5 (19.5)	0.034	0.900	82.5 (27.7)	92.6 (18.7)	89.4 (23.5)	79.8 (28.4)	0.376
Initial eGFR <60 ml/min/1.73m ²		16 (15.4%)	19 (11.9%)	1 (3.4%)	0.460	0.322	2 (11.8%)	4 (7.1%)	8 (12.3%)	5 (23.8%)	0.256
Peak creatinine, µmol/l	Mean (SD)	92 (40)	103 (95)	90 (46)	0.123	0.752	99 (54)	89 (57)	96 (79)	168 (187)	0.506
Minimum eGFR, ml/min/1.73m ²	Mean (SD)	79.2 (22.8)	81.6 (27.4)	77.8 (29.0)	0.124	0.799	79.8 (27.1)	84.8 (23.4)	83.5 (26.5)	68.7 (37.0)	0.454
Minimum eGFR <60 ml/min/1.73m ²		23 (22.3%)	28 (17.6%)	1 (20.0%)	0.425	1.000	2 (11.8%)	7 (12.5%)	12 (18.5%)	7 (33.3%)	0.200
Peak ferritin, µg/l	Median (IQR)	348 (107, 919)	360 (180, 864)	118 (69, 166)	0.333	<0.001	454 (184, 835)	359 (212, 1082)	332 (154, 692)	562 (198, 1860)	0.441
Initial high sensitivity troponin I, ng/l	Median (IQR)	4 (2, 7)	4 (3, 10)	4 (4, 4)	0.003	0.519	5 (4, 8)	4 (3, 8)	4 (3, 7)	16 (4, 51)	0.146
<i>Initial high sensitivity troponin I, males</i>											
<1.2 pg/ml		7 (11.5%)	0 (0.0%)	0 (0.0%)			0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
1.2 - 5.0 pg/ml		33 (54.1%)	44 (55.7%)	15 (88.2%)			6 (54.5%)	16 (57.1%)	18 (58.1%)	4 (44.4%)	
>5 - 34 pg/ml		16 (26.2%)	25 (31.6%)	2 (11.8%)	0.017	0.052	3 (27.3%)	9 (32.1%)	8 (25.8%)	5 (55.6%)	0.745
>34 pg/ml		5 (8.2%)	10 (12.7%)	0 (0.0%)			2 (18.2%)	3 (10.7%)	5 (16.1%)	0 (0.0%)	
<i>Initial high sensitivity troponin I, females</i>											
<1.2 pg/ml		6 (15.8%)	5 (8.5%)	0 (0.0%)			1 (25.0%)	2 (10.0%)	1 (3.8%)	1 (11.1%)	
1.2 - 5.0 pg/ml		23 (60.5%)	37 (62.7%)	7 (77.8%)			1 (25.0%)	14 (70.0%)	21 (80.8%)	1 (11.1%)	
>5 - 16 pg/ml		7 (18.4%)	8 (13.6%)	2 (22.2%)		0.320	2 (50.0%)	3 (15.0%)	3 (11.5%)	0 (0.0%)	
>16 pg/ml		2 (5.3%)	9 (15.3%)	0 (0.0%)			0 (0.0%)	1 (5.0%)	1 (3.8%)	7 (77.8%)	<0.001

		COVID-19		Control	p-value ^(a)	p-value ^(b)	Adjudicated Myocarditis Diagnosis				p-value
		Without Primary Outcome	With Primary Outcome				Not present	Unlikely	Probable	Very likely	
		n = 104	n = 159				n = 17	n = 56	n = 65	n = 21	
Initial high sensitivity troponin I	<1.2 pg/ml	13 (13.1%)	5 (3.6%)	0 (0.0%)	0.031	0.061	1 (6.7%)	2 (4.2%)	1 (1.8%)	1 (5.6%)	0.041
	1.2 - 5.0 pg/ml	56 (56.6%)	81 (58.7%)	22 (84.6%)			7 (46.7%)	30 (62.5%)	39 (68.4%)	5 (27.8%)	
	>5 pg/ml - 99 th percentile	23 (23.2%)	33 (23.9%)	4 (15.4%)			5 (33.3%)	12 (25.0%)	11 (19.3%)	5 (27.8%)	
	>99 th percentile	7 (7.1%)	19 (13.8%)	0 (0.0%)			2 (13.3%)	4 (8.3%)	6 (10.5%)	7 (38.9%)	
Peak high sensitivity troponin I, ng/l	Median (IQR)	4.0 (2.0, 10.5)	4.0 (3.0, 12.8)	4.0 (4.0, 4.0)	0.009	0.358	6.0 (4.0, 10.0)	4.0 (3.0, 10.0)	4.0 (3.0, 9.0)	30.0 (3.5, 83.8)	0.158
Peak high sensitivity troponin I, males	<1.2 pg/ml	6 (9.8%)	0 (0.0%)	0 (0.0%)	0.020	0.044	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0.906
	1.2 - 5.0 pg/ml	29 (47.5%)	43 (54.4%)	14 (87.5%)			6 (54.5%)	15 (53.6%)	18 (58.1%)	4 (44.4%)	
	>5 - 34 pg/ml	19 (31.1%)	21 (26.6%)	2 (12.5%)			3 (27.3%)	9 (32.1%)	6 (19.4%)	3 (33.3%)	
	>34 pg/ml	7 (11.5%)	15 (19.0%)	0 (0.0%)			2 (18.2%)	4 (14.3%)	7 (22.6%)	2 (22.2%)	
Peak high sensitivity troponin I, females	<1.2 pg/ml	6 (15.8%)	5 (8.5%)	0 (0.0%)	0.559	0.362	1 (25.0%)	2 (10.0%)	1 (3.8%)	1 (11.1%)	<0.001
	1.2 - 5.0 pg/ml	21 (55.3%)	35 (59.3%)	7 (77.8%)			0 (0.0%)	14 (70.0%)	20 (76.9%)	1 (11.1%)	
	>5 - 16 pg/ml	6 (15.8%)	7 (11.9%)	2 (22.2%)			2 (50.0%)	3 (15.0%)	2 (7.7%)	0 (0.0%)	
	>16 pg/ml	5 (13.2%)	12 (20.3%)	0 (0.0%)			1 (25.0%)	1 (5.0%)	3 (11.5%)	7 (77.8%)	
Peak high sensitivity troponin I	<1.2 pg/ml	12 (12.1%)	5 (3.6%)	0 (0.0%)	0.034	0.021	1 (6.7%)	2 (4.2%)	1 (1.8%)	1 (5.6%)	0.013
	1.2 - 5.0 pg/ml	50 (50.5%)	78 (56.5%)	21 (84.0%)			6 (40.0%)	29 (60.4%)	38 (66.7%)	5 (27.8%)	
	>5 pg/ml - 99 th percentile	25 (25.3%)	28 (20.3%)	4 (16.0%)			5 (33.3%)	12 (25.0%)	8 (14.0%)	3 (16.7%)	
	>99 th percentile	12 (12.1%)	27 (19.6%)	0 (0.0%)			3 (20.0%)	5 (10.4%)	10 (17.5%)	9 (50.0%)	
Peak fibrinogen, g/l	Mean (SD)	7.13 (2.37)	6.04 (2.54)	3.41 (0.48)	0.376	0.003	6.70 (2.75)	4.69 (2.70)	5.65 (2.22)	7.83 (1.72)	0.240
Peak CRP, mg/l	Median (IQR)	131 (40, 206)	104 (37, 181)	2 (1, 5)	0.385	<0.001	130 (77, 180)	107 (45, 164)	91 (35, 181)	121 (17, 350)	0.584
Peak CRP, mg/l	<5 mg/l	9 (8.7%)	10 (6.3%)	20 (71.4%)	0.529	<0.001	0 (0.0%)	2 (3.6%)	4 (6.2%)	4 (19.0%)	0.274
	≥5, <10 mg/l	2 (1.9%)	6 (3.8%)	4 (14.3%)			0 (0.0%)	3 (5.4%)	3 (4.6%)	0 (0.0%)	
	≥10 mg/l	92 (89.3%)	143 (89.9%)	4 (14.3%)			17 (100.0%)	51 (91.1%)	58 (89.2%)	17 (81.0%)	
Peak sodium, mmol/l	Mean (SD)	140.1 (4.3)	140.3 (4.0)	139.4 (2.5)	0.336	0.157	138.8 (4.2)	139.8 (3.2)	140.8 (3.8)	141.8 (5.6)	0.233
Albumin, g/l	Mean (SD)	34.0 (4.9)	34.2 (5.2)	40.0 (5.3)	0.756	<0.001	32.1 (5.0)	35.0 (4.4)	33.8 (6.0)	34.8 (4.8)	0.273
Initial MCV, fl	Mean (SD)	- (-)	89.6 (6.1)	91.6 (4.1)	-	0.109	89.0 (4.5)	88.4 (7.2)	90.5 (5.7)	90.5 (5.0)	0.551
Initial RDW, %	Mean (SD)	- (-)	13.1 (1.4)	13.1 (0.5)	-	0.164	13.1 (0.8)	13.3 (1.9)	13.1 (1.0)	13.1 (1.2)	0.825
Peak MCV, fl	Mean (SD)	- (-)	92.8 (6.3)	91.9 (4.2)	-	0.221	92.9 (4.8)	90.9 (6.6)	93.6 (5.8)	95.2 (6.7)	0.094
RDW at peak MCV, %	Mean (SD)	- (-)	13.3 (1.3)	13.2 (0.6)	-	0.602	13.3 (0.9)	13.3 (1.5)	13.3 (1.2)	13.8 (1.6)	0.399

769 Table S1: ^(a) COVID-19 patients, without primary outcome vs. with primary outcome. ^(b): COVID-19 patients with primary outcome vs. controls. Ethnicity: Indian (0), Pakistani
770 (0), Bangladeshi (0), Other Asian n=14 (8.8%), Black Caribbean (0), Black African n=2 (1.2%), Chinese n=1 (0.6%), Other (n=1 (0.9%), White, n=139 (87.4%). Missing data
771 in post-COVID-19 patients: D-Dimer, n=62; HbA1c, n=23; ferritin, n=18; troponin I, n=21. Missing data in Control patients: D-Dimer, n=15; HbA1c, n=5; ferritin, n=5;
772 troponin I, n=4. CCS – Canadian Cardiovascular Society; CRP – C-reactive protein; GFR - glomerular filtration rate was estimated using the Chronic Kidney Disease
773 Epidemiology equation⁶; ISARIC-4C - Coronavirus Clinical Characterisation Consortium; MCV – mean cell volume; PCR – polymerase chain reaction; SD – standard
774 deviation; RDW – red cell distribution width; SIMD - Scottish Index of Multiple Deprivation; TIA – transient ischaemic attack; WHO – World Health Organization. In the
775 control group, the Abbott Architect CMIA SARS-CoV-2 IgG assay⁸ was used to confirm absence of prior infection with COVID-19. The primary outcome evaluation (visit 2)
776 was scheduled 28-60 days post-discharge. *Categorical data are summarized as frequency and percentage, and compared between groups using Fisher's Exact tests. Continuous
777 data are summarized as mean and standard deviation, or median and interquartile range (IQR, defined as the upper and lower quartiles), and compared between groups using
778 Kruskal-Wallis tests. All p-values are two-sided. No adjustments were made for multiple comparisons.

779 **Table S2. Expanded - Multisystem phenotyping: serial electrocardiography, biomarkers of inflammation, metabolism, renal function,**
 780 **and hemostasis, and heart, lung, and kidney imaging at 28-60 days post-discharge.**

	COVID-19 n = 159	Controls n = 29	P-value	Not present n = 17 (11%)	Unlikely n = 56 (35%)	Probable n = 65 (41%)	Very likely n = 21 (13%)	Myocarditis P-value*
<i>Electrocardiogram, n (%)</i>								
<i>Admission (n = 150)</i>								
Myopericarditis criteria	31 (20.7)	0 (0)	0.003	3 (17.6)	9 (16.7)	13 (22.4)	6 (28.6)	0.646
Premature atrial contraction	2 (1.3)	0 (0)	1.000	0 (0.0)	0 (0.0)	2 (3.1)	0 (0.0)	0.710
Premature ventricular contraction	3 (1.9)	0 (0)	1.000	1 (5.9)	0 (0.0)	0 (0.0)	2 (9.5)	0.013
Atrial fibrillation or flutter, n = 9 missing	5 (3.3)	0 (0)	1.000	0 (0.0)	2 (3.6)	2 (3.4)	1 (4.8)	1.000
<i>Enrolment (n = 147)</i>								
Myopericarditis criteria	47 (32.0)	0 (0)	<0.001	3 (21.4)	16 (30.2)	20 (32.3)	8 (44.4)	0.586
Premature atrial contraction	7 (4.4)	0 (0)	0.598	1 (5.9)	3 (5.4)	2 (3.1)	1 (4.8)	0.840
Premature ventricular contraction	1 (0.6)	0 (0)	1.000	0 (0.0)	1 (1.8)	0 (0.0)	0 (0.0)	0.591
Atrial fibrillation or flutter	3 (2.0)	0 (0)	1.000	0 (0.0)	1 (1.9)	1 (1.6)	1 (5.9)	0.623
<i>28-60 days post-discharge (n = 143)</i>								
Myopericarditis criteria	33 (23.1)	0 (0)	0.002	2 (14.3)	10 (20.4)	14 (23.3)	7 (35.0)	0.546
Premature atrial contraction	8 (5.0)	0 (0)	0.612	1 (5.9)	3 (5.4)	3 (4.6)	1 (4.8)	1.000
Premature ventricular contraction	2 (1.3)	0 (0)	1.000	1 (5.9)	0 (0.0)	1 (1.5)	0 (0.0)	0.220
Atrial fibrillation or flutter	2 (1.4)	0 (0)	1.000	0 (0.0)	0 (0.0)	1 (1.6)	1 (5.0)	0.535

	COVID-19 n = 159	Controls n = 29	P-value	Not present n = 17 (11%)	Unlikely n = 56 (35%)	Probable n = 65 (41%)	Very likely n = 21 (13%)	Myocarditis P-value*
<i>CT chest 28-60 days post-discharge</i>								
Ground glass opacity and/or consolidation, n (%)	70 (44.6)	1 (4.2)	<0.001	10 (66.7)	26 (46.4)	24 (36.9)	10 (47.6)	0.201
Reticulation and/or architectural distortion, n (%)	47 (29.9)	1 (4.2)	0.006	6 (40.0)	15 (26.8)	18 (27.7)	8 (38.1)	0.600
Atelectasis, n (%)	13 (8.3)	0 (0.0)	0.222	1 (6.7)	7 (12.5)	4 (6.2)	1 (4.8)	0.600
Pulmonary arterial thrombus, n (%)	5 (3.3)	0 (0.0)	1.000	0 (0.0)	2 (3.6)	2 (3.1)	1 (5.3)	0.905
Visual estimate of % of total lung area abnormal, mean (SD)	14.3 (19.0)	0.1 (0.5)	<0.001	19.3 (22.5)	12.7 (17.6)	12.3 (17.5)	21.1 (23.4)	0.286
<20%, n (%)	107 (68.2)	23 (95.8)	0.006	9 (60.0)	38 (67.9)	48 (73.8)	12 (57.1)	0.324
≥20%, n (%)	38 (24.2)	0 (0.0)		3 (20.0)	15 (26.8)	14 (21.5)	6 (28.6)	
≥50%, n (%)	12 (7.6)	1 (4.2)		3 (20.0)	3 (5.4)	3 (4.6)	3 (14.3)	
<i>CT coronary angiogram 28-60 days post-discharge</i>								
Coronary calcium - Agatston score, mean (SD)	144 (502)	95 (297)	0.238	44 (61)	253 (730)	91 (363)	91 (162)	0.071
CADS-RADS score, n (%)								
Level 0	77 (50.0)	13 (52.0)		3 (21.4)	23 (41.1)	40 (63.5)	11 (52.4)	
Level 1	42 (27.3)	7 (32.0)		7 (50.0)	19 (33.9)	11 (17.5)	5 (23.8)	
Level 2	11 (7.1)	2 (8.0)	0.974	0 (0.0)	5 (8.9)	4 (6.3)	2 (9.5)	0.013
Level 3	7 (4.5)	1 (4.0)		2 (14.3)	1 (1.8)	4 (6.3)	0 (0.0)	
Level 4	14 (9.1)	1 (4.0)		1 (7.1)	8 (14.3)	2 (3.2)	3 (14.3)	
Level 5	3 (1.9)	0 (0.0)		1 (7.1)	0 (0.0)	2 (3.2)	0 (0.0)	
Obstructive coronary artery disease, n (%)	21 (13.9)	1 (4.0)	0.325	3 (20.0)	7 (12.7)	7 (11.3)	4 (21.1)	0.555
<i>FFR_{CT}, patient-level (all coronary arteries)</i>								
Median FFR _{CT} , mean (SD)	0.93 (0.03)	0.93 (0.02)	0.109	0.92 (0.03)	0.93 (0.04)	0.93 (0.03)	0.92 (0.04)	0.308
Minimum FFR _{CT} , mean (SD)	0.80 (0.10)	0.84 (0.09)	0.006	0.82 (0.08)	0.79 (0.11)	0.81 (0.09)	0.76 (0.13)	0.541

	COVID-19 n = 159	Controls n = 29	P-value	Not present n = 17 (11%)	Unlikely n = 56 (35%)	Probable n = 65 (41%)	Very likely n = 21 (13%)	Myocarditis P-value*
Minimum FFR _{CT} ≤0.80, n (%)	50 (37.9)	7 (28.0)	0.377	5 (41.7)	16 (34.8)	20 (35.7)	9 (50.0)	0.655
<i>Cardiovascular MRI 28-60 days post-discharge</i>								
LV end-diastolic volume index, mean (SD), mL/m ²	75.9 (17.0)	73.9 (18.7)	0.326	77.2 (17.9)	74.1 (16.6)	77.4 (17.1)	75.2 (17.7)	0.881
LV end-systolic volume index, mean (SD), mL/m²	35.3 (12.8)	30.2 (13.7)	0.012	34.6 (11.1)	33.7 (11.7)	36.3 (14.2)	36.6 (12.4)	0.815
LV ejection fraction, mean (SD), %	54.1 (9.7)	60.4 (9.3)	<0.001	54.8 (9.8)	55.1 (10.1)	54.0 (8.6)	51.3 (11.5)	0.433
LV ejection fraction reduced, males <48%, n (%)	19 (21.3)	2 (12.5)	0.518	2 (15.4)	6 (17.1)	8 (24.2)	3 (33.3)	0.665
LV ejection fraction reduced, females <51%, n (%)	12 (17.6)	0 (0.0)	0.346	1 (25.0)	0 (0.0)	6 (18.8)	5 (41.7)	0.012
LV mass index, mean (SD), g/m²	91.8 (25.6)	119.4 (26.8)	<0.001	100.9 (18.9)	93.2 (21.5)	89.7 (27.8)	87.6 (31.6)	0.170
LV global longitudinal strain, mean (SD), %	-14.2 (3.0)	-14.5 (6.9)	0.031	-13.7 (3.1)	-13.9 (2.9)	-14.6 (3.0)	-14.1 (3.0)	0.375
LV global circumferential strain, mean (SD), %	-17.0 (3.3)	-17.9 (3.0)	0.153	-17.4 (3.6)	-17.0 (3.4)	-17.1 (3.1)	-16.6 (3.4)	0.801
LV global radial strain, mean (SD), %	28.3 (7.8)	30.6 (7.7)	0.164	29.3 (8.4)	28.4 (8.3)	28.3 (7.3)	27.1 (7.9)	0.780
RV end-diastolic volume index, mean (SD), mL/m²	73.3 (17.7)	79.7 (14.1)	0.019	77.8 (18.7)	72.7 (19.7)	72.6 (17.0)	73.3 (13.9)	0.760
RV end-systolic volume index, mean (SD), mL/m ²	35.9 (11.3)	34.4 (10.0)	0.948	34.6 (12.4)	36.6 (11.9)	35.0 (11.5)	38.1 (8.3)	0.487
RV ejection fraction, mean (SD), %	50.9 (10.5)	58.5 (9.3)	<0.001	54.6 (15.9)	49.9 (9.5)	51.9 (9.0)	47.5 (11.4)	0.210
RV global longitudinal strain, mean (SD), %	-17.2 (5.9)	-19.0 (5.8)	0.165	-14.4 (9.5)	-17.5 (5.4)	-18.1 (5.3)	-15.6 (4.3)	0.130
<i>Myocardial tissue characterization</i>								
Increased global T1 (>1233 ms), n (%)	55 (34.8)	5 (19.2)	0.174	2 (12.5)	14 (25.0)	31 (46.2)	9 (42.9)	0.016
Increased global T2 (>44 ms), n (%)	10 (6.3)	0 (0.0)	0.312	0 (0.0)	0 (0.0)	6 (9.2)	4 (19.0)	0.007
T2 ratio (myocardium/ serratus anterior muscle)	1.7 (0.2)	1.6 (0.1)	0.180	1.6 (0.2)	1.6 (0.2)	1.8 (0.2)	1.8 (0.3)	<0.001
Increased global extracellular volume (>27.4%), n (%)	71 (49.7)	5 (20.8)	0.014	1 (7.7)	22 (41.5)	35 (60.3)	13 (68.4)	<0.001
<i>Late gadolinium enhancement</i>								
Myocardial late gadolinium enhancement, n (%)	30 (19.0)	2 (7.7)	0.262	4 (25.0)	7 (12.5)	13 (20.0)	6 (28.6)	0.329

	COVID-19 n = 159	Controls n = 29	P-value	Myocarditis				P-value*
				Not present n = 17 (11%)	Unlikely n = 56 (35%)	Probable n = 65 (41%)	Very likely n = 21 (13%)	
Ischemic distribution, n (%)	8 (5.5)	0 (0.0)	0.658	0 (0.0)	2 (3.9)	5 (8.1)	1 (5.6)	0.769
Non-ischemic distribution, n (%)	24 (16.3)	2 (8.0)	0.606	4 (28.6)	5 (9.8)	8 (12.9)	7 (35.0)	0.033
Pericardial thickening, n (%)	33 (21.2)	0 (0.0)	0.176	1 (5.9)	10 (18.5)	15 (23.4)	7 (33.3)	0.197
Pericardial effusion, n (%)	16 (10.2)	0 (0.0)	0.135	0 (0.0)	5 (9.1)	7 (10.9)	4 (19.0)	0.297
Right atrial area, mean (SD), cm ²	18.7 (4.7)	19.5 (4.2)	0.369	19.2 (3.9)	18.8 (4.9)	18.1 (5.1)	19.9 (3.2)	0.202
Left atrial area, mean (SD), cm ²	20.7 (4.7)	21.8 (4.7)	0.210	22.3 (5.0)	19.9 (4.2)	20.5 (5.1)	21.9 (4.1)	0.204
<i>Myocardial inflammation (Lake Louise criteria), n (%)</i>								
Probable (1/2)	74 (46.8)	1 (3.4)	<0.001	4 (25.0)	49 (87.5)	21 (32.3)	0 (0.0)	<0.001
Definite (2/2)	67 (42.4)	1 (3.4)	<0.001	0 (0)	2 (3.6)	44 (67.7)	21 (100.0)	<0.001
<i>Renal MRI, mean (SD)</i>								
Average volume of right and left kidneys, ml	153 (31)	156 (33)	0.761	158 (37)	154 (25)	150 (33)	153 (38)	0.656
Average cortex T1 of right and left kidneys, ms	1544 (62)	1519 (70)	0.118	1548 (66)	1535 (58)	1541 (63)	1585 (60)	0.110
Average medulla T1 of right and left kidneys, ms	1934 (69)	1953 (59)	0.161	1935 (66)	1924 (65)	1925 (66)	2008 (57)	0.003
Average T1 corticomedullary differentiation of kidneys	0.80 (0.03)	0.78 (0.03)	<0.001	0.80 (0.03)	0.79 (0.03)	0.80 (0.03)	0.79 (0.02)	0.535
<i>Biomarkers at enrolment, central laboratory</i>								
eGFR, median [IQR], ml/min/1.73m²	96 (85, 106)	89 (70, 98)	0.025	95 (88, 103)	94 (84, 107)	97 (83, 105)	98 (94, 104)	0.931
eGFR <60 ml/min/1.73m ² , n (%)	7 (4.7)	1 (4.8)	1.000	1 (6.2)	1 (1.9)	4 (6.6)	1 (5.3)	0.523
C-reactive protein, median [IQR], mg/L	5.5 (1.6, 22.3)	1.5 (0.8, 3.5)	<0.001	6.0 (1.6, 18.2)	5.5 (1.4, 22.8)	4.9 (1.8, 21.6)	14.0 (0.9, 21.5)	0.971
<i>Peak CRP category</i>								
<5.0 mg/l	75 (48.4%)	24 (88.9%)	<0.001	7 (41.2%)	27 (49.1%)	33 (52.4%)	8 (40.0%)	0.591
≥5.0, <10.0 mg/l	20 (12.9%)	2 (7.4%)		4 (23.5%)	8 (14.5%)	7 (11.1%)	1 (5.0%)	
≥10.0 mg/l	60 (38.7%)	1 (3.7%)		6 (35.3%)	20 (36.4%)	23 (36.5%)	11 (55.0%)	

	COVID-19 n = 159	Controls n = 29	P-value	Myocarditis				P-value*
				Not present n = 17 (11%)	Unlikely n = 56 (35%)	Probable n = 65 (41%)	Very likely n = 21 (13%)	
High sensitivity troponin I, median [IQR], ng/L	3 (2, 6)	4 (4, 6)	0.059	4 (3, 5)	3 (2, 7)	3 (2, 5)	4 (3, 7)	0.609
NT-proBNP, median [IQR], pg/mL	114 (57, 262)	58 (38, 99)	0.004	108 (57, 246)	116 (65, 258)	93 (49, 278)	139 (65, 274)	0.546
Ferritin, median [IQR], µg/L	366 (202, 675)	186 (96, 254)	<0.001	428 (143, 576)	398 (281, 658)	313 (172, 683)	379 (187, 637)	0.529
Haptoglobin, median [IQR], g/L	2.1 (1.3, 3.1)	1.5 (1.2, 1.8)	0.001	2.2 (1.8, 3.2)	2.0 (1.5, 2.8)	1.9 (1.2, 3.1)	2.6 (1.6, 3.2)	0.738
Total cholesterol, mean (SD), mmol/L	4.8 (1.4)	4.9 (1.0)	0.530	4.8 (1.4)	4.6 (1.3)	5.0 (1.5)	5.0 (1.1)	0.384
Triglycerides, mean (SD), mmol/L	2.3 (1.3)	1.8 (1.3)	0.016	2.3 (1.3)	2.4 (1.4)	2.2 (1.2)	2.1 (0.8)	0.868
HDL cholesterol, mean (SD), mmol/L	1.1 (0.3)	1.2 (0.3)	0.014	1.0 (0.3)	1.0 (0.3)	1.1 (0.4)	1.1 (0.4)	0.432
Prothrombin time, mean (SD), s	12.1 (3.7)	11.2 (0.8)	0.106	12.1 (2.0)	12.7 (5.5)	11.7 (2.4)	12.0 (1.5)	0.042
D-Dimer, mean (SD), ng/mL	259 (221)	152 (149)	<0.001	290 (195)	263 (178)	246 (168)	260 (192)	0.374
Fibrinogen, mean (SD), g/L	4.1 (1.7)	3.2 (1.1)	0.006	3.9 (1.5)	4.1 (1.7)	4.0 (1.6)	4.5 (1.9)	0.659
Factor VIII, mean (SD), IU/dL	184 (94)	99 (39)	<0.001	208 (88)	183 (97)	181 (99)	173 (73)	0.527
Antithrombin, mean (SD), IU/dL	111 (18)	108 (14)	0.570	108 (18)	112 (18)	112 (17)	109 (20)	0.687
Protein C, mean (SD), IU/dL	123 (30)	116 (21)	0.194	121 (30)	120 (30)	127 (32)	123 (21)	0.574
Protein S, mean (SD), IU/dL	85 (28)	96 (26)	0.116	94 (33)	87 (32)	85 (25)	75 (23)	0.252
VWF:GP1bR, mean (SD), IU/dL	236 (127)	137 (70)	<0.001	257 (176)	241 (118)	224 (123)	246 (122)	0.755
VWF:Ag, mean (SD), IU/dL	243 (145)	204 (251)	0.002	310 (235)	233 (118)	228(135)	261 (140)	0.380
<i>Biomarkers at 28-60 days post-discharge, central laboratory</i>								
eGFR, median [IQR], ml/min/1.73m²	95 (83, 106)	88 (70, 98)	0.047	91 (79, 103)	95 (82, 106)	95 (87, 105)	98 (79, 105)	0.954
eGFR <60 ml/min/1.73m ² , n (%)	7 (4.7)	1 (4.8)	1.000	1 (6.7)	1 (1.9)	4 (6.5)	1 (5.3)	0.514
C-reactive protein, median [IQR], mg/L	1.9 (0.9, 3.5)	1.5 (0.8, 3.5)	0.572	1.7 (1.1, 3.5)	2.0 (1.2, 3.4)	1.8 (0.8, 4.4)	1.9 (0.9, 3.1)	0.996
Peak CRP category								
<5.0 mg/l	124 (81.0%)	24 (88.9%)	0.770	14 (87.5%)	45 (81.8%)	2 (3.6%)	16 (80.0%)	0.562
≥5.0, <10.0 mg/l	16 (10.5%)	2 (7.4%)		1 (6.2%)	8 (14.5%)	6 (9.7%)	1 (5.0%)	

	COVID-19 n = 159	Controls n = 29	P-value	Myocarditis				P-value*
≥10.0 mg/l	13 (8.5%)	1 (3.7%)		1 (6.2%)	2 (3.6%)	7 (11.3%)	3 (15.0%)	
High sensitivity troponin I, median [IQR], ng/L	2 (1, 4)	4 (4, 6)	<0.001	2 (2, 4)	2 (1, 5)	2 (1, 4)	2 (1, 4)	0.941
NT-proBNP, median [IQR], pg/mL	83 (54, 198)	58 (38, 99)	0.022	60 (30, 172)	112 (65, 207)	87 (56, 148)	75 (52, 213)	0.290
Ferritin, median [IQR], ug/L	144 (72, 282)	186 (96, 254)	0.529	145 (86, 299)	158 (94, 296)	129 (57, 206)	157 (99, 319)	0.360
Haptoglobin, median [IQR], g/L	1.3 (0.9, 1.6)	1.5 (1.2, 1.8)	0.031	1.4 (1.1, 1.6)	1.2 (0.8, 1.6)	1.3 (0.8, 1.6)	1.3 (0.9, 1.8)	0.709
D-Dimer, mean (SD), ng/mL	205 (252)	152 (149)	0.146	171 (111)	207 (200)	194 (193)	266 (517)	0.965
Fibrinogen, mean (SD), g/L	3.4(1.4)	3.2 (1.1)	0.439	3.6 (2.4)	3.2 (0.9)	3.5 (1.3)	3.8 (1.7)	0.468
Factor VIII, mean (SD), IU/dL	149 (65)	108 (58)	<0.001	151 (96)	141 (54)	151 (70)	160 (50)	0.606
Protein S, mean (SD), IU/dL	100 (22)	100 (26)	0.260	107 (21)	105 (21)	96 (22)	92 (17)	0.062
VWF:GP1bR, mean (SD), IU/dL	143 (80)	137 (70)	0.912	138 (104)	136 (76)	144 (79)	165 (77)	0.345
VWF:Ag, mean (SD), IU/dL	164 (97)	204 (251)	0.479	151 (79)	155 (85)	157 (88)	224 (144)	0.091
<i>Urine Biomarkers</i>								
Albumin: creatinine ratio at enrolment, mean (SD)	3.2 (8.0)	6.2 (26.9)	0.173	1.6 (3.1)	4.5 (12.0)	2.3 (4.3)	4.1 (6.5)	0.758
Albumin: creatinine ratio at 28-60 days post-discharge, mean (SD)	3.8 (10.9)	6.2 (26.9)	0.257	5.1 (13.4)	5.1 (15.2)	3.4 (5.4)	3.8 (6.4)	0.900

781 Table S2: Missing data in post-COVID-19 patients (admission, enrolment, 28-60 days) and controls – myopericarditis criteria – n=9, n=12, n=16, n=0; Missing data in post-
782 COVID-19 patients at 28-60 days and controls – CT chest atelectasis, reticulation, ground glass – n=2, n=5; pulmonary arterial thrombus – n=8, n=6; CT coronary angiogram
783 28-60 days and controls: Agatston score – n=7, n=4; CAD-RADS score – n=5, n=4; FFR_{CT} – n=27, n=4; Cardiovascular magnetic resonance imaging 28-60 days post-discharge:
784 left ventricular end-diastolic volume index, left ventricular end-systolic volume index, left ventricular ejection fraction, left ventricular strain – n=2, n=3; left ventricular mass
785 – n=2, n=3; right ventricular end-diastolic volume index, right ventricular systolic volume index, n=4, n=3; right ventricular ejection fraction, n=3, n=3; global T1 – n=1, n=3;
786 global T2 – n=1, n=3; global extracellular volume – n=16, n=5; late gadolinium enhancement – n=1, n=3; ischemic distribution – n=14, n=4; non-ischemic distribution – n=12,
787 n=4; mixed distribution – n=14, n=4; pericardial thickening – n=3, n=3; pericardial effusion – n=2, n=3; right and left atrial area – n=1, n=3; myocardial inflammation – n=1,
788 n=0. Missing data in blood biomarkers, post-COVID-19 patients (enrolment and 28-60 days) and controls – eGFR – n=9, n=10, n= 8; C-reactive protein – n=4, n=6, n=2; High
789 sensitivity troponin I – n=6, n=8, n=2; NT-proBNP – n=6, n=10, n=2; Total cholesterol, triglycerides, HDL cholesterol – n=4, n=6, n=2; Fibrinogen – n=5, n=10, n=2; D-Dimer
790 – n=5, n=9, n=2; Factor VIII – n=5, n=9, n=2; Antithrombin – n=5, n=10, n=3; Protein C – n=5, n=10, n=3; Protein S – n=3, n=11, n=3; VWF:GP1bR – n=6, n=9, n=2;
791 VWF:Ag – n=5, n=9, n=2. Abbreviations – aPTT - activated partial thromboplastin time; CAD-RADS - Coronary Artery Disease - Reporting and Data System; ECV -
792 extracellular volume; eGFR (CKD-EPI) – estimated glomerular filtration rate using the Chronic Kidney Disease Epidemiology (CKD-EPI) equation⁶; EF – ejection fraction;
793 EDV – end-diastolic volume; ESV – end-systolic volume; FFR_{CT} – fractional flow reserve computed tomography; HbA1c - hemoglobin A1c; HDL – high density lipoprotein;
794 LV – left ventricle; MESA - Multi-ethnic study of atherosclerosis; NT-proBNP - N-terminal pro B-type natriuretic peptide; PT – prothrombin time; RV - right ventricle; T1 –

795 longitudinal relaxation time; T2 – transverse relaxation time; TCT - thrombin clotting time; vWF:Ag - von Willebrand factor antigen.
796 * Categorical data are summarized as frequency and percentage and compared between groups using Fisher's Exact tests. Continuous data are summarized as mean and standard
797 deviation, or median and interquartile range (IQR, defined as the upper and lower quartiles), and compared between groups using Kruskal-Wallis tests. All p-values are two-
798 sided. No adjustments were made for multiple comparisons. The Kendall's tau rank correlation between Lake Louise criteria, and the final adjudication (4 levels) was 0.75, or
799 with 2 levels, was 0.72, representing moderately strong correlations.

800 **Table S3. SARS-CoV-2 vaccination status.**

	COVID-19		Controls
	Without Primary Outcome	With Primary outcome	
	(n = 104)	(n = 159)	(n = 29)
<i>SARS-CoV-2 vaccination status, n (%)</i>			
First dose prior to admission	1 (1.0)	2 (1.3)	-
Second dose prior to admission	0 (0.0)	0 (0.0)	-
First dose prior to imaging	-	11 (6.9)	23 (79.3)
Second dose prior to imaging	-	0 (0.0)	15 (51.7)
Third dose prior to imaging	-	0 (0.0)	0 (0.0)

801 Table S3 shows numbers and percentages. SARS-CoV-2 vaccines used in the NHS Scotland vaccination
 802 programme are the messenger RNA vaccines, BNT162b2 and mRNA-1273 manufactured by Biotech GmbH
 803 (Mainz, Germany) and Moderna (Cambridge, MA), respectively, and the AZD1222 (ChAdOx1 nCoV-19)
 804 recombinant SARS-CoV-2 vaccine manufactured by AstraZeneca (Cambridge, UK).

Table S4. Primary and secondary outcomes.

Myocarditis						
	All cases n=159	Not present n=17 (11%)	Unlikely n=56 (35%)	Probable n=65 (41%)	Very likely n=21 (13%)	P-value
<i>Cause of myocardial inflammation (endotype)</i>						
<i>SARS-COV-2 Myocarditis, n (%)</i>						
Not present	16 (10.1)	16 (94.1)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001
Unlikely	59 (37.1)	1 (5.9)	56 (100.0)	2 (3.1)	0 (0.0)	
Probable	76 (47.8)	0 (0.0)	0 (0.0)	62 (95.4)	14 (66.7)	
Very Likely	8 (5.0)	0 (0.0)	0 (0.0)	1 (1.5)	7 (33.3)	
<i>Acute Stress Cardiomyopathy, n (%)</i>						
Not present	89 (56.0)	15 (88.2)	37 (66.1)	26 (40.0)	11 (52.4)	<0.001
Unlikely	69 (43.4)	1 (5.9)	19 (33.9)	39 (60.0)	10 (47.6)	
Probable	1 (0.6)	1 (5.9)	0 (0.0)	0 (0.0)	0 (0.0)	
Very Likely	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
<i>Ischemia/ Impaired Perfusion as a Stressor of Inflammation, n (%)</i>						
Not present	75 (47.2)	14 (82.4)	35 (62.5)	21 (32.3)	5 (23.8)	<0.001
Unlikely	73 (45.9)	2 (11.8)	17 (30.4)	39 (60.0)	15 (71.4)	
Probable	9 (5.7)	0 (0.0)	3 (5.4)	5 (7.7)	1 (4.8)	
Very Likely	2 (1.3)	1 (5.9)	1 (1.8)	0 (0.0)	0 (0.0)	
<i>Infective Myopericarditis (non-Covid infection), n (%)</i>						
Not present	42 (26.4)	17 (100.0)	12 (21.4)	11 (16.9)	2 (9.5)	<0.001
Unlikely	117 (73.6)	0 (0.0)	44 (78.6)	54 (83.1)	19 (90.5)	
Probable	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
Very Likely	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	

	Myocarditis					P-value
	All cases n=159	Not present n=17 (11%)	Unlikely n=56 (35%)	Probable n=65 (41%)	Very likely n=21 (13%)	
<i>Drug Induced (Toxic) Myocardial Inflammation, n (%)</i>						
Not present	71 (44.7)	17 (100.0)	29 (51.8)	20 (30.8)	5 (23.8)	<0.001
Unlikely	88 (55.3)	0 (0.0)	27 (48.2)	45 (69.2)	16 (76.2)	
Probable	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
Very Likely	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
<i>Idiopathic Myocardial and/or Pericardial Inflammation, n (%)</i>						
Not present	29 (18.2)	17 (100.0)	6 (10.7)	4 (6.2)	2 (9.5)	<0.001
Unlikely	130 (81.8)	0 (0.0)	50 (89.3)	61 (93.8)	19 (90.5)	
Probable	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
Very Likely	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	

806
807
808
809

Table S4 shows numbers and percentages, with p-values from Fisher's Exact tests. All p-values are two-sided. No adjustments were made for multiple comparisons. The exact p-values are : SARS-COV-2 Myocarditis, p=0.0001; Acute Stress Cardiomyopathy, p=0.0001; Ischemia/ Impaired Perfusion as a Stressor of Inflammation, p=0.0001; Infective Myopericarditis (non-Covid infection), p=0.0001; Drug Induced (Toxic) Myocardial Inflammation, p=0.0001; Idiopathic Myocardial and/or Pericardial Inflammation, p=0.0001.

810 **Table S5. Diagnoses given by all raters in relation to the adjudicated diagnosis.**

	All COVID-19	Adjudicated Myocarditis Diagnosis				p-value
		Not present	Unlikely	Probable	Very likely	
		(n=795)	(n=85)	(n=280)	(n=335)	
Individual rater assessment of the likelihood of myocarditis	Not Present	111 (14.0%)	74 (87.1%)	36 (12.9%)	1 (0.3%)	0 (0.0%)
	Unlikely	223 (28.1%)	9 (10.6%)	171 (61.1%)	41 (12.6%)	2 (1.9%)
	Probable	314 (39.5%)	2 (2.4%)	68 (24.3%)	214 (65.8%)	30 (28.6%)
	Very Likely	147 (18.5%)	0 (0.0%)	5 (1.8%)	69 (21.2%)	73 (69.5%)

811 Table S5 shows numbers and percentages, with p-value from Fisher's Exact tests (two-sided). The denominators at the top of the table are five times the numbers of patients
 812 with each adjudicated diagnosis. E.g. with 21 patients adjudicated as being highly likely to have myocarditis, there were 105 individual assessment of this likelihood for these
 813 patients. The data illustrate a high level of concordance between the raters for myocarditis not present, and good discrimination between probably and very likely.

Table S6. Clinical and Diagnostic criteria for myocarditis, split by adjudicated likelihood of myocarditis diagnosis.

	COVID-19 n=159	Adjudicated Myocarditis Diagnosis				p-value	
		Not present n=17	Unlikely n=56	Probable n=65	Very likely n=21		
<i>Clinical criteria</i>							
Chest pain	38 (23.9%)	7 (41.2%)	10 (17.9%)	13 (20.0%)	8 (38.1%)	0.081	
New symptoms of breathlessness	149 (93.7%)	17 (100.0%)	51 (91.1%)	61 (93.8%)	20 (95.2%)	0.804	
Chronic symptoms of breathlessness	2 (1.3%)	0 (0.0%)	1 (1.8%)	1 (1.5%)	0 (0.0%)	1.000	
Cardiogenic shock	2 (1.3%)	1 (5.9%)	0 (0.0%)	1 (1.5%)	0 (0.0%)	0.220	
Palpitations / Syncope / Arrhythmia	14 (8.8%)	1 (5.9%)	4 (7.1%)	7 (10.8%)	2 (9.5%)	0.905	
Number of clinical criteria	0 1 2 3 4	7 (4.4%) 104 (65.4%) 44 (27.7%) 3 (1.9%) 1 (0.6%)	0 (0.0%) 8 (47.1%) 9 (52.9%) 0 (0.0%) 0 (0.0%)	4 (7.1%) 40 (71.4%) 10 (17.9%) 2 (3.6%) 0 (0.0%)	3 (4.6%) 44 (67.7%) 16 (24.6%) 1 (1.5%) 1 (1.5%)	0 (0.0%) 12 (57.1%) 9 (42.9%) 0 (0.0%) 0 (0.0%)	0.206
<i>Diagnostic criteria</i>							
ECG Myocarditis	30 (18.9%)	3 (17.6%)	9 (16.1%)	12 (18.5%)	6 (28.6%)	0.651	
T2 Criteria	89 (56.0%)	0 (0.0%)	16 (28.6%)	52 (80.0%)	21 (100.0%)	<0.001	
T1 Criteria	114 (71.7%)	0 (0.0%)	36 (64.3%)	57 (87.7%)	21 (100.0%)	<0.001	
Lake Louise Criteria	1 of 2 2 of 2	73 (45.9%) 65 (40.9%)	0 (0.0%) 0 (0.0%)	52 (92.9%) 0 (0.0%)	21 (32.3%) 44 (67.7%)	0 (0.0%) 21 (100.0%)	<0.001
Elevated high sensitivity troponin	77 (48.4%)	6 (35.3%)	26 (46.4%)	31 (47.7%)	14 (66.7%)	0.267	
Global left ventricular dysfunction	32 (20.1%)	3 (17.6%)	6 (10.7%)	14 (21.5%)	9 (42.9%)	0.022	
Regional left ventricular dysfunction	34 (21.4%)	3 (17.6%)	5 (8.9%)	17 (26.2%)	9 (42.9%)	0.006	

Pericardial changes	32 (20.1%)	0 (0.0%)	10 (17.9%)	15 (23.1%)	7 (33.3%)	0.046
New left ventricular dysfunction	29 (18.2%)	2 (11.8%)	5 (8.9%)	14 (21.5%)	8 (38.1%)	0.022
0	10 (6.8%)	10 (58.8%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
1-2	28 (19.0%)	4 (23.5%)	20 (35.7%)	4 (6.7%)	0 (0.0%)	
Number of diagnostic criteria	3-4	62 (42.2%)	3 (17.6%)	32 (57.1%)	24 (40.0%)	3 (21.4%)
	5-6	37 (25.2%)	0 (0.0%)	3 (5.4%)	26 (43.3%)	8 (57.1%)
	7+	10 (6.8%)	0 (0.0%)	1 (1.8%)	6 (10.0%)	3 (21.4%)

815 Table S6 shows numbers and percentages, with p-values from Fisher's Exact tests. All p-values are two-sided. No adjustments were made for multiple comparisons.

816 **Table S7. Serious adverse events occurring during the index admission.**

	COVID-19 population		
	Without Primary Outcome (n = 104)	With Primary outcome (n = 159)	Fisher Test p-value
<i>Index admission, n (%)</i>			
Death	1 (1.0)	0 (0.0)	0.40
Myocardial infarction	1 (1.0)	7 (4.4)	0.16
Stroke	1 (1.0)	0 (0.0)	0.40
Heart failure	2 (1.9)	1 (0.6)	0.57
Pulmonary embolism	5 (4.8)	5 (3.1)	0.53
Deep vein thrombosis	0 (0.0)	1 (0.6)	1.00
Atrial fibrillation	6 (5.8)	5 (3.1)	0.36
Ventricular arrhythmia	0 (0.0)	0 (0.0)	-
Non-cardiovascular serious adverse event	0 (0.0)	0 (0.0)	-

817 Table S7 shows numbers and percentages, with p-values from Fisher's Exact tests. All p-values are two-sided. No adjustments were made for multiple comparisons.

Table S8. Clinical outcomes in the primary analysis population.

	With Primary Outcome n = 159	Control n = 27*	p-value ^(a)	Myocarditis				p-value ^(b)
				Not present n = 17 (11%)	Unlikely n = 56 (35%)	Probable n = 65 (41%)	Very likely n = 21 (13%)	
<i>Duration of follow-up</i>								
Days to Visit 3 or death, (median IQR)	419 (369, 451)	218 (208, 236)	<0.001	429 (416, 545)	414 (369, 437)	417 (369, 444)	454 (365, 546)	0.185
<i>Outcomes, n (%)</i>								
Death or Hospitalization (Any Cause)	24 (15.1)	2 (7.4)	0.356	3 (17.6)	10 (17.9)	9 (13.8)	2 (9.5)	0.845
Death (Any Cause)	2 (1.3)	0 (0.0)	0.663	0 (0.0)	0 (0.0)	2 (3.1)	0 (0.0)	0.413
Cardiovascular Death	1 (0.6)	0 (0.0)	0.888	0 (0.0)	0 (0.0)	1 (1.5)	0 (0.0)	0.701
Renal Death	0 (0.0)	0 (0.0)	-	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-
Respiratory Death	0 (0.0)	0 (0.0)	-	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-
Hospitalization (Any Cause)	24 (15.1)	2 (7.4)	0.356	3 (17.6)	10 (17.9)	9 (13.8)	2 (9.5)	0.845
Cardiovascular Hospitalization	7 (4.4)	2 (7.4)	0.340	1 (5.9)	5 (8.9)	1 (1.5)	0 (0.0)	0.177
Renal Hospitalization	2 (1.3)	0 (0.0)	0.680	1 (5.9)	1 (1.8)	0 (0.0)	0 (0.0)	0.526
Respiratory Hospitalization	8 (5.0)	0 (0.0)	0.242	2 (11.8)	2 (3.6)	2 (3.1)	2 (9.5)	0.335
<i>Cardiovascular outcomes, n (%)</i>								
Myocardial infarction	1 (0.6)	0 (0.0)	0.679	1 (5.9)	0 (0.0)	0 (0.0)	0 (0.0)	0.040
Percutaneous Coronary Intervention	3 (1.9)	0 (0.0)	0.481	1 (5.9)	1 (1.8)	0 (0.0)	1 (4.8)	0.300
Coronary Artery Bypass Grafting	0 (0.0)	0 (0.0)	-	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-
Cerebrovascular accident	0 (0.0)	1 (3.7)	0.016	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-
Heart Failure	2 (1.3)	1 (3.7)	0.232	0 (0.0)	1 (1.8)	1 (1.5)	0 (0.0)	0.886
Deep vein thrombosis	0 (0.0)	0 (0.0)	-	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-
New atrial fibrillation	3 (1.9)	1 (3.7)	0.204	0 (0.0)	1 (1.8)	2 (3.1)	0 (0.0)	0.739

Ventricular tachycardia or fibrillation	0 (0.0)	0 (0.0)	-	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-
<i>Respiratory outcomes, n (%)</i>								
Pulmonary fibrosis	9 (5.7)	0 (0.0)	0.336	0 (0.0)	0 (0.0)	4 (6.2)	5 (23.8)	<0.001
New diagnosis asthma	1 (0.6)	0 (0.0)	0.680	1 (5.9)	0 (0.0)	0 (0.0)	0 (0.0)	0.039
Pulmonary embolism	4 (2.5)	0 (0.0)	0.407	0 (0.0)	1 (1.8)	2 (3.1)	1 (4.8)	0.778
Long-term oxygen therapy	2 (1.3)	0 (0.0)	0.559	0 (0.0)	0 (0.0)	1 (1.5)	1 (4.8)	0.391
<i>Secondary care (outpatients)</i>								
Any outpatient care	108 (67.9)	7 (25.9)	0.017	12 (70.6)	35 (62.5)	44 (67.7)	17 (81.0)	0.184
Acute COVID-19 (<28 days)	15 (9.4)	0 (0.0)	0.102	2 (11.8)	3 (5.4)	9 (13.8)	1 (4.8)	0.340
Ongoing COVID-19 (28-84 days)	20 (12.6)	1 (3.7)	0.194	1 (5.9)	5 (8.9)	9 (13.8)	5 (23.8)	0.225
Long COVID-19 (>84 days)	58 (36.5)	1 (3.7)	0.017	9 (52.9)	13 (23.2)	27 (41.5)	9 (42.9)	0.122
Cardiology	22 (13.8)	1 (3.7)	0.320	3 (17.6)	9 (16.1)	6 (9.2)	4 (19.0)	0.650
Respiratory	55 (34.6)	1 (3.7)	0.013	8 (47.1)	12 (21.4)	27 (41.5)	8 (38.1)	0.102
Physiotherapy	22 (13.8)	1 (3.7)	0.232	2 (11.8)	4 (7.1)	12 (18.5)	4 (19.0)	0.264

819 Table S8: For clinical outcomes, data are the number and percentage of participants with at least one event during follow-up. P-values are from log rank tests of the time to first event.
820 All p-values are two-sided. No adjustments were made for multiple comparisons. (a): COVID-19 patients with primary outcome vs. controls. (b): Comparing subgroups defined by
821 likelihood of myocarditis. *Two controls developed COVID-19 after the imaging visit, hence they were excluded from this group.

822

Table S9. Deaths and adjudicated causes of death.

Participant ID	Date of consent	Date of death	Death during index hospitalisation, Y/N	Visit 2, Y/N	Myocarditis adjudicated very likely, Y/N	Acute kidney injury, Y/N	Adjudicated cause of death
1074	21/09/2020	16/12/2020	N	Y	N	N	Urosepsis
1104	08/10/2020	02/11/2020	N	N	N/A	N	Metastatic prostate cancer
1117	20/10/2020	15/08/2021	N	Y	N	N	Diffuse hypoxic brain injury secondary to out of hospital cardiac arrest and lower respiratory tract infection
2025	23/10/2020	16/11/2020	Y	N	N/A	N	COVID-19 pneumonitis

823

Table S9: The reporting period extends from the date of the first patient in (22 May 2020) to the last day of follow-up (13 December 2021). Participant details: Patient 1074 had chronic kidney disease stage 3B and the adjudicated likelihood of myocarditis for this patient was ‘probably’ (median score = 3; adjudicator scores: 2,3,3,3,3). The adjudicated likelihood of myocarditis for patient 1117 was ‘probably’ (median score = 3; adjudicator scores: 3,3,3,4,4).

824

825

Table S10. Medication prescription at follow-up.

	COVID-19 N=159	Control N=27*	p-value ^(a)	Myocarditis				p-value ^(b)
				Not present n = 17 (11%)	Unlikely n = 56 (35%)	Probable n = 65 (41%)	Very likely n = 21 (13%)	
<i>Medication, n (%)</i>								
Aspirin	22 (13.8%)	0 (0.0%)	0.048	4 (23.5)	7 (12.5)	6 (9.2)	5 (23.8)	0.204
P2Y12 inhibitor	8 (5.0%)	3 (11.1%)	0.202	2 (11.8)	3 (5.4)	2 (3.1)	1 (4.8)	0.413
Calcium channel blocker	31 (19.5%)	6 (22.2%)	0.795	6 (35.3)	13 (23.2)	6 (9.2)	6 (28.6)	0.021
Angiotensin converting enzyme inhibitor	37 (23.3%)	5 (18.5%)	0.804	5 (29.4)	13 (23.2)	12 (18.5)	7 (33.3)	0.463
Angiotensin receptor blocker	15 (9.4%)	2 (7.4%)	1.000	1 (5.9)	7 (12.5)	6 (9.2)	1 (4.8)	0.802
Sacubitril/Valsartan	5 (3.1%)	0 (0.0%)	1.000	1 (5.9)	1 (1.8)	2 (3.1)	1 (4.8)	0.625
Statin	76 (47.8%)	15 (55.6%)	0.534	11 (64.7)	32 (57.1)	25 (38.5)	8 (38.1)	0.074
Mineralocorticoid receptor antagonist	2 (1.3%)	0 (0.0%)	1.000	0 (0.0)	0 (0.0)	2 (3.1)	0 (0.0)	0.710
Beta Blocker	33 (20.8%)	3 (11.1%)	0.301	4 (23.5)	9 (16.1)	12 (18.5)	8 (38.1)	0.190
Thiazide	14 (8.8%)	2 (7.4%)	1.000	2 (11.8)	10 (17.9)	1 (1.5)	1 (4.8)	0.008
Warfarin	1 (0.6%)	0 (0.0%)	1.000	0 (0.0)	1 (1.8)	0 (0.0)	0 (0.0)	0.591
Novel oral anticoagulant therapy	12 (7.5%)	3 (11.1%)	0.461	1 (5.9)	4 (7.1)	5 (7.7)	2 (9.5)	0.968
Inhaled Steroid	24 (15.1%)	2 (7.4%)	0.379	2 (11.8)	8 (14.3)	9 (13.8)	5 (23.8)	0.691
Inhaled Bronchodilator	31 (19.5%)	3 (11.1%)	0.422	2 (11.8)	11 (19.6)	11 (16.9)	7 (33.3)	0.349
Inhaled Anti-muscarinics	15 (9.4%)	1 (3.7%)	0.475	0 (0.0)	5 (8.9)	6 (9.2)	4 (19.0)	0.285
Sodium-glucose co-transporter-2 inhibitor	17 (10.7%)	1 (3.7%)	0.479	0 (0.0)	8 (14.3)	7 (10.8)	2 (9.5)	0.456
Glucagon-like peptide-1 agonist	4 (2.5%)	1 (3.7%)	0.548	1 (5.9)	1 (1.8)	1 (1.5)	1 (4.8)	0.388
Insulin	8 (5.0%)	0 (0.0%)	0.605	1 (5.9)	5 (8.9)	1 (1.5)	1 (4.8)	0.222
Other anti-diabetic	24 (15.1%)	1 (3.7%)	0.135	2 (11.8)	12 (21.4)	7 (10.8)	3 (14.3)	0.436
Loop diuretic	5 (3.1%)	1 (3.7%)	1.000	1 (5.9)	1 (1.8)	1 (1.5)	2 (9.5)	0.180

Oral Steroid	1 (0.6%)	1 (3.7%)	0.270	0 (0.0)	0 (0.0)	0 (0.0)	1 (4.8)	0.239
Antidepressant	28 (17.6%)	5 (18.5%)	1.000	2 (11.8)	5 (8.9)	15 (23.1)	6 (28.6)	0.087

827
828
829

Table S10: n (%) of participants who were taking the medication at follow-up. *Two controls developed COVID-19 after the imaging visit, hence they were excluded from this group.
P-values are from Fisher's Exact tests. All p-values are two-sided. No adjustments were made for multiple comparisons. (a): COVID-19 patients with primary outcome vs. controls. (b): Comparing subgroups defined by likelihood of myocarditis.