## **Supporting Information**

Inactivated Cowpea mosaic virus in combination with OX40 agonist primes potent antitumor immunity in bilateral melanoma mouse model

Edward C. Koellhoffer<sup>1</sup>, Chenkai Mao<sup>2</sup>, Veronique Beiss<sup>3</sup>, Lu Wang<sup>4</sup>, Steven N. Fiering<sup>2,5</sup>, Christine E. Boone<sup>\*1</sup>, Nicole F. Steinmetz<sup>1,3,4,6,7,8\*</sup>

<sup>1</sup>Department of Radiology, University of California, San Diego, La Jolla CA 92093, United States; <sup>2</sup>Department of Microbiology and Immunology, Geisel School of Medicine, Dartmouth College, Hanover, NH 03755 United States; <sup>3</sup>Department of NanoEngineering, University of California, San Diego, La Jolla CA 92093, United States; <sup>4</sup>Department of Bioengineering, University of California, San Diego, La Jolla CA 92093, United States; <sup>5</sup>Norris Cotton Cancer Center, Geisel School of Medicine and Dartmouth Hitchcock Medical System, NH 03755 United States; <sup>6</sup>Moores Cancer Center, University of California, San Diego, La Jolla CA 92093, United States; <sup>7</sup>Center for Nano-ImmunoEngineering, University of California, San Diego, La Jolla CA 92093, United States; <sup>8</sup>Institute for Materials Design and Discovery, University of California, San Diego, La Jolla CA 92093, United States

## \* Co-Corresponding authors:

Christine E. Boone, M.D., Ph.D. <u>ceboone@health.ucsd.edu</u> and Nicole F. Steinmetz, Ph.D., <u>nsteinmetz@ucsd.edu</u>



Figure S1. CPMV vs. inCPMV leads to higher induction of IL-6 and IFN- $\beta$  due to additional TLR7 stimulation by packaged ssRNA cargo.

Direct comparison of splenocyte assays established previously<sup>19</sup> set up from C57BL6 and TLR7-/- mice (n=3/strain) respectively **from Figs. 3A and 3D**. **A**) Supernatant was analyzed by IL-6 ELISA and **B**) IFN- $\beta$  ELISA. Data for bar graphs were calculated using unpaired Student's *t*test with *p* <0.05 as \*, *p* <0.01 as \*\*, and *p* <0.001 as \*\*\*.