

Supplemental 1. Bivariable and Multivariable Survival Analysis for Patients with pT4b Locally Advanced Gastric Cancer who Underwent Gastrectomy with Multivisceral Resection

	Bivariable HR (95% CI)*	Multivariable HR (95% CI)*
Age (years)		
18-49	Reference	Reference
50-64	0.94 (0.63-1.39)	1.00 (0.66-1.53)
65-74	0.92 (0.60-1.42)	1.14 (0.72-1.81)
≥ 75	1.48 (0.97-2.26)	1.44 (0.91-2.28)
Female Sex	1.31 (1.01-1.69)	1.29 (0.97-1.71)
Race		
White	Reference	
African-American	0.82 (0.60-1.14)	
Asian-American	0.67 (0.41-1.11)	
Hispanic Ethnicity		
Yes	1.04 (0.68-1.60)	
Charlson-Deyo Score		
0	Reference	Reference
1	1.28 (0.95-1.73)	1.19 (0.86-1.64)
≥ 2	0.91 (0.54-1.51)	0.90 (0.51-1.58)
Gastrectomy Type		
Subtotal	Reference	
Total	1.35 (1.03-1.79)	1.16 (0.86-1.56)
Gastrectomy Unspecified	1.48 (1.00-2.17)	1.20 (0.80-1.81)

Pathologic Nodal Stage		
0	Reference	Reference
1	1.51 (0.98-2.35)	1.54 (0.98-2.42)
2	1.35 (0.88-2.08)	1.38 (0.88-2.16)
3	2.07 (1.43-2.98)	1.97 (1.33-2.91)
Resection Margin Status		
Positive	1.87 (1.44-2.43)	1.49 (1.11-2.00)
Neoadjuvant Chemotherapy		
Yes	0.93 (0.70-1.25)	
Neoadjuvant Radiation		
Yes	0.84 (0.31-2.26)	
Adjuvant Chemotherapy		
Yes	0.53 (0.41-0.69)	0.58 (0.42-0.80)
Adjuvant Radiation Therapy		
Yes	0.60 (0.44-0.82)	0.84 (0.58-1.21)
Treatment at a High-Volume Center		
Yes	0.76 (0.57-0.99)	0.72 (0.54-0.96)

* Hazard ratios and 95% confidence intervals