

Supplementary information

Bitter taste receptor T2R38 is expressed on skin-infiltrating lymphocytes and regulates lymphocyte migration

Moe Sakakibara¹, Hayakazu Sumida^{1*}, Keisuke Yanagida², Sosuke Miyasato³, Motonao Nakamura³,
Shinichi Sato¹

Affiliations of the authors

¹Department of Dermatology, Faculty of Medicine, The University of Tokyo, Bunkyo-Ku, Tokyo, Japan.

²Department of Lipid Signaling, National Center for Global Health and Medicine, Shinjuku-ku, Tokyo, Japan.

³Department of Bioscience, Graduate School of Life Science, Okayama University of Science, Okayama, Japan.

Corresponding author: Hayakazu Sumida, MD, PhD

Department of Dermatology, Faculty of Medicine, The University of Tokyo, 7-3-1, Hongo, Bunkyo-Ku, Tokyo, 113-8655, Japan.

E-mail: sumida-tky@umin.ac.jp

Tel.: +81 3 5800 8661

Fax: +81 3 3814 1503

Supplementary Table 1

Row data for *GAPDH* in PCR assays from skin samples

sample name	Target	CT value
Healthy skin No.1	<i>GAPDH</i>	23.96975708
Healthy skin No.2	<i>GAPDH</i>	22.54921341
Healthy skin No.3	<i>GAPDH</i>	22.74416351
Healthy skin No.4	<i>GAPDH</i>	21.17991638
Healthy skin No.5	<i>GAPDH</i>	21.6879158
AD skin No.1	<i>GAPDH</i>	22.31202698
AD skin No.2	<i>GAPDH</i>	22.04991722
AD skin No.3	<i>GAPDH</i>	23.56932449
AD skin No.4	<i>GAPDH</i>	23.06023407
AD skin No.5	<i>GAPDH</i>	20.09181595
AD skin No.6	<i>GAPDH</i>	21.82665062
AD skin No.7	<i>GAPDH</i>	23.96928406
AD skin No.8	<i>GAPDH</i>	22.99201775
AD skin No.9	<i>GAPDH</i>	22.86565018
AD skin No.10	<i>GAPDH</i>	21.79933929
AD skin No.11	<i>GAPDH</i>	22.20566368

AD, Atopic dermatitis