

Supplementary Table 2. Subgroup Analysis of Patients with Appendicitis

	Pre-SLRW (n=1238)	Post-SLRW (n=886)	<i>p</i> value
Age (yr)	44 (31–55)	46 (31–58)	<0.001
Sex (male)	599 (48.4)	427 (48.2)	0.965
Laboratory test			
Hemoglobin <10 (g/dL)	18 (1.5)	14 (1.6)	0.857
Abnormal WBC ($10^3/\mu\text{L}$)	904 (73.3)	604 (68.9)	0.028
Neutrophil count >80 (%)	548 (44.4)	397 (45.3)	0.696
C-reactive protein >1.0 (mg/dL)	753 (61.6)	564 (64.8)	0.134
Blood urea nitrogen >20 (mg/dL)	72 (5.9)	66 (7.6)	0.119
Creatinine >1.0 (mg/dL)	146 (11.9)	102 (11.7)	0.898
Albumin <3.0 (g/dL)	17 (1.4)	9 (1.2)	0.720
Total bilirubin >2.0 (mg/dL)	91 (7.4)	33 (3.8)	0.001
Vital sign			
SBP <90 (mm Hg)	14 (1.1)	7 (0.8)	0.434
Pulse rate >100 bpm	122 (9.9)	101 (11.4)	0.252
Body temperature >37.8 (°C)	204 (16.5)	153 (17.3)	0.631
Time from ER arrival to surgery (min)	423.5 (330.5–598.5)	435.0 (331.0–636.5)	0.169
Length of hospital stay (day)	4 (4–5)	4 (3–5)	0.316
Mortality	9 (0.7)	3 (0.3)	0.239

SLRW, special law on restriction of working hours in training; ER, emergency room; SBP, systolic blood pressure.

Data are presented as interquartile range or n (%).