

Supplementary Table 3. Subgroup Analysis of Patients with Cholecystitis

	Pre-SLRW (n=65)	Post-SLRW (n=128)	<i>p</i> value
Age (yr)	52.6±12.8	57.7±15.4	0.016
Sex (male)	35 (53.8)	73 (57.0)	0.674
Laboratory test			
Hemoglobin <10 (g/dL)	0 (0.0)	1 (0.8)	0.475
Abnormal WBC (10 ³ /uL)	35 (53.8)	88 (68.8)	0.042
Neutrophil count >80 (%)	24 (36.9)	56 (43.8)	0.363
C-reactive protein >1.0 (mg/dL)	28 (45.9)	79 (62.7)	0.030
Blood urea nitrogen >20 (mg/dL)	5 (8.1)	26 (21.1)	0.025
Creatinine >1.0 (mg/dL)	9 (14.5)	21 (17.2)	0.640
Albumin <3.0 (g/dL)	1 (1.6)	3 (2.9)	0.605
Total bilirubin >2.0 (mg/dL)	8 (12.3)	9 (7.0)	0.222
Vital sign			
SBP <90 (mm Hg)	1 (1.5)	0 (0.0)	0.159
Pulse rate >100 bpm	2 (3.1)	16 (12.5)	0.033
Body temperature >37.8 (°C)	8 (12.3)	24 (18.8)	0.255
Time from ER arrival to surgery (min)	717.4±311.8	681.1±339.6	0.471
Length of hospital stay (day)	4.5±2.2	4.6±2.0	0.802
Mortality	1 (1.5)	0 (0.0)	0.159

SLRW, special law on restriction of working hours in training; ER, emergency room; SBP, systolic blood pressure.

Data are presented as interquartile range or n (%).