Study	Events Total		Proportion	95%-CI	Weight (common)	Weight (random)
Type = Multicentric						
García-Salido A	9 45		0.20	[0.11; 0.34]	0.9%	9.7%
Deep A	48 116		0.41	[0.33; 0.51]	3.7%	13.0%
Lima-Setta F	5 56 —	•	0.09	[0.04; 0.20]	0.6%	8.2%
Falah NU	1 10	+ 1	0.10	[0.01; 0.47]	0.1%	2.9%
Lorena Acevedo	23 78	¦ <u></u>	0.29	[0.20; 0.40]	2.1%	12.0%
Fatih Haslak	8 76 —		0.11	[0.05; 0.20]	0.9%	9.7%
Allison D. Miller	849 4470	+	0.19	[0.18; 0.20]	89.6%	14.6%
Common effect model	4851	\$	0.20	[0.19; 0.21]	98.0%	
Random effects model			0.20	[0.13; 0.29]		70.1%
Heterogeneity: $I^2 = 87\%$, τ	² = 0.3274, <i>p</i> < 0.01					
Type = Single						
Dhanalakshmi K	3 19 —		0.16	[0.05; 0.39]	0.3%	6.0%
Shobhavat L	8 21		- 0.38	[0.20; 0.60]	0.6%	8.5%
Alkan G	5 36 -			[0.06; 0.29]		8.0%
Gupta S	5 20			[0.11; 0.48]		7.5%
Common effect model	96			[0.16; 0.33]	2.0%	
Random effects model				[0.13; 0.36]		29.9%
Heterogeneity: $I^2 = 38\%$, τ	² = 0.1585, <i>p</i> = 0.19					
Common effect model	4947	 ↓ ↓	0.20	[0.19; 0.21]	100.0%	
Random effects model			0.21	[0.15; 0.28]		100.0%
Heterogeneity: $I^2 = 80\%$, τ	² = 0.2755, <i>p</i> < 0.01					
Test for subgroup difference	ses (fixed effect): $\chi_1^2 = 0$	0.168, 0.12 = 10(0.3 = 0.441) 0.5				
Test for subgroup difference		0				

Test for subgroup differences (random effects): $\chi_1^2 = 0.14$, df = 1 (p = 0.71)