

1
2 **Table S3 Plasmids used in this study**
3

Plasmid	Parent	Description	Source
pREP3x		<i>leu2</i> ⁺ P _{nmt1}	[1]
AB 553-1	pREP3x	<i>leu2</i> ⁺ with a NotI between PstI and P _{nmt1}	[2]
EMT 256-1	pREP3x	<i>leu2</i> ⁺ P _{trm6} <i>trm6</i> ⁺	This study
EMT 257-2	pREP3x	<i>leu2</i> ⁺ P _{trm61} <i>trm61</i> ⁺	This study
EMT 265-1	AB 553-1	<i>leu2</i> ⁺ <i>imt06</i> ⁺	This study
EMT 264-1	AB 553-1	<i>leu2</i> ⁺ <i>sup9</i> ⁺ - <i>imt07</i> ⁺	This study
EMT 327-1	AB 553-1	<i>leu2</i> ⁺ P _{tol1} <i>tol1</i> ⁺	This study
EMT 292		<i>ura4</i> ⁺ single integrating vector (SIV)	[3]
EMT 301-3	EMT 292	<i>ura4</i> ⁺ :: <i>dhp1</i> ⁺	This study
EMT 304-1	EMT 292	<i>ura4</i> ⁺ :: <i>imt06</i> ⁺	This study
EMT 324		URA3 P _{GAL} <i>TRM6</i> 2μ	[4]
EMP 881		URA3 2μ	Laboratory strain
EMT 306-1		<i>LEU2</i> 2μ <i>IMT1</i>	this study
EMP 824		<i>LEU2</i> 2μ	Laboratory strain
EMT 386		<i>leu2</i> ⁺ P _{cid14} <i>cid14</i> ⁺	this study

4
5 **References**
6

- 7 1. Forsburg SL. Comparison of *Schizosaccharomyces pombe* expression systems. Nucleic Acids
8 Res. 1993;21(12):2955-6.
- 9 2. De Zoysa T, Phizicky EM. Hypomodified tRNA in evolutionarily distant yeasts can trigger rapid
10 tRNA decay to activate the general amino acid control response, but with different consequences.
11 PLoS Genet. 2020;16(8):e1008893.
- 12 3. Vjestica A, Marek M, Nkosi PJ, Merlini L, Liu G, Berard M, et al. A toolbox of stable integration
13 vectors in the fission yeast *Schizosaccharomyces pombe*. J Cell Sci. 2020;133(1).
- 14 4. Gelperin DM, White MA, Wilkinson ML, Kon Y, Kung LA, Wise KJ, et al. Biochemical and
15 genetic analysis of the yeast proteome with a movable ORF collection. Genes Dev.
16 2005;19(23):2816-26.
- 17