

## Supplemental materials

### Genetic alterations of *Keap1* confers chemotherapeutic resistance through functional activation of *Nrf2* and Notch pathway in head and neck squamous cell carcinoma

Abu Shadat M Noman<sup>1</sup>, Khawlah Qassem<sup>2</sup>, Shafiqul Islam<sup>1</sup>, Rashed R Parag<sup>1</sup>, Mohammed Z Rahman<sup>3</sup>, Walid A Farhat<sup>4</sup>, Herman Yeger<sup>5</sup>, Abdelillah Aboussekhra<sup>2</sup>, Bedri Karakas<sup>2</sup>, Syed S Islam<sup>2,6\*</sup>

### Raw western blots for Figures 3/ 5/7

Original Raw Western blots for Figures 3B

Figure 3B Raw uncropped Western blots





Figure 3B Raw uncropped Western blots

Figure 5A Raw uncropped Nrf2 Western blot




Figure 5A Raw uncropped GAPDH Western blot




Figure 7B Raw uncropped Notch1 Western blot




Figure 7B Raw uncropped GAPDH Western blot



Figure 7B Raw uncropped Hes1 Western blot




Figure 7D Raw uncropped Notch1 Western blot




Figure 7D Raw uncropped GAPDH Western blot




Figure 7D Raw uncropped Hes1 Western blot




Figure 7F Raw uncropped Notch1 Western blot




Figure 7F Raw uncropped GAPDH Western blot




Figure 7I Raw uncropped Hes1 Western blot




Figure 7I Raw uncropped GAPDH Western blot

