Pinocembrin suppresses oxidized low-density lipoprotein-triggered NLRP3 inflammasome/GSDMD-mediated endothelial cell pyroptosis through an

Nrf2-dependent signaling pathway

Tong Wang¹§, Hua Tian^{2,3}§[∞], Tianqi Pan⁴, Shutong Yao⁴, Huayun Yu², Yumei Wu⁵ & Shijun Wang²[∞]

¹ College of nursing, Shandong University of Traditional Chinese Medicine, Jinan 250355, Shandong, China.

² College of Traditional Chinese Medicine, Shandong University of Traditional Chinese Medicine, Jinan 250355, Shandong, China.

³ Key Laboratory of Atherosclerosis in Universities of Shandong and Institute of Atherosclerosis, Shandong First Medical University & Shandong Academy of Medical Sciences, Taian 271000, Shandong, China.

⁴ College of Basic Medical Sciences, Shandong First Medical University & Shandong Academy of Medical Sciences, Taian 271000, Shandong, China.


⁵ Pharmacy Department, Dongying district hospital of Traditional Chinese Medicine,

Dongying 257000, Shandong, China.


§ Equal contribution to this study

 \square To whom correspondence should be addressed:

¹ Shijun Wang, Email: wsj@sdutcm.edu.cn, ² Hua Tian, Email:liutianfangyu @163.com, College of Traditional Chinese Medicine, Shandong University of Traditional Chinese Medicine, No.4655 Daxue Road, Changqing District, Jinan 250355, Shandong, China. Tel.: +86 0531 89628750; Fax: +86 0531 89628077.















Figure 4 c







Model Pin

