

Supplemental Table 1. Patients' characteristics based on the presence of NAFLD

	Patients without NAFLD (n = 92)	Patients with NAFLD (n = 183)	P value
Age, years	58 ± 11	52 ± 11	< 0.001
Gender, male/female %	73%/27%	74%/26%	0.79
Ethnicity			< 0.001
Caucasian, %	50%	44%	
Hispanic, %	23%	46%	
African-American, %	27%	8%	
Other, %	0%	2%	
Weight, kg	98 ± 18	100 ± 16	0.30
Body mass index, kg/m²	33.2 ± 5.3	34.3 ± 4.8	0.08
Presence of diabetes, %	64%	60%	0.52
Diabetes medications			
Metformin, %	49%	43%	0.34
Sulfonylureas, %	26%	25%	0.82
Insulin, %	15%	14%	0.73
Statin use, %	60%	57%	0.67
Blood pressure medication use, %	74%	68%	0.31
A1c, %	6.5 ± 1.2	6.5 ± 1.1	0.72
Fasting plasma glucose, mg/ml	125 ± 39	128 ± 34	0.49
Intrahepatic triglyceride content, %	3 ± 1	16 ± 9	< 0.001
Aspartate aminotransferase, U/L	23 ± 12	41 ± 26	< 0.001
Alanine aminotransferase, U/L	23 ± 18	58 ± 37	< 0.001