

Supplementary Table 1: CBCL score differences by pubertal stage, diagnosis, and sex

Internalizing behavior score difference: pre-puberty vs early puberty				
	ADHD	ASD	OCD	TD
Males	5.1 (-0.4, 10.7)	4.4 (-0.5, 9.3)	0.3 (-4.7, 5.3)	0.5 (-5.4, 6.4)
Difference from TD	4.6 (-3.5, 12.7)	3.9 (-3.8, 11.6)	-0.2 (-7.9, 7.5)	
Internalizing behavior score difference: early puberty vs late puberty				
	ADHD	ASD	OCD	TD
Males	-3.4 (-10.6, 3.8)	3.0 (-0.9, 6.8)	-1.5 (-6.4, 3.5)	6.0 (-0.5, 12.4)
Difference from TD	-9.4 (-19.0, 0.3)	-3.0 (-10.5, 4.5)	-7.4 (-15.5, 0.7)	
Females	0.9 (-3.3, 5.0)	1.0 (-1.5, 3.5)	-1.2 (-6.8, 4.4)	-2.7 (-7.4, 2.1)
Difference from TD	3.6 (-2.8, 9.9)	3.6 (-1.8, 9.0)	1.5 (-5.9, 8.9)	
Externalizing behavior score difference: pre-puberty vs early puberty				
	ADHD	ASD	OCD	TD
Males	4.1 (-1.1, 9.2)	0.4 (-5.0, 5.7)	1.0 (-4.2, 6.3)	4.1 (-1.4, 9.7)
Difference from TD	-0.1 (-7.6, 7.5)	-3.8 (-11.5, 3.9)	-3.1 (-10.7, 4.5)	
Females	0.1 (-2.9, 3.1)	-3.1 (-5.7, -0.5)	-3.5 (-9.6, 2.7)	-3.3 (-8.3, 1.8)
Difference from TD	3.3 (-2.6, 9.2)	0.2 (-5.5, 5.9)	-0.2 (-8.2, 7.8)	
Externalizing behavior score difference: early puberty vs late puberty				
	ADHD	ASD	OCD	TD
Males	-3.7 (-10.5, 3.1)	0.6 (-3.6, 4.8)	-3.4 (-9.1, 2.4)	0.0 (-5.4, 5.5)
Difference from TD	-3.7 (-12.4, 5.0)	0.6 (-6.3, 7.5)	-3.4 (-11.3, 4.5)	
Females	-5.3 (-10.2, -0.3)	-1.0 (-3.7, 1.7)	1.1 (-4.0, 6.3)	-0.9 (-5.3, 3.5)
Difference from TD	-4.4 (-11.0, 2.3)	-0.1 (-5.3, 5.0)	2.0 (-4.8, 8.8)	

ADHD: attention-deficit hyperactivity disorder; ASD: autism spectrum disorder; CBCL: Child Behavior Checklist; OCD: obsessive compulsive disorder; TD: typically developing.