

Supplemental Table 2. Association between EVALI knowledge and perceived impact of EVALI on e-cigarette interest, among smokers aware of EVALI (n=542)

	Perceived Impact of EVALI on Future E-cigarette Interest						pvalue
	Less interested in using e-cigarettes in future		More interested in using e-cigarettes in future		No effect on interest in using e-cigarettes in future		
	%	95% CI	%	95% CI	%	95% CI	
Believed main cause of EVALI was:							
E-cigarettes used to vape nicotine, like JUUL	58.4	(52.4-64.3)	1.0	(0.0-2.2)	40.6	(34.7-46.5)	<.0001
Vaping products used for vaping marijuana/THC	34.9	(25.8-44.0)	9.2	(2.2-16.3)	55.8	(46.1-65.6)	
Both were responsible	48.2	(41.4-54.9)	1.7	(0.0-3.4)	50.2	(43.4-56.9)	
I don't know	35.0	(27.3-42.8)	3.3	(0.3-6.2)	61.7	(53.7-69.7)	
Heard of Vitamin E Acetate as associated with EVALI							
Yes	48.5	(41.4-55.7)	3.9	(0.3-7.4)	47.6	(40.5-54.7)	0.85
No	46.9	(41.7-52.1)	2.3	(0.6-4.1)	50.8	(45.6-56.0)	
Not sure	45.8	(38.0-53.5)	3.1	(0.5-5.7)	51.1	(43.2-59.0)	

Note: Read as row percentages. P-values are based on bivariate Chi-square tests of association between EVALI knowledge and perceived impact variables