

S2 Table. Bivariate analysis: Predictors of 12-month mental health service utilisation.

Predictors	Any service use		Health service use		Education service use		Social care and criminal justice service use	
	OR (95%CI)	P	OR (95%CI)	p	OR (95%CI)	p	OR (95%CI)	p
<i>Sociodemographic characteristics</i>								
Female gender	0.82 0.57-1.17	0.271	0.90 0.62-1.31	0.583	0.47 0.18-1.19	0.111	1.34 0.53-3.40	0.535
Age	0.99 0.91-1.09	0.892	0.97 0.88-1.06	0.498	0.97 0.79-1.20	0.770	1.22 0.96-1.55	0.101
SEG	(Reference)							
High								
Low	1.16 0.81-1.67	0.408	1.00 0.67-1.45	0.979	2.39 0.88-6.47	0.087	3.32 0.96-11.51	0.059
Ethnicity	(Reference)							
White								
Non-White	1.00 0.70-1.41	0.981	0.84 0.58-1.23	0.371	0.45 0.18-1.16	0.098	2.64 0.98-7.06	0.054
<i>Guardian characteristics</i>								
Mother's Education	(Reference)							
No education/basic								
Secondary	1.27 0.88-1.85	0.205	1.40 0.94-2.01	0.100	1.86 0.744-6.9	0.190	0.45 0.15-1.29	0.136
University	1.34 0.76-2.39	0.315	1.52 0.84-2.78	0.168	1.21 0.25-5.88	0.815	0.76 0.17-3.48	0.728
Lower parental stigma-RIBS	1.08 1.02-1.15	0.010	1.07 1.00-1.14	0.038	1.20 0.99-1.46	0.060	1.04 0.89-1.21	0.619
<i>Clinical characteristics</i>								
<i>Psychiatric diagnosis trajectories</i>								
No diagnosis	(Reference)							
Incident	4.20 2.57-6.86	<0.001	4.49 2.67-7.55	<0.001	4.95 1.23-20.00	0.025	3.96 1.05-14.89	0.042
Remittent	2.76 1.66-4.58	<0.001	2.77 1.61-4.79	<0.001	6.21 1.73-22.21	0.005	2.45 0.58-10.32	0.223
Persistent	7.22 4.50-11.58	<0.001	7.85 4.77-12.90	<0.001	13.77 4.19-45.33	<0.001	7.18 2.16-23.85	0.001
SDQ impact score	1.49 1.36-1.62	<0.001	1.51 1.38-1.65	<0.001	1.55 1.35-1.78	<0.001	1.31 1.01-1.57	0.003
<i>Interview method</i>								
In-person	(Reference)							
Telephone	0.43 0.30-0.60	<0.001	0.40 0.28-0.58	<0.001	0.55 0.24-1.26	0.156	0.42 0.17-1.07	0.068