

Table S1 related to Fig.1. Analysis of clinical and microbiological characteristics of patients with septic shock

Pt#	SEX	AGE	SEPSIS SOURCE	Identified Pathogen	PCT
1	F	49	Intra-abdominal	<i>Staphylococcus aureus</i>	13.2
2	F	64	Intra-abdominal	<i>Staphylococcus aureus</i>	16.3
3	M	36	Intra-abdominal		3.3
4	F	51	Skin and Soft Tissue Infection		2.5
5	F	46	Intra-abdominal		17.0
6	M	66	Urinary tract infection	<i>Staphylococcus aureus</i>	6.8
7	M	72	Intra-abdominal		5.0
8	F	44	Intra-abdominal	<i>Staphylococcus aureus</i>	19.2
9	F	38	Intra-abdominal	<i>Escherichia coli</i>	14.9
10	M	62	Intra-abdominal	<i>Streptococcus pyogenes</i>	1.9
11	M	46	Urinary tract infection		3.6
12	M	71	Intra-abdominal		9.3
13	M	84	Pneumonia	<i>Streptococcus pneumoniae</i>	11.6
14	F	77	Intra-abdominal		4.2
15	F	52	Intra-abdominal		5.1
16	M	47	Intra-abdominal		6.6
17	M	60	Pneumonia	<i>Pseudomonas aeruginosa</i>	4.9
18	M	51	Intra-abdominal		2.6

Pt#	SEX	AGE	SEPSIS SOURCE	Identified Pathogen	PCT
19	F	56	Skin and Soft Tissue Infection	<i>Escherichia coli</i>	3.0
20	F	62	Intra-abdominal	<i>Staphylococcus aureus</i>	21.3
21	M	27	Intra-abdominal		19.2
22	F	42	Skin and Soft Tissue Infection		5.6
23	F	66	Intra-abdominal		7.3
24	M	66	Intra-abdominal	<i>Staphylococcus aureus</i>	6.8
25	M	49	Intra-abdominal		5.0
26	F	64	Intra-abdominal	<i>Staphylococcus aureus</i>	19.2
27	F	67	Pneumonia	<i>Escherichia coli</i>	14.9
28	M	61	Intra-abdominal		1.9
29	M	51	Intra-abdominal	<i>Staphylococcus aureus</i>	6.9
30	M	70	Intra-abdominal	<i>Streptococcus pyogenes</i>	11.9
31	M	80	Pneumonia	<i>Streptococcus pneumoniae</i>	13.2
34	M	33	Intra-abdominal		8.4
36	M	94	Intra-abdominal		11.2
