

Full wwPDB X-ray Structure Validation Report (i)

Sep 17, 2021 – 05;08 pm BST

PDB ID : 7PPY

Title : CARM1 in complex with EML709

Deposited on : 2021-09-15

Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity /: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.23.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.23.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\# \text{Entries}, \text{resolution range}(\mathring{ ext{A}}))$
R_{free}	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1 /	A	361	87%	8% 5%
1	В	361	2% 84%	10% 5%
1	C	361	90%	5% 5%
1	D	361	2%	7% • 6%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 22277 atoms, of which 10974 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.


Mol	Chain	Residues	Atoms	AltConf	Trace
1	A	342	Total C H N O S 5436 1772 2692 453 505 14	0	0
1	В	342	Total C H N O S 5436 1772 2692 453 505 14 0	0	0
1	С	343	Total C H N O S 5450 1776 2699 454 507 14 0	0	0
1	D	341	Total C H N O S 5415 1763 2683 452 503 14 0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	GLY		expression tag	UNP Q9WVG6
A	128	HIS	-	expression tag	UNP Q9WVG6
A	129	MET	_	expression tag	UNP Q9WVG6
В	127	GLY	_	expression tag	UNP Q9WVG6
В	128	HIS 🦰	_	expression tag	UNP Q9WVG6
В	129 /	MET	_	expression tag	UNP Q9WVG6
С	127 /	GLY	- /	expression tag	UNP Q9WVG6
С	128	HIS	-/	expression tag	UNP Q9WVG6
С	129	MET	/	expression tag	UNP Q9WVG6
D	/127	GLY	/ -	expression tag	UNP Q9WVG6
D	128	HIS	/ -	expression tag	UNP Q9WVG6
D	129	MET	_	expression tag	UNP Q9WVG6


• Molecule 2 is methyl 6-[4-[[{N}-[[(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methyl]carbamimidoyl]amino]butylcarbamoylamino]-4-oxidanyl-napht halene-2-carboxylate (three-letter code: 7ZH) (formula: C₂₈H₃₄N₁₀O₇) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms ZeroOcc AltConf
2	A	1	Total C H N O 0 0
2	В	1	Total C H N O 0 0
2	С	1	Total C H N O 0 0
2	D	1	Total C H N O 0 0

 \bullet Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$





Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
3	A	1	Total	С	Н	О	0	0
J 3	A	1	10	2	6	2	U	0 /
3	A	1	Total	С	Н	Ο	0	6
	Λ	1	10	2	6	2	U	/0
3	A	1	Total	С	Н	Ο	0	0
	71	1	10	2	6	2		
3	A	1	Total	С	Η	Ο	0 /	0
	71	1	10	2	6	2	0 /	
3	В	1	Total	С	Н	Ο	a/	0
	Б	1	10	2	6	2		A
3	В	1	Total	С	Η	Ο	0	0
	D	1	10	2	6	2		Ü
3	$^{\rm C}$	1	Total	С	Н	0 /	0.	0 /
	<u> </u>	-	10	2	6	2/		0 /
3	$^{\rm C}$	1	Total	С	Н	0	0	0/
	Ü	-	10	2	6/	2		/
3	$^{\rm C}$	1	Total	С	H	О	0	0
	Ü	-	10	2/	6	2		/ 0
3	$^{\rm C}$	1	Total	C	Н	O	0 /	0
	Ü	-	10	2	6	2		0
3	D	1	Total	С	H	0	0 /	0
		_	10	2	6	2		
3	D	1	Total	C	H	O	$\sqrt{0}$	0
	_	_	/ 10	2	6	2	/ "	Ü

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	31	Total O 31 31	0	0
4	В	24	Total O 24 24	0	0
4	C	31	Total O 31 31	0	0
4	D	18	Total O 18 18	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Histone-arginine methyltransferase CARM1







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	74.68Å 98.82Å 206.98Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.06 - 2.42	Depositor
resolution (A)	48.06 - 2.42	EDS
% Data completeness	98.1 (48.06-2.42)	Depositor
(in resolution range)	98.3 (48.06-2.42)	EDS /
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.47 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.205 , 0.235	Depositor
10, 10 free	0.205 , 0.235	DCC
R_{free} test set	2880 reflections (4.96%)	wwPDB-VP
Wilson B-factor (\mathring{A}^2)	44.2	Xtriage
Anisotropy	0.468	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38, 29.1	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	22277	wwPDB-VP
Average B, all atoms (A^2)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 35.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.5235e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, 7ZH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.51	0/2814	0.68	1/3813 (0.0%)	
1	В	0.48	0/2814	0.67	0/3813	
1	С	0.50	0/2821	0.68	0/3823	
1	D	0.47	0/2801	0.66/	0/3795	
All	All	0.49	0/11250	0.67	1/15244 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mo	Chain	Res	Type/	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	365	GLU	CA-CB-CG	5.08	124.57	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
/1	A	2744 /	2692	2692	16	0
1	В	2744	2692	2692	29	0
1	C	2751	2699	2699	14	0
1	D	2/732	2683	2683	17	0
2	A	45	34	0	0	0
2	В	45	34	0	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	С	45	34	0	0	0
2	D	45	34	0	0	0
3	A	16	24	24	0	0
3	В	8	12	12	1	0
3	С	16	24	24	0 /	0
3	D	8	12	12	0 /	0
4	A	31	0	0	4/	0
4	В	24	0	0	/5	0
4	С	31	0	0	7	0
4	D	18	0	0	6	0
All	All	11303	10974	10838	76	0 /

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A	• • •	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	$\text{overlap } (\mathring{\mathbf{A}})$
1:D:474:PHE:CD2	4:D:601:HOH:O	2.11	1.02
1:B:390:PHE:HE1	1:B:421:CYS:HG	0.99/	0.98
1:B:421:CYS:HB2	1:B:468:LEU:HD11	1.43	0.97
1:D:474:PHE:CE2	4:D:601:HOH:O	2.20	0.94
1:B:421:CYS:HB2	1:B:468:LEU:CD1	1.99	0.93
1:B:224:GLU:OE2	4:B:601:HOH:O	1.95	0.84
1:A:404:TRP:O	4:A:601:HOH:O	1.95	0.84
1:D:307:GLN:OE1/	4:D:601:HOH:O	2.01	0.77
1:B:417:TYR:O	4:B:602:HOH:O	2.03	0.74
1:B:338:GLN:OE1	4:B:603:HOH:O	2.06	0.73
1:A:250:GLU:OE2	4:A:602:HOH:O	2.05	0.73
1:D:175:ARG:NH1	4:D:602:HOH:O	2.22	0.72
1:B:421:CYS:CB	1:B:468:LEU:CD1	2.68	0.72
1:B:337:ARG:HA	1:B:468:LEU:HD23	1.73	0.71
1:C:271:GLU:OE1	4:C:601:HOH:O	2.11	0.67
1:B:421:CYS:SG	1:B:453:ILE:CD1	2.82	0.67
1:B:421:CYS:SG	1:B:453:ILE:HD13	2.36	0.66
1:C:267:GLU:OE2	4:C:602:HOH:O	2.14	0.65
1:C:283:SER:HA	/4:C:604:HOH:O	2.00	0.62
1:B:421:CYS:CB	1:B:468:LEU:HD13	2.31	0.60
1:A:166:ASP:OD1	1:A:168:VAL:HG22	2.02	0.60
1:D:307:GLN:CD	4:D:601:HOH:O	2.38	0.60
1:D:307:GLN:NE2	4:D:601:HOH:O	2.36	0.59



Continued from previous page...

Continued from previous page								
Atom-1	Atom-2	Interatomic	Clash					
		$\operatorname{distance}\left(\mathrm{\AA}\right)$	overlap (Å)					
1:B:390:PHE:HE1	1:B:421:CYS:SG	2.15	0.58					
1:C:283:SER:N	4:C:604:HOH:O	2.37	0.58					
1:C:282:PRO:C	4:C:604:HOH:O	2.41	0.57					
1:B:314:TRP:O	1:B:325:SER:HA	2.06	0.56					
1:C:283:SER:CA	4:C:604:HOH:O	2.53	0.56					
1:D:314:TRP:O	1:D:325:SER:HA	2.06	0.56					
1:D:315:TYR:CE1	1:D:328:ARG:HG3	2.42	0.54					
1:B:421:CYS:HB3	1:B:468:LEU:HD13	1.90	0.54					
1:C:424:GLN:NE2	4:C:605:HOH:O	2.44	0.51					
1:D:282:PRO:O	1:D:283:SER:HB2	2.12	0.50					
1:C:379:MET:CE	1:C:385:VAL:HG21	2.41	0.50					
1:B:175:ARG:HD2	4:B:608:HOH:O	2.13	0.48					
1:A:376:LYS:HG3	1:A:434:THR:CG2	2.44	0.47					
1:D:143:GLU:HG3	1:D:445:LYS:HB3	1.96	0.47					
1:D:268:ARG:HD3	1:D:447:GLN:HA	1.97	0.47					
1:B:467:LEU:O	1:B:468:LEU:HD 2 3	2.15	0.47					
1:B:268:ARG:HD3	1:B:447:GLN:HA	1.98	0.46					
1:A:268:ARG:HD3	1:A:447:GLN:HA	1.98	0.46					
1:B:141:ARG:HG3	1:B:142:THR:HG23	1.99	0.45					
1:B:188:ILE:HD12	1:B:210:LYS:HB3	1.99	0.45					
1:B:437:GLY:HA3	1:B:454:VAL:O	2.18	0.44					
1:D:437:GLY:HA3	1:D:454:VAL:O	2,18	0.44					
1:A:255:ILE:HB	1:A:286:MET:HE3	1.98	0.43					
1:A:471:LYS:HA	4:A:613:HOH:O	2.17	0.43					
1:B:268:ARG:CZ	1:B:443:ALA:HB1	2.48	0.43					
1:A:188:ILE:HG22	1:A:252:VAL:HG12	2.01	0.43					
1:B:282:PRO:O	1:B:283:SER:CB	2.65	0.43					
1:B:282:PRO:O	1:B:283:SER:HB3	2.18	0.43					
1:B:441:LEU:HB3	1:B:449:TYR:CD2	2.54	0.43					
1:C:268:ARG:HD3	1:C:447:GLN:HA	1.99	0.43					
1:C:437:GLY:HA3	1:C:454:VAL:O	2.19	0.43					
1:B:370:ARG:HG3	1:B:442:ILE:CD1	2.48	0.43					
1:A:168:VAL:HG22	4:A:608:HOH:O	2.19	0.43					
1:A:437:GLY:HA3	1:A:454:VAL:O	2.19	0.42					
1:B:315:TYR:CE1	1:B:328:ARG:HB2	2.54	0.42					
1:A:299:THR:HG23	1;A:384:LEU:O	2.19	0.42					
1:D:188:ILE:HD12	1:D:210:LYS:HB3	2.02	0.42					
1:D:263:MET:CE	1:D:419:VAL:HG11	2.49	0.42					
1:D:469:ASP:HB3	1:D:471:LYS:HG2	2.01	0.42					
1:B:193:GLY:HA3	2:B:501:7ZH:O7	2.19	0.42					
1:C:453:ILE:O	1:C:465:SER:HA	2.20	0.42					
		C ti	<u> </u>					



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} ({\rm \AA}) \end{array}$	$egin{array}{c} ext{Clash} \ ext{overlap } (ext{Å}) \end{array}$
1:D:268:ARG:CZ	1:D:443:ALA:HB1	2.50	0.42
1:C:471:LYS:HD2	1:C:471:LYS:HA	1.84	0.42
1:D:364:LYS:HD3	1:D:366:GLY:H	1.85	0.41
3:B:502:EDO:O2	4:B:604:HOH:O	2.20	0.41
1:A:186:ASP:HA	1:A:209:ARG:HB2	2.02	0.41
1:A:184:PHE:O	1:A:208:ALA:HA	2.21	0.41
1:B:423:PHE:HA	1:B:466:ASN:ND2	2.36	0.41
1:A:145:SER:HA	1:B:145:SER:CB	2.51	0.41
1:C:270:LEU:HD22	1:C:360:PHE:CE1	2,56	0.41
1:A:268:ARG:CZ	1:A:443:ALA:HB1	2.50	0.40
1:A:270:LEU:HD22	1:A:360:PHE:CE1	2.57	0.40
1:C:268:ARG:CZ	1:C:443:ALA:HB1	2.50	0,40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	${f ntiles}$
1	Α	340/361 (94%)	329 (97%)	11 (3%)	0	100	100
1	В	340/361 (94%)	327 (96%)	13 (4%)	0	100	100
1	C	341/361 (94%)	327 (96%)	14 (4%)	0	100	100
1	D	339/361 (94%)	329 (97%)	10 (3%)	0	100	100
All	All	1360/1444 (94%)	1312 (96%)	48 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$298/313\ (95\%)$	295 (99%)	3 (1%)	76 87
1	В	$298/313\ (95\%)$	293 (98%)	5 (2%)	60 77
1	С	299/313~(96%)	299 (100%)	0 /	100 100
1	D	$297/313\ (95\%)$	290 (98%)	7 (2%)	49 67
All	All	$1192/1252 \ (95\%)$	1177 (99%)	15 (1%)	69 83

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	136	SER
1	A	144	GLU
1	A	415	HIS
1	В	221	GLN
1	В	308	PHE
1	В	415	HIS
1	В	466	ASN /
1	В	469	ASP
1	D	136	SER
1	D	209	ARG
1	D	308	PHE
1	D	328	ARG
1	D	333/	ASP
1	D	415	HIS
1	D	471	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

								/		
Mal	Т	Chain	Des	Bond lengths			ths	Bond angles		
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	$7\mathrm{ZH}$	A	501	_	44,49,49	0.46	0/	54,69,69	0.62	1 (1%)
3	EDO	A	502	- /	3,3,3	0.07	/0	2,2,2	0.24	0
3	EDO	D	503	-/	3,3,3	0.16	0	2,2,2	0.42	0
3	EDO	С	503	+	3,3,3	0.10	0	2,2,2	0.27	0
3	EDO	С	505	/-	3,3,3	0.07	0	2,2,2	0.17	0
3	EDO	A	504	-	3,3,3	0.06	0	2,2,2	0.14	0
3	EDO	С	504		3,3,3	0.06	0	2,2,2	0.17	0
3	EDO	A	503		3,3,3	0.07	0	2,2,2	0.07	0
2	$7\mathrm{ZH}$	C	$\sqrt{501}$	_	44,49,49	0.44	0	54,69,69	0.65	1 (1%)
2	7ZH	В	501	<u> </u>	44,49,49	0.46	0	54,69,69	0.75	2 (3%)
3	EDO	В/	503		3,3,3	0.06	0	2,2,2	0.23	0
3	EDO	Ď	502	77	3,3,3	0.14	0	2,2,2	0.25	0
3	EDO	/ C	502) -	/3,3,3	0.05	0	2,2,2	0.17	0
2	7ZH	D	501		44,49,49	0.48	0	54,69,69	0.68	1 (1%)
3	EDO	В	502	- /	3,3,3	0.09	0	2,2,2	0.16	0
3	EDO	A	505	-/	3,3,3	0.14	0	2,2,2	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol T	ype	Chain	Res	Link	Chirals	Torsions	Rings
2	7ZH	A	501	-	-	10/24/44/44	0/5/5/5
3 E	EDO	/ A	502	_	_	1/1/1/1	-
3 E	EDO	/ D	503	-	=	1/1/1/1	_



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	С	503	-	-	1/1/1/1	-
3	EDO	С	505	-	-	1/1/1/1	/ -
3	EDO	A	504	_	-	1/1/1/1	-
3	EDO	С	504	_	-	1/1/1/1	-
3	EDO	A	503	_	-	1/1/1/1	-22
2	7ZH	С	501	_	-	10/24/44/44	0/5/5/5
2	7ZH	В	501	-	-	8/24/44/44	0/5/5/5
3	EDO	В	503	-	-	1/1/1/1	-
3	EDO	D	502	_	_	1/1/1/1	-
3	EDO	С	502	_	-	0/1/1/1	-
2	7ZH	D	501	_	-	5/24/44/44	0/5/5/5
3	EDO	В	502	_	- /	1/1/1/1	- /
3	EDO	A	505	-	- /	1/1/1/1	7

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$ \operatorname{Ideal}({}^o) $
2	В	501	7ZH	N3-C17-N5	2.76	125.44	120.26
2	В	501	7ZH	C25-C26-N8	2.41	/124.02	120.35
2	D	501	7ZH /	C25-C26-N8	2.36	123.93	120.35
2	С	501	7ZH	C25-C26-N8	2.22	123.73	120.35
2	A	501	7ZH	C25-C26-N8	2.21	123.71	120.35

There are no chirality outliers.

All (44) torsion outliers are listed below:

Mol	Chain/	Res	Type	Atoms
2	A /	501	7ZH	N4-C19-C20-C21
2	Ą	501	7ZH	N4-C19-C20-O7
2	/B	501	$7\mathrm{ZH}$	C1-C11-O3-C18
2	/ B	501	7ZH	N3-C17-N4-C19
2 /	В	501	$7\mathrm{ZH}$ $/$	N5-C17-N4-C19
2/	D	501	7ZH	N4-C19-C20-C21
/2	D	501	$7\mathrm{ZH}$	N4-C19-C20-O7
/2	A	501	7ZH	O4-C11-O3-C18
2	A	501	/ 7ZH	C1-C11-O3-C18
2	D	501/	$7\mathrm{ZH}$	N2-C13-C14-C15
2	В	501	7ZH	O4-C11-O3-C18
2	В	/ 501	7ZH	N2-C13-C14-C15



Continued from previous page...

Mol	Chain	Res	$\overline{\text{Type}}$	Atoms
2	С	501	7ZH	N2-C13-C14-C15
2	В	501	7ZH	C14-C15-C16-N3
2	A	501	7ZH	C2-C1-C11-O3
2	С	501	7ZH	C9-C8-N1-C12
2	A	501	7ZH	C6-C1-C11-O3
2	С	501	7ZH	C7-C8-N1-C12
2	С	501	7ZH	C14-C15-C16-N3
2	В	501	7ZH	C13-C14-C15-C16
3	A	504	EDO	O1-C1-C2-O2
3	A	505	EDO	O1-C1-C2-O2
3	В	502	EDO	O1-C1-C2-O2
3	A	502	EDO	O1-C1-C2-O2
3	С	503	EDO	O1-C1-C2-O2
3	С	504	EDO	O1-C1-C2-O2
3	С	505	EDO	O1-C1-C2-O2
2	A	501	7ZH	C2-C1-C11-O4
3	A	503	EDO	O1-C1-C2-O2
3	D	502	EDO	O1-C1-C2-O2
2	D	501	7ZH	N5-C17-N4-C19
2	A	501	7ZH	C6-C1-C11-O4
2	D	501	7ZH	N3-C17-N4-C19
2	С	501	7ZH	C2-C1-C11-O3
2	С	501	7ZH/	C6-C1-C11-O3
3	В	503	EDO	O1-C1-C2-O2
3	D	503	EDO	O1-C1-C2-O2
2	С	501	$/7\mathrm{ZH}$	C13-C14-C15-C16
2	В	501/	7ZH	C15-C16-N3-C17
2	С	501	7ZH	C2-C1-C11-O4
2	A	501	7ZH	C9-C8-N1-C12
2	С	501	7ZH	C6-C1-C11-O4
2	С /	501	7ZH	C14-C13-N2-C12
2	A/	501	7ZH	C7-C8-N1-C12

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
 2	В	501	/7ZH	1	0
3	В	502	EDO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In



addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues 🕦

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	342/361 (94%)	-0.06	3 (0%) 84 82	32, 45, 62, 69	0
1	В	342/361 (94%)	0.04	8 (2%) 60 57	33, 49, 66, 80	0
1	С	343/361 (95%)	0.02	7 (2%) 65 62	34, 47, 63, 76	0
1	D	341/361 (94%)	-0.00	6 (1%) 68 65	36, 48, 66, 80	0
All	All	1368/1444 (94%)	-0.00	24 (1%) 68 65	32, 47, 65, 80	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	178	LEU /	3.6
1	D	380	LEU	3.5
1	В	477	TYR	3.3
1	D	167	TYR	3.0
1	D	347	ARG	2.9
1	В	144/	GLU	2.8
1	В	167	TYR	2.7
1	С	276	ALA	2.7
1	С /	144	GLU	2.7
1	В	136	SER	2.7/
1	Ç	415	HIS	2,6
1	/C	477	TYR	2.6
1	/ B	347	ARG	2.6
1 /	A	477	TYR /	2.6
1/	В	140	GLU	2.3
/1	В	145	SER	2.3
/ 1	A	140	GLU	2.3
1	D	328	ARG	2.3
1	D	140/	GLU	2.2
1_	A	138	PHE	2.2
1	C	/136	SER	2.2



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	136	SER	2.1
1	В	381	HIS	2.1
1	С	204	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q < 0.9
3	EDO	С	504	4/4	0.74	0.29	63,76,84,84	0
3	EDO	D	502	4/4	0.76	0.38	57,69,74,74	0
3	EDO	С	$\sqrt{505}$	4/4	0.81	0.17	57,69,73,73	0
3	EDO	С /	503	4/4	0.81	0.22	61,73,81,82	0
3	EDO	A/	505	4/4	0,82	0.18	61,73,78,78	0
3	EDO	A	504	4/4	0.84	0.24	59,72,74,78	0
3	EDO	D	503	4/4	0.84	0.10	51,62,65,65	0
3	EDO	C	502	4/4	0.86	0.29	53,64,72,72	0
3	EDO	A	503	4/4/	0.86	0.18	61,73,74,80	0
3	EDO	A	502	4/4	0.87	0.34	58,71,75,78	0
3	EDO	В	502	/4/4	0.88	0.27	$56,\!68,\!72,\!72$	0
3	ÉDO	В	503	4/4	0.92	0.17	51,62,64,67	0
2 /	7ZH	A	501/	45/45	0.93	0.15	43,63,91,94	0
2/	7ZH	В	50/1	45/45	0.93	0.19	44,63,90,93	0
/2	7ZH	C	501	45/45	0.94	0.17	48,68,99,103	0
2	7ZH	D	/501	45/45	0.94	0.18	44,59,86,96	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 7ZH A 501: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive) Electron density around 7ZH B 501: $2\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

